

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
7 April 2005 (07.04.2005)

PCT

(10) International Publication Number
WO 2005/030124 A2

(51) International Patent Classification⁷: **A61K**

(21) International Application Number:
PCT/US2004/029390

(22) International Filing Date:
9 September 2004 (09.09.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/502,163 10 September 2003 (10.09.2003) US

(71) Applicants (for all designated States except US):
WARNER-LAMBERT COMPANY LLC [US/US];
201 Tabor Road, Morris Plains, New Jersey 07950 (US).
ABGENIX, INC. [US/US]; 6701 Kaiser Drive, Fremont,
California 94555 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BEDIAN, Vahe**
[US/US]; 3 Frog Hollow Road, East Lyme, Connecticut
06333 (US). **DEVALARAJA, Madhav Narasimha**
[IN/US]; 690 Trego Circle, Ann Arbor, Michigan 48103
(US). **LOW, Joseph Edwin** [US/US]; 4259 Fairway
Drive, Ann Arbor, Michigan 48103 (US). **MOBLEY,**

James Leslie [US/US]; 10195 Covington, Brighton,
Michigan 48114 (US). **KELLERMANN, Sird-Aimee**
[DE/US]; 1000 Alamanor Avenue, Menlo Park, California
94025 (US). **FOLTZ, Ian** [CA/CA]; 2108 Knightswood
Place, Burnaby, British Columbia V5A4B9 (CA).
HAAK-FREDSCHO, Mary [US/US]; 8472 Willow
Place, Newark, California 94560 (US).


(74) Agents: **HALEY, James E., Jr.** et al.; c/o FISH & NEAVE,
1251 Avenue of the Americas, New York, NY 10020 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PI, PT, RO, RU, SC, SD, SF, SG, SK, SI, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,

[Continued on next page]

(54) Title: ANTIBODIES TO M-CSF



(57) Abstract: The present invention relates to antibodies and antigen-binding portions thereof that specifically bind to a M-CSF, preferably human M-CSF, and that function to inhibit a M-CSF. The invention also relates to human anti-M-CSF antibodies and antigen-binding portions thereof. The invention also relates to antibodies that are chimeric, bispecific, derivatized, single chain antibodies or portions of fusion proteins. The invention also relates to isolated heavy and light chain immunoglobulins derived from human anti-M-CSF antibodies and nucleic acid molecules encoding such immunoglobulins. The present invention also relates to methods of making human anti-M-CSF antibodies, compositions comprising these antibodies and methods of using the antibodies and compositions for diagnosis and treatment. The invention also provides gene therapy methods using nucleic acid molecules encoding the heavy and/or light immunoglobulin molecules that comprise the human anti-M-CSF antibodies. The invention also relates to transgenic animals and transgenic plants comprising nucleic acid molecules of the present invention.



FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI,
SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Published:

- *without international search report and to be republished upon receipt of that report*

ANTIBODIES TO M-CSF

[0001] This application claims the benefit of United States Provisional Application 60/502,163, filed September 10, 2003.

Background of the Invention

- 5 [0002] Macrophage colony stimulating factor (M-CSF) is a member of the family of proteins referred to as colony stimulating factors (CSFs). M-CSF is a secreted or a cell surface glycoprotein comprised of two subunits that are joined by a disulfide bond with a total molecular mass varying from 40 to 90 kD ((Stanley E.R., *et al.*, *Mol. Reprod. Dev.*, 46:4-10 (1997)). Similar to other CSFs, M-CSF is
- 10 produced by macrophages, monocytes, and human joint tissue cells, such as chondrocytes and synovial fibroblasts, in response to proteins such as interleukin-1 or tumor necrosis factor-alpha. M-CSF stimulates the formation of macrophage colonies from pluripotent hematopoietic progenitor stem cells (Stanley E.R., *et al.*, *Mol. Reprod. Dev.*, 46:4-10 (1997)).
- 15 [0003] M-CSF typically bind to its receptor, *c-fms*, in order to exert a biological effect. *c-fms* contains five extracellular Ig domains, one transmembrane domain, and an intracellular domain with two kinase domains. Upon M-CSF binding to *c-fms*, the receptor homo-dimerizes and initiates a cascade of signal transduction pathways including the JAK/STAT, PI3K, and ERK pathways.
- 20 [0004] M-CSF is an important regulator of the function, activation, and survival of monocytes/macrophages. A number of animal models have confirmed the role of M-CSF in various diseases, including rheumatoid arthritis (RA) and cancer. Macrophages comprise key effector cells in RA. The degree of synovial

- macrophage infiltration in RA has been shown to closely correlate with the extent of underlying joint destruction. M-CSF, endogenously produced in the rheumatoid joint by monocytes/macrophages, fibroblasts, and endothelial cells, acts on cells of the monocyte/macrophage lineage to promote their survival and differentiation into
- 5 bone destroying osteoclasts, and enhance pro-inflammatory cellular functions such as cytotoxicity, superoxide production, phagocytosis, chemotaxis and secondary cytokine production. For example, treatment with M-CSF in the rat streptococcus agalactiae sonicate-induced experimental arthritis model lead to enhanced pathology (Abd, A.H., *et al.*, *Lymphokine Cytokine Res.* 10:43-50 (1991)).
- 10 Similarly, subcutaneous injections of M-CSF in a murine model of collagen-induced arthritis (CIA), which is a model for RA, resulted in a significant exacerbation of the RA disease symptoms (Campbell I.K., *et al.*, *J. Leuk. Biol.* 68:144-150 (2000)). Furthermore, MRL/lpr mice that are highly susceptible to RA and other autoimmune diseases have elevated basal M-CSF serum concentrations
- 15 (Yui M.A., *et al.*, *Am. J. Pathol.* 139:255-261 (1991)). The requirement for endogenous M-CSF in maintaining CIA was demonstrated by a significant reduction in the severity of established disease by M-CSF neutralizing mouse monoclonal antibody (Campbell I.K., *et al.*, *J. Leuk. Biol.* 68:144-150 (2000)).
- [0005] With respect to cancer, inhibition of colony stimulating factors by
- 20 antisense oligonucleotides suppresses tumor growth in embryonic and colon tumor xenografts in mice by decelerating macrophage-mediated ECM breakdown (Seyedhossein, A., *et al.*, *Cancer Research*, 62:5317-5324 (2002)).
- [0006] M-CSF binding to *c-fms* and its subsequent activation of monocyte/macrophages is important in a number of disease states. In addition to
- 25 RA and cancer, the other examples of M-CSF-related disease states include osteoporosis, destructive arthritis, atherogenesis, glomerulonephritis, Kawasaki disease, and HIV-1 infection, in which monocytes/macrophages and related cell types play a role. For instance, osteoclasts are similar to macrophages and are regulated in part by M-CSF. Growth and differentiation signals induced by
- 30 M-CSF in the initial stages of osteoclast maturation are essential for their subsequent osteoclastic activity in bone.

- [0007] Osteoclast mediated bone loss, in the form of both focal bone erosions and more diffuse juxta-articular osteoporosis, is a major unsolved problem in RA. The consequences of this bone loss include joint deformities, functional disability, increased risk of bone fractures and increased mortality. M-CSF is uniquely
- 5 essential for osteoclastogenesis and experimental blockade of this cytokine in animal models of arthritis successfully abrogates joint destruction. Similar destructive pathways are known to operate in other forms of destructive arthritis such as psoriatic arthritis, and could represent venues for similar intervention.
- [0008] Postmenopausal bone loss results from defective bone remodeling
- 10 secondary to an uncoupling of bone formation from exuberant osteoclast mediated bone resorption as a consequence of estrogen deficiency. *In-vivo* neutralization of M-CSF using a blocking antibody has been shown in mice to completely prevent the rise in osteoclast numbers, the increase in bone resorption and the resulting bone loss induced by ovariectomy.
- 15 [0009] Several lines of evidence point to a central role for M-CSF in atherogenesis, and in proliferative intimal hyperplasia after mechanical trauma to the arterial wall. All the major cell types in atherosclerotic lesions have been shown to express M-CSF, and this is further up-regulated by exposure to oxidized lipoprotein. Blockade of M-CSF signaling with a neutralizing *c-fms* antibody
- 20 reduces the accumulation of macrophage-derived foam cells in the aortic root of apolipoprotein E deficient mice maintained on a high fat diet.
- [0010] In both experimental and human glomerulonephritis, glomerular M-CSF expression has been found to co-localize with local macrophage accumulation, activation and proliferation and correlate with the extent of glomerular injury and
- 25 proteinuria. Blockade of M-CSF signaling via an antibody directed against its receptor *c-fms* significantly down-regulates local macrophage accumulation in mice during the renal inflammatory response induced by experimental unilateral ureteric obstruction.
- [0011] Kawasaki disease (KD) is an acute, febrile, pediatric vasculitis of
- 30 unknown cause. Its most common and serious complications involve the coronary vasculature in the form of aneurismal dilatation. Serum M-CSF levels are significantly elevated in acute phase Kawasaki's disease, and normalize following

treatment with intravenous immunoglobulin. Giant cell arthritis (GCA) is an inflammatory vasculopathy mainly occurring in the elderly in which T cells and macrophages infiltrate the walls of medium and large arteries leading to clinical consequences that include blindness and stroke secondary to arterial occlusion. The

- 5 active involvement of macrophages in GCA is evidenced by the presence of elevated levels of macrophage derived inflammatory mediators within vascular lesions.

- [0012] M-CSF has been reported to render human monocyte derived macrophages more susceptible to HIV-1 infection *in vitro*. In a recent study,
- 10 M-CSF increased the frequency with which monocyte-derived macrophages became infected, the amount of HIV mRNA expressed per infected cell, and the level of proviral DNA expressed per infected culture.

[0013] Given the role of M-CSF in various diseases, a method for inhibiting M-CSF activity is desirable.

- 15 [0014] There is a critical need for therapeutic anti-M-CSF antibodies.

SUMMARY OF THE INVENTION

[0015] The present invention provides isolated human antibodies or antigen-binding portions thereof that specifically bind human M-CSF and acts as a M-CSF antagonist and compositions comprising said antibody or portion.

- 5 [0016] The invention also provides for compositions comprising the heavy and/or light chain, the variable regions thereof, or antigen-binding portions thereof an anti-M-CSF antibody, or nucleic acid molecules encoding an antibody, antibody chain or variable region thereof the invention effective in such treatment and a pharmaceutically acceptable carrier. In certain embodiments, the compositions
- 10 may further comprise another component, such as a therapeutic agent or a diagnostic agent. Diagnostic and therapeutic methods are also provided by the invention. In certain embodiments, the compositions are used in a therapeutically effective amount necessary to treat or prevent a particular disease or condition.

- [0017] The invention also provides methods for treating or preventing a variety
- 15 of diseases and conditions such as, but not limited to, inflammation, cancer, atherogenesis, neurological disorders and cardiac disorders with an effective amount of an anti-M-CSF antibody of the invention, or antigen binding portion thereof, nucleic acids encoding said antibody, or heavy and/or light chain, the variable regions, or antigen-binding portions thereof.

- 20 [0018] The invention provides isolated cell lines, such as a hybridomas, that produce anti-M-CSF antibodies or antigen-binding portions thereof.

[0019] The invention also provides nucleic acid molecules encoding the heavy and/or light chains of anti-M-CSF antibodies, the variable regions thereof, or the antigen-binding portions thereof.

- 25 [0020] The invention provides vectors and host cells comprising the nucleic acid molecules, as well as methods of recombinantly producing the polypeptides encoded by the nucleic acid molecules.

- [0021] Non-human transgenic animals or plants that express the heavy and/or light chains, or antigen-binding portions thereof, of anti-M-CSF antibodies are also
- 30 provided.

BRIEF DESCRIPTION OF THE DRAWINGS

- [0022] Figures 1A and 1B are graphs illustrating that the anti-M-CSF antibodies resulted in a dose-related decrease in total monocyte counts in male and female monkeys over time. The monocyte counts were determined by light scatter using an Abbott Diagnostics Inc. Cell Dyn system. Monocyte counts were monitored from 24 hours through 3 weeks after administration of vehicle or antibody 8.10.3 at 0, 0.1, 1 or 5 mg/kg in a dose volume of 3.79 mL/kg over an approximately 5 minute period.
- [0023] Figure 1A male monkeys.
- 10 [0024] Figure 1B female monkeys.
- [0025] Figures 2A and 2B are graphs illustrating that anti-M-CSF treatment resulted in a reduction in the percentage of CD14+CD16+ monocytes, in male and female monkeys. 0-21 days after administration of vehicle or antibody 8.10.3 at 0, 0.1, 1 or 5 mg/kg in a dose volume of 3.79 mL/kg over an approximately 5 minute period. For each monkey tested, the percentage of monocytes within the CD14+CD16+ subset was determined after each blood draw, on days 1, 3, 7, 14 and 21 after 8.10.3 injection.
- [0026] Figure 2A male monkeys.
- [0027] Figure 2B female monkeys.
- 20 [0028] Figures 3A and 3B are graphs illustrating that anti-M-CSF treatment resulted in a decrease in the percentage change of total monocytes at all doses of antibody 8.10.3F and antibody 9.14.4I as compared to pre-test levels of monocytes.
- [0029] Figure 3A shows data collected from experiments using antibody 8.10.3F.
- [0030] Figure 3B shows data collected from experiments using antibody 9.14.4I.
- 25 [0031] Figure 4 is a sequence alignment of the predicted amino acid sequences of light and heavy chain variable regions from twenty-six anti-M-CSF antibodies compared with the germline amino acid sequences of the corresponding variable region genes. Differences between the antibody sequences and the germline gene sequences are indicated in bold-faced type. Dashes represent no change from germline. The underlined sequences in each alignment represent, from left to right, the FR1, CDR1, FR2, CDR2, FR3, CDR3 AND FR4 sequences.
- 30

- [0032] Figure 4A shows an alignment of the predicted amino acid sequence of the light chain variable region for antibody 252 (residues 21-127 of SEQ ID NO: 4) to the germline V_κO12, J_κ3 sequence (SEQ ID NO: 103).
- [0033] Figure 4B shows an alignment of the predicted amino acid sequence of the light chain variable region for antibody 88 (residues 21-127 of SEQ ID NO: 8) to the germline V_κO12, J_κ3 sequence (SEQ ID NO: 103).
- [0034] Figure 4C shows an alignment of the predicted amino acid sequence of the light chain variable region for antibody 100 (residues 21-127 of SEQ ID NO: 12) to the germline V_κL2, J_κ3 sequence (SEQ ID NO: 107).
- [0035] Figure 4D shows an alignment of the predicted amino acid sequence of the light chain variable region for antibody 3.8.3 (residues 23-130 of SEQ ID NO: 16) to the germline V_κL5, J_κ3 sequence (SEQ ID NO: 109).
- [0036] Figure 4E shows an alignment of the predicted amino acid sequence of the light chain variable region for antibody 2.7.3 (residues 23-130 of SEQ ID NO: 20) to the germline V_κL5, J_κ4 sequence (SEQ ID NO: 117).
- [0037] Figure 4F shows an alignment of the predicted amino acid sequence of the light chain variable region for antibody 1.120.1 (residues 21-134 of SEQ ID NO: 24) to the germline V_κB3, J_κ1 sequence (SEQ ID NO: 112).
- [0038] Figure 4G shows an alignment of the predicted amino acid sequence of the heavy chain variable region for antibody 252 (residues 20-136 of SEQ ID NO: 2) to the germline V_H3-11, D_H7-27 J_H6 sequence (SEQ ID NO: 106).
- [0039] Figure 4H shows an alignment of the predicted amino acid sequence of the heavy chain variable region for antibody 88 (residues 20-138 of SEQ ID NO: 6) to the germline V_H3-7, D_H6-13, J_H4 sequence (SEQ ID NO: 105).
- [0040] Figure 4I shows the alignment of the predicted amino acid sequence of the heavy chain variable region for antibody 100 (residues 20-141 of SEQ ID NO: 10) to the germline V_H3-23, D_H1-26, J_H4 sequence (SEQ ID NO: 104).
- [0041] Figure 4J shows an alignment of the predicted amino acid sequence of the heavy chain variable region for antibody 3.8.3 (residues 20-135 of SEQ ID NO: 14) to the germline V_H3-11, D_H7-27, J_H4 sequence (SEQ ID NO: 108).

- [0042] Figure 4K shows an alignment of the predicted amino acid sequence of the heavy chain variable region for antibody 2.7.3 (residues 20-137 of SEQ ID NO: 18) to the germline V_H3-33, D_H1-26, J_H4 sequence (SEQ ID NO: 110).
- 5 [0043] Figure 4L shows an alignment of the predicted amino acid sequence of the heavy chain variable region for antibody 1.120.1 (residues 20-139 of SEQ ID NO: 22) to the germline V_H1-18, D_H4-23, J_H4 sequence (SEQ ID NO: 111).
- [0044] Figure 4M shows an alignment of the predicted amino acid sequence of the light chain variable region for antibody 8.10.3 (residues 21-129 of SEQ ID NO: 44) to the germline V_κA27, J_κ4 sequence (SEQ ID NO: 114).
- 10 [0045] Figure 4N shows an alignment of the predicted amino acid sequence of the heavy chain variable region for antibody 8.10.3 (residues 20-141 of SEQ ID NO: 30) to the germline V_H3-48, D_H1-26, J_H4b sequence (SEQ ID NO: 113).
- [0046] Figure 4O shows an alignment of the predicted amino acid sequence of the light chain variable region for antibody 9.14.4 (residues 23-130 of SEQ ID NO: 28) to the germline V_κO12, J_κ3 sequence (SEQ ID NO: 103).
- 15 [0047] Figure 4P shows an alignment of the predicted amino acid sequence of the heavy chain variable region for antibody 9.14.4 (residues 20-135 of SEQ ID NO: 38) to the germline V_H3-11, D_H7-27, J_H4b sequence (SEQ ID NO: 116).
- [0048] Figure 4Q shows an alignment of the predicted amino acid sequence of the light chain variable region for antibody 9.7.2 (residues 23-130 of SEQ ID NO: 48) to the germline V_κO12, J_κ3 sequence (SEQ ID NO: 103).
- 20 [0049] Figure 4R shows an alignment of the predicted amino acid sequence of the heavy chain variable region for antibody 9.7.2 (residues 20-136 of SEQ ID NO: 46) to the germline V_H3-11, D_H6-13, J_H6b sequence (SEQ ID NO: 115).
- 25 [0050] Figure 4S shows an alignment of the predicted amino acid sequence of the light chain variable region for antibody 9.14.4I (residues 23-130 of SEQ ID NO: 28) to the germline V_κO12 J_κ3 sequence (SEQ ID NO: 103).
- [0051] Figure 4T shows an alignment of the predicted amino acid sequence of the heavy chain variable region for antibody 9.14.4I (residues 20-135 of SEQ ID NO: 26) to the germline V_H3-11, D_H7-27, J_H4b sequence (SEQ ID NO: 116).
- 30

- [0052] Figure 4U shows an alignment of the predicted amino acid sequence of the light chain variable region for antibody 8.10.3F (residues 21-129 of SEQ ID NO: 32) to the germline V_κA27, J_κ4 sequence (SEQ ID NO: 114).
- 5 [0053] Figure 4V shows an alignment of the predicted amino acid sequence of the heavy chain variable region for antibody 8.10.3F (residues 20-141 of SEQ ID NO: 30) to the germline V_H3-48, D_H1-26, J_H4b sequence (SEQ ID NO: 113).
- [0054] Figure 4W shows an alignment of the predicted amino acid sequence of the light chain variable region for antibody 9.7.2IF (residues 23-130 of SEQ ID NO: 36) to the germline V_κO12, J_κ3 sequence (SEQ ID NO: 103).
- 10 [0055] Figure 4X shows an alignment of the predicted amino acid sequence of the heavy chain variable region for antibody 9.7.2IF (residues 20-136 of SEQ ID NO: 34) to the germline V_H3-11, D_H6-13, J_H6b sequence (SEQ ID NO: 115).
- [0056] Figure 4Y shows an alignment of the predicted amino acid sequence of the light chain variable region for antibody 9.7.2C-Ser (residues 23-130 of SEQ ID NO: 52) to the germline V_κO12, J_κ3 sequence (SEQ ID NO: 103).
- 15 [0057] Figure 4Z shows an alignment of the predicted amino acid sequence of the heavy chain variable region for antibody 9.7.2C-Ser (residues 20-136 of SEQ ID NO: 50) to the germline V_H3-11, D_H6-13, J_H6b sequence (SEQ ID NO: 115).
- [0058] Figure 4AA shows an alignment of the predicted amino acid sequence of the light chain variable region for antibody 9.14.4C-Ser (residues 23-130 of SEQ ID NO: 56) to the germline V_κO12, J_κ3 sequence (SEQ ID NO: 103).
- 20 [0059] Figure 4BB shows an alignment of the predicted amino acid sequence of the heavy chain variable region for antibody 9.14.4C-Ser (residues 20-135 of SEQ ID NO: 54) to the germline V_H3-11, D_H7-27, J_H4b sequence (SEQ ID NO: 116).
- 25 [0060] Figure 4CC shows an alignment of the predicted amino acid sequence of the light chain variable region for antibody 8.10.3C-Ser (residues 21-129 of SEQ ID NO: 60) to the germline V_κA27, J_κ4 sequence (SEQ ID NO: 114).
- [0061] Figure 4DD shows an alignment of the predicted amino acid sequence of the heavy chain variable region for antibody 8.10.3C-Ser (residues 20-141 of SEQ ID NO: 58) to the germline V_H3-48, D_H1-26, J_H4b sequence (SEQ ID NO: 113).
- 30

- [0062] Figure 4EE shows an alignment of the predicted amino acid sequence of the light chain variable region for antibody 8.10.3-CG2 (residues 21-129 of SEQ ID NO: 60) to the germline V_κA27, J_κ4 sequence (SEQ ID NO: 114).
- 5 [0063] Figure 4FF shows an alignment of the predicted amino acid sequence of the heavy chain variable region for antibody 8.10.3-CG2 (residues 20-141 of SEQ ID NO: 62) to the germline V_H3-48, D_H1-26, J_H4b sequence (SEQ ID NO: 113).
- [0064] Figure 4GG shows an alignment of the predicted amino acid sequence of the light chain variable region for antibody 9.7.2-CG2 (residues 23-130 of SEQ ID NO: 52) to the germline V_κO12, J_κ3 sequence (SEQ ID NO: 103).
- 10 [0065] Figure 4HH shows an alignment of the predicted amino acid sequence of the heavy chain variable region for antibody 9.7.2-CG2 (residues 20-136 of SEQ ID NO: 66) to the germline V_H3-11, D_H6-13, J_H6b sequence (SEQ ID NO: 115).
- [0066] Figure 4II shows an alignment of the predicted amino acid sequence of the light chain variable region for antibody 9.7.2-CG4 (residues 23-130 of SEQ ID NO: 52) to the germline V_κO12, J_κ3 sequence (SEQ ID NO: 103).
- 15 [0067] Figure 4JJ shows an alignment of the predicted amino acid sequence of the heavy chain variable region for antibody 9.7.2-CG4 (residues 20-135 of SEQ ID NO: 70) to the germline V_H3-11, D_H6-13, J_H6b sequence (SEQ ID NO: 115).
- [0068] Figure 4KK shows an alignment of the predicted amino acid sequence of the light chain variable region for antibody 9.14.4-CG2 (residues 23-130 of SEQ ID NO: 56) to the germline V_κO12, J_κ3 sequence (SEQ ID NO: 103).
- 20 [0069] Figure 4LL shows an alignment of the predicted amino acid sequence of the heavy chain variable region for antibody 9.14.4-CG2 (residues 20-135 of SEQ ID NO: 74) to the germline V_H3-11, D_H7-27, J_H4b sequence (SEQ ID NO: 116).
- 25 [0070] Figure 4MM shows an alignment of the predicted amino acid sequence of the light chain variable region for antibody 9.14.4-CG4 (residues 23-130 of SEQ ID NO: 56) to the germline V_κO12, J_κ3 sequence (SEQ ID NO: 103).
- [0071] Figure 4NN shows an alignment of the predicted amino acid sequence of the heavy chain variable region for antibody 9.14.4-CG4 (residues 20-135 of SEQ ID NO: 78) to the germline V_H3-11, D_H7-27, J_H4b sequence (SEQ ID NO: 116).
- 30

- [0072] Figure 4OO shows an alignment of the predicted amino acid sequence of the light chain variable region for antibody 9.14.4-Ser (residues 23-130 of SEQ ID NO: 28) to the germline V_HO12, J_K3 sequence (SEQ ID NO: 103).
- 5 [0073] Figure 4PP shows an alignment of the predicted amino acid sequence of the heavy chain variable region for antibody 9.14.4-Ser (residues 20-135 of SEQ ID NO: 82) to the germline V_H3-11, D_H7-27, J_H4b sequence (SEQ ID NO: 116).
- [0074] Figure 4QQ shows an alignment of the predicted amino acid sequence of the light chain variable region for antibody 9.7.2-Ser (residues 23-130 of SEQ ID NO: 48) to the germline V_KO12, J_K3 sequence (SEQ ID NO: 103).
- 10 [0075] Figure 4RR shows an alignment of the predicted amino acid sequence of the heavy chain variable region for antibody 9.7.2-Ser (residues 20-136 of SEQ ID NO: 86) to the germline V_H3-11, D_H6-13, J_H6b sequence (SEQ ID NO: 115).
- [0076] Figure 4SS shows an alignment of the predicted amino acid sequence of the light chain variable region for antibody 8.10.3-Ser (residues 21-129 of SEQ ID NO: 44) to the germline V_KA27, J_K4 sequence (SEQ ID NO: 114).
- 15 [0077] Figure 4TT shows an alignment of the predicted amino acid sequence of the heavy chain variable region for antibody 8.10.3-Ser (residues 20-141 of SEQ ID NO: 90) to the germline V_H3-48, D_H1-26, J_H4b sequence (SEQ ID NO: 113).
- [0078] Figure 4UU shows an alignment of the predicted amino acid sequence of the light chain variable region for antibody 8.10.3-CG4 (residues 21-129 of SEQ ID NO: 60) to the germline V_KA27, J_K4 sequence (SEQ ID NO: 114).
- 20 [0079] Figure 4VV shows an alignment of the predicted amino acid sequence of the heavy chain variable region for antibody 8.10.3-CG4 (residues 20-141 of SEQ ID NO: 94) to the germline V_H3-48, D_H1-26, J_H4b sequence (SEQ ID NO: 113).
- 25 [0080] Figure 4WW shows an alignment of the predicted amino acid sequence of the light chain variable region for antibody 9.14.4G1 (residues 23-130 of SEQ ID NO: 28) to the germline V_KO12 J_K3 sequence (SEQ ID NO: 103).
- [0081] Figure 4XX shows an alignment of the predicted amino acid sequence of the heavy chain variable region for antibody 9.14.4G1 (residues 20-135 of SEQ ID NO: 102) to the germline V_H3-11, D_H7-27, J_H4b sequence (SEQ ID NO: 116).
- 30

[0082] Figure 4YY shows an alignment of the predicted amino acid sequence of the light chain variable region for antibody 8.10.3FG1 (residues 21-129 of SEQ ID NO:32) to the germline V_HA27, J_H4 sequence (SEQ ID NO: 114).

- 5 [0083] Figure 4ZZ shows an alignment of the predicted amino acid sequence of the heavy chain variable region for antibody 8.10.3FG1 (residues 20-141 of SEQ ID NO: 98) to the germline V_H3-48, D_H1-26, J_H4b sequence (SEQ ID NO: 113).

DETAILED DESCRIPTION OF THE INVENTION

Definitions and General Techniques

- 10 [0084] Unless otherwise defined herein, scientific and technical terms used in connection with the present invention shall have the meanings that are commonly understood by those of ordinary skill in the art. Further, unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular. Generally, nomenclatures used in connection with, and techniques of, cell and tissue culture, molecular biology, immunology, microbiology, genetics and protein and nucleic acid chemistry and hybridization described herein are those well known and commonly used in the art.

- 15 [0085] The methods and techniques of the present invention are generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification unless otherwise indicated. See, e.g., Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2d ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989) and Ausubel *et al.*, *Current Protocols in Molecular Biology*, Greene Publishing Associates (1992), and Harlow and Lane *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1990), which are incorporated herein by reference. Enzymatic reactions and purification techniques are performed according to manufacturer's specifications, as commonly accomplished in the art or as described herein. The nomenclatures used in connection with, and the laboratory procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well known and commonly used in the art. Standard techniques are used
- 20
- 25
- 30

for chemical syntheses, chemical analyses, pharmaceutical preparation, formulation, and delivery, and treatment of patients.

[0086] The following terms, unless otherwise indicated, shall be understood to have the following meanings:

- 5 [0087] The term "polypeptide" encompasses native or artificial proteins, protein fragments and polypeptide analogs of a protein sequence. A polypeptide may be monomeric or polymeric.

- [0088] The term "isolated protein", "isolated polypeptide" or "isolated antibody" is a protein, polypeptide or antibody that by virtue of its origin or source of
10 derivation has one to four of the following: (1) is not associated with naturally associated components that accompany it in its native state, (2) is free of other proteins from the same species, (3) is expressed by a cell from a different species, or (4) does not occur in nature. Thus, a polypeptide that is chemically synthesized or synthesized in a cellular system different from the cell from which it naturally
15 originates will be "isolated" from its naturally associated components. A protein may also be rendered substantially free of naturally associated components by isolation, using protein purification techniques well known in the art.

- [0089] Examples of isolated antibodies include an anti-M-CSF antibody that has been affinity purified using M-CSF, an anti-M-CSF antibody that has been
20 synthesized by a hybridoma or other cell line *in vitro*, and a human anti-M-CSF antibody derived from a transgenic mouse.

- [0090] A protein or polypeptide is "substantially pure," "substantially homogeneous," or "substantially purified" when at least about 60 to 75% of a sample exhibits a single species of polypeptide. The polypeptide or protein may be
25 monomeric or multimeric. A substantially pure polypeptide or protein will typically comprise about 50%, 60%, 70%, 80% or 90% W/W of a protein sample, more usually about 95%, and preferably will be over 99% pure. Protein purity or homogeneity may be indicated by a number of means well known in the art, such as polyacrylamide gel electrophoresis of a protein sample, followed by visualizing
30 a single polypeptide band upon staining the gel with a stain well known in the art. For certain purposes, higher resolution may be provided by using HPLC or other means well known in the art for purification.

[0091] The term "polypeptide fragment" as used herein refers to a polypeptide that has an amino-terminal and/or carboxy-terminal deletion, but where the remaining amino acid sequence is identical to the corresponding positions in the naturally-occurring sequence. In some embodiments, fragments are at least 5, 6, 8 or 10 amino acids long. In other embodiments, the fragments are at least 14, at least 20, at least 50, or at least 70, 80, 90, 100, 150 or 200 amino acids long.

[0092] The term "polypeptide analog" as used herein refers to a polypeptide that comprises a segment that has substantial identity to a portion of an amino acid sequence and that has at least one of the following properties: (1) specific binding to M-CSF under suitable binding conditions, (2) ability to inhibit M-CSF.

[0093] Typically, polypeptide analogs comprise a conservative amino acid substitution (or insertion or deletion) with respect to the normally-occurring sequence. Analogs typically are at least 20 or 25 amino acids long, preferably at least 50, 60, 70, 80, 90, 100, 150 or 200 amino acids long or longer, and can often be as long as a full-length polypeptide.

[0094] In certain embodiments, amino acid substitutions of the antibody or antigen-binding portion thereof are those which: (1) reduce susceptibility to proteolysis, (2) reduce susceptibility to oxidation, (3) alter binding affinity for forming protein complexes, or (4) confer or modify other physicochemical or functional properties of such analogs. Analogs can include various muteins of a sequence other than the normally-occurring peptide sequence. For example, single or multiple amino acid substitutions (preferably conservative amino acid substitutions) may be made in the normally-occurring sequence, preferably in the portion of the polypeptide outside the domain(s) forming intermolecular contacts.

[0095] A conservative amino acid substitution should not substantially change the structural characteristics of the parent sequence; e.g., a replacement amino acid should not alter the anti-parallel β -sheet that makes up the immunoglobulin binding domain that occurs in the parent sequence, or disrupt other types of secondary structure that characterizes the parent sequence. In general, glycine and proline analogs would not be used in an anti-parallel β -sheet. Examples of art-recognized polypeptide secondary and tertiary structures are described in *Proteins, Structures and Molecular Principles* (Creighton, Ed., W. H. Freeman

and Company, New York (1984)); *Introduction to Protein Structure* (C. Branden and J. Tooze, eds., Garland Publishing, New York, N.Y. (1991)); and Thornton *et al.*, *Nature* 354:105 (1991), which are each incorporated herein by reference.

- [0096] Non-peptide analogs are commonly used in the pharmaceutical industry as drugs with properties analogous to those of the template peptide. These types of non-peptide compound are termed "peptide mimetics" or "peptidomimetics." Fauchere, *J. Adv. Drug Res.* 15:29 (1986); Veber and Freidinger, *TINS* p.392 (1985); and Evans *et al.*, *J. Med. Chem.* 30:1229 (1987), which are incorporated herein by reference. Such compounds are often developed with the aid of
- 5 computerized molecular modeling. Peptide mimetics that are structurally similar to therapeutically useful peptides may be used to produce an equivalent therapeutic or prophylactic effect. Generally, peptidomimetics are structurally similar to a paradigm polypeptide (i.e., a polypeptide that has a desired biochemical property or pharmacological activity), such as a human antibody, but have one or more
- 10 peptide linkages optionally replaced by a linkage selected from the group consisting of: --CH₂NH--, --CH₂S--, --CH₂-CH₂--, --CH=CH--(cis and trans), --COCH₂--, --CH(OH)CH₂--, and --CH₂SO--, by methods well known in the art. Systematic substitution of one or more amino acids of a consensus sequence with a D-amino acid of the same type (e.g., D-lysine in place of L-lysine) may also be
- 20 used to generate more stable peptides. In addition, constrained peptides comprising a consensus sequence or a substantially identical consensus sequence variation may be generated by methods known in the art (Rizo and Gierasch, *Ann. Rev. Biochem.* 61:387 (1992), incorporated herein by reference); for example, by adding internal cysteine residues capable of forming intramolecular disulfide
- 25 bridges which cyclize the peptide.
- [0097] An "antibody" refers to an intact antibody or an antigen-binding portion that competes with the intact antibody for specific binding. See generally, Fundamental Immunology, Ch. 7 (Paul, W., ed., 2nd ed. Raven Press, N.Y. (1989)) (incorporated by reference in its entirety for all purposes). Antigen-binding
- 30 portions may be produced by recombinant DNA techniques or by enzymatic or chemical cleavage of intact antibodies. In some embodiments, antigen-binding portions include Fab, Fab', F(ab')₂, Fd, Fv, dAb, and complementarity determining

region (CDR) fragments, single-chain antibodies (scFv), chimeric antibodies, diabodies and polypeptides that contain at least a portion of an antibody that is sufficient to confer specific antigen binding to the polypeptide.

- [0098] From N-terminus to C-terminus, both the mature light and heavy chain variable domains comprise the regions FR1, CDR1, FR2, CDR2, FR3, CDR3 and FR4. The assignment of amino acids to each domain is in accordance with the definitions of Kabat, *Sequences of Proteins of Immunological Interest* (National Institutes of Health, Bethesda, Md. (1987 and 1991)), Chothia & Lesk, *J. Mol. Biol.* 196:901-917 (1987), or Chothia *et al.*, *Nature* 342:878-883 (1989).
- 5 [0099] As used herein, an antibody that is referred to by number is the same as a monoclonal antibody that is obtained from the hybridoma of the same number. For example, monoclonal antibody 3.8.3 is the same antibody as one obtained from hybridoma 3.8.3.
- [0100] As used herein, a Fd fragment means an antibody fragment that consists of the V_H and C_H 1 domains; an Fv fragment consists of the V_L and V_H domains of a single arm of an antibody; and a dAb fragment (Ward *et al.*, *Nature* 341:544-546 (1989)) consists of a V_H domain.
- [0101] In some embodiments, the antibody is a single-chain antibody (scFv) in which a V_L and V_H domains are paired to form a monovalent molecules via a synthetic linker that enables them to be made as a single protein chain. (Bird *et al.*, *Science* 242:423-426 (1988) and Huston *et al.*, *Proc. Natl. Acad. Sci. USA* 85:5879-5883 (1988).) In some embodiments, the antibodies are diabodies, i.e., are bivalent antibodies in which V_H and V_L domains are expressed on a single polypeptide chain, but using a linker that is too short to allow for pairing between
- 20 the two domains on the same chain, thereby forcing the domains to pair with complementary domains of another chain and creating two antigen binding sites. (See e.g., Holliger P. *et al.*, *Proc. Natl. Acad. Sci. USA* 90:6444-6448 (1993), and Poljak R. J. *et al.*, *Structure* 2:1121-1123 (1994).) In some embodiments, one or more CDRs from an antibody of the invention may be incorporated into a molecule
- 25 either covalently or noncovalently to make it an immunoadhesin that specifically binds to M-CSF. In such embodiments, the CDR(s) may be incorporated as part of
- 30

a larger polypeptide chain, may be covalently linked to another polypeptide chain, or may be incorporated noncovalently.

[0102] In embodiments having one or more binding sites, the binding sites may be identical to one another or may be different.

- 5 [0103] As used herein, the term "human antibody" means any antibody in which the variable and constant domain sequences are human sequences. The term encompasses antibodies with sequences derived from human genes, but which have been changed, e.g. to decrease possible immunogenicity, increase affinity, eliminate cysteines that might cause undesirable folding, etc. The term
- 10 encompasses such antibodies produced recombinantly in non-human cells, which might impart glycosylation not typical of human cells. These antibodies may be prepared in a variety of ways, as described below.

- [0104] The term "chimeric antibody" as used herein means an antibody that comprises regions from two or more different antibodies. In one embodiment, one
- 15 or more of the CDRs are derived from a human anti-M-CSF antibody. In another embodiment, all of the CDRs are derived from a human anti-M-CSF antibody. In another embodiment, the CDRs from more than one human anti-M-CSF antibodies are combined in a chimeric antibody. For instance, a chimeric antibody may comprise a CDR1 from the light chain of a first human anti-M-CSF antibody, a
- 20 CDR2 from the light chain of a second human anti-M-CSF antibody and a CDR3 from the light chain of a third human anti-M-CSF antibody, and the CDRs from the heavy chain may be derived from one or more other anti-M-CSF antibodies. Further, the framework regions may be derived from one of the anti-M-CSF antibodies from which one or more of the CDRs are taken or from one or more
- 25 different human antibodies.

- [0105] Fragments or analogs of antibodies or immunoglobulin molecules can be readily prepared by those of ordinary skill in the art following the teachings of this specification. Preferred amino- and carboxy-termini of fragments or analogs occur near boundaries of functional domains. Structural and functional domains can be
- 30 identified by comparison of the nucleotide and/or amino acid sequence data to public or proprietary sequence databases. Preferably, computerized comparison methods are used to identify sequence motifs or predicted protein conformation

domains that occur in other proteins of known structure and/or function. Methods to identify protein sequences that fold into a known three-dimensional structure are known. See Bowie *et al.*, *Science* 253:164 (1991).

- [0106] The term "surface plasmon resonance", as used herein, refers to an optical phenomenon that allows for the analysis of real-time biospecific interactions by detection of alterations in protein concentrations within a biosensor matrix, for example using the BIACORE™ system (Pharmacia Biosensor AB, Uppsala, Sweden and Piscataway, N.J.). For further descriptions, see Jonsson U. *et al.*, *Ann. Biol. Clin.* 51:19-26 (1993); Jonsson U. *et al.*, *Biotechniques* 11:620-627 (1991); Jonsson B. *et al.*, *J. Mol. Recognit.* 8:125-131 (1995); and Johnsson B. *et al.*, *Anal. Biochem.* 198:268-277 (1991).
- [0107] The term " K_D " refers to the equilibrium dissociation constant of a particular antibody-antigen interaction.
- [0108] The term "epitope" includes any protein determinant capable of specific binding to an immunoglobulin or T-cell receptor or otherwise interacting with a molecule. Epitopic determinants generally consist of chemically active surface groupings of molecules such as amino acids or sugar side chains and generally have specific three dimensional structural characteristics, as well as specific charge characteristics. An epitope may be "linear" or "conformational." In a linear epitope, all of the points of interaction between the protein and the interacting molecule (such as an antibody) occur linearly along the primary amino acid sequence of the protein. In a conformational epitope, the points of interaction occur across amino acid residues on the protein that are separated from one another. An antibody is said to specifically bind an antigen when the dissociation constant is ≤ 1 mM, preferably ≤ 100 nM and most preferably ≤ 10 nM. In certain embodiments, the K_D is 1 pM to 500 pM. In other embodiments, the K_D is between 500 pM to 1 μ M. In other embodiments, the K_D is between 1 μ M to 100 nM. In other embodiments, the K_D is between 100 mM to 10 nM. Once a desired epitope on an antigen is determined, it is possible to generate antibodies to that epitope, e.g., using the techniques described in the present invention. Alternatively, during the discovery process, the generation and characterization of antibodies may elucidate information about desirable epitopes. From this

information, it is then possible to competitively screen antibodies for binding to the same epitope. An approach to achieve this is to conduct cross-competition studies to find antibodies that competitively bind with one another, e.g., the antibodies compete for binding to the antigen. A high throughput process for "binning" antibodies based upon their cross-competition is described in International Patent Application No. WO 03/48731.

- [0109] As used herein, the twenty conventional amino acids and their abbreviations follow conventional usage. See *Immunology - A Synthesis* (2nd Edition, E.S. Golub and D.R. Gren, Eds., Sinauer Associates, Sunderland, Mass. (1991)), which is incorporated herein by reference.

[0110] The term "polynucleotide" as referred to herein means a polymeric form of nucleotides of at least 10 bases in length, either ribonucleotides or deoxynucleotides or a modified form of either type of nucleotide. The term includes single and double stranded forms.

- [0111] The term "isolated polynucleotide" as used herein means a polynucleotide of genomic, cDNA, or synthetic origin or some combination thereof, which by virtue of its origin or source of derivation, the "isolated polynucleotide" has one to three of the following: (1) is not associated with all or a portion of a polynucleotides with which the "isolated polynucleotide" is found in nature, (2) is operably linked to a polynucleotide to which it is not linked in nature, or (3) does not occur in nature as part of a larger sequence.

- [0112] The term "oligonucleotide" as used herein includes naturally occurring, and modified nucleotides linked together by naturally occurring and non-naturally occurring oligonucleotide linkages. Oligonucleotides are a polynucleotide subset generally comprising a length of 200 bases or fewer. Preferably oligonucleotides are 10 to 60 bases in length and most preferably 12, 13, 14, 15, 16, 17, 18, 19, or 20 to 40 bases in length. Oligonucleotides are usually single stranded, e.g. for primers and probes; although oligonucleotides may be double stranded, e.g. for use in the construction of a gene mutant. Oligonucleotides of the invention can be either sense or antisense oligonucleotides.

[0113] The term "naturally occurring nucleotides" as used herein includes deoxyribonucleotides and ribonucleotides. The term "modified nucleotides" as

- used herein includes nucleotides with modified or substituted sugar groups and the like. The term "oligonucleotide linkages" referred to herein includes oligonucleotides linkages such as phosphorothioate, phosphorodithioate, phosphoroselenoate, phosphorodiselenoate, phosphoroanilothioate,
- 5 phosphoraniladate, phosphoroamidate, and the like. See e.g., LaPlanche *et al.*, *Nucl. Acids Res.* 14:9081 (1986); Stec *et al.*, *J. Am. Chem. Soc.* 106:6077 (1984); Stein *et al.*, *Nucl. Acids Res.* 16:3209 (1988); Zon *et al.*, *Anti-Cancer Drug Design* 6:539 (1991); Zon *et al.*, *Oligonucleotides and Analogues: A Practical Approach*, pp. 87-108 (F. Eckstein, Ed., Oxford University Press, Oxford England (1991));
- 10 U.S. Patent No. 5,151,510; Uhlmann and Peyman, *Chemical Reviews* 90:543 (1990), the disclosures of which are hereby incorporated by reference. An oligonucleotide can include a label for detection, if desired.
- [0114] "Operably linked" sequences include both expression control sequences that are contiguous with the gene of interest and expression control sequences that
- 15 act in *trans* or at a distance to control the gene of interest. The term "expression control sequence" as used herein means polynucleotide sequences that are necessary to effect the expression and processing of coding sequences to which they are ligated. Expression control sequences include appropriate transcription initiation, termination, promoter and enhancer sequences; efficient RNA
- 20 processing signals such as splicing and polyadenylation signals; sequences that stabilize cytoplasmic mRNA; sequences that enhance translation efficiency (i.e., Kozak consensus sequence); sequences that enhance protein stability; and when desired, sequences that enhance protein secretion. The nature of such control sequences differs depending upon the host organism; in prokaryotes, such control
- 25 sequences generally include promoter, ribosomal binding site, and transcription termination sequence; in eukaryotes, generally, such control sequences include promoters and transcription termination sequence. The term "control sequences" is intended to include, at a minimum, all components whose presence is essential for expression and processing, and can also include additional components whose
- 30 presence is advantageous, for example, leader sequences and fusion partner sequences.

- [0115] The term "vector", as used herein, means a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. In some embodiments, the vector is a plasmid, i.e., a circular double stranded DNA loop into which additional DNA segments may be ligated. In some embodiments, the vector is a viral vector, wherein additional DNA segments may be ligated into the viral genome. In some embodiments, the vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). In other embodiments, the vectors (e.g., non-episomal mammalian vectors) can be integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as "recombinant expression vectors" (or simply, "expression vectors").
- [0116] The term "recombinant host cell" (or simply "host cell"), as used herein, means a cell into which a recombinant expression vector has been introduced. It should be understood that "recombinant host cell" and "host cell" mean not only the particular subject cell but also the progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term "host cell" as used herein.
- [0117] The term "selectively hybridize" referred to herein means to detectably and specifically bind. Polynucleotides, oligonucleotides and fragments thereof in accordance with the invention selectively hybridize to nucleic acid strands under hybridization and wash conditions that minimize appreciable amounts of detectable binding to nonspecific nucleic acids. "High stringency" or "highly stringent" conditions can be used to achieve selective hybridization conditions as known in the art and discussed herein. One example of "high stringency" or "highly stringent" conditions is the incubation of a polynucleotide with another polynucleotide, wherein one polynucleotide may be affixed to a solid surface such as a membrane, in a hybridization buffer of 6X SSPE or SSC, 50% formamide, 5X Denhardt's reagent, 0.5% SDS, 100 μ g/ml denatured, fragmented salmon sperm

DNA at a hybridization temperature of 42°C for 12-16 hours, followed by twice washing at 55°C using a wash buffer of 1X SSC, 0.5% SDS. See also Sambrook *et al., supra*, pp. 9.50-9.55.

- [0118] The term “percent sequence identity” in the context of nucleic acid sequences means the percent of residues when a first contiguous sequence is compared and aligned for maximum correspondence to a second contiguous sequence. The length of sequence identity comparison may be over a stretch of at least about nine nucleotides, usually at least about 18 nucleotides, more usually at least about 24 nucleotides, typically at least about 28 nucleotides, more typically at least about 32 nucleotides, and preferably at least about 36, 48 or more nucleotides. There are a number of different algorithms known in the art which can be used to measure nucleotide sequence identity. For instance, polynucleotide sequences can be compared using FASTA, Gap or Bestfit, which are programs in Wisconsin Package Version 10.0, Genetics Computer Group (GCG), Madison, Wisconsin. FASTA, which includes, e.g., the programs FASTA2 and FASTA3, provides alignments and percent sequence identity of the regions of the best overlap between the query and search sequences (Pearson, *Methods Enzymol.* 183:63-98 (1990); Pearson, *Methods Mol. Biol.* 132:185-219 (2000); Pearson, *Methods Enzymol.* 266:227-258 (1996); Pearson, *J. Mol. Biol.* 276:71-84 (1998); herein incorporated by reference). Unless otherwise specified, default parameters for a particular program or algorithm are used. For instance, percent sequence identity between nucleic acid sequences can be determined using FASTA with its default parameters (a word size of 6 and the NOPAM factor for the scoring matrix) or using Gap with its default parameters as provided in GCG Version 6.1, herein incorporated by reference.
- [0119] A reference to a nucleotide sequence encompasses its complement unless otherwise specified. Thus, a reference to a nucleic acid having a particular sequence should be understood to encompass its complementary strand, with its complementary sequence.
- [0120] The term “percent sequence identity” means a ratio, expressed as a percent of the number of identical residues over the number of residues compared.

[0121] The term "substantial similarity" or "substantial sequence similarity," when referring to a nucleic acid or fragment thereof, means that when optimally aligned with appropriate nucleotide insertions or deletions with another nucleic acid (or its complementary strand), there is nucleotide sequence identity in at least about 85%, preferably at least about 90%, and more preferably at least about 95%, 96%, 97%, 98% or 99% of the nucleotide bases, as measured by any well-known algorithm of sequence identity, such as FASTA, BLAST or Gap, as discussed above.

[0122] As applied to polypeptides, the term "substantial identity" means that two peptide sequences, when optimally aligned, such as by the programs GAP or BESTFIT using default gap weights, as supplied with the programs, share at least 70%, 75% or 80% sequence identity, preferably at least 90% or 95% sequence identity, and more preferably at least 97%, 98% or 99% sequence identity. In certain embodiments, residue positions that are not identical differ by conservative amino acid substitutions. A "conservative amino acid substitution" is one in which an amino acid residue is substituted by another amino acid residue having a side chain R group with similar chemical properties (e.g., charge or hydrophobicity). In general, a conservative amino acid substitution will not substantially change the functional properties of a protein. In cases where two or more amino acid sequences differ from each other by conservative substitutions, the percent sequence identity may be adjusted upwards to correct for the conservative nature of the substitution. Means for making this adjustment are well-known to those of skill in the art. See, e.g., Pearson, *Methods Mol. Biol.* 243:307-31 (1994). Examples of groups of amino acids that have side chains with similar chemical properties include 1) aliphatic side chains: glycine, alanine, valine, leucine, and isoleucine; 2) aliphatic-hydroxyl side chains: serine and threonine; 3) amide-containing side chains: asparagine and glutamine; 4) aromatic side chains: phenylalanine, tyrosine, and tryptophan; 5) basic side chains: lysine, arginine, and histidine; 6) acidic side chains: aspartic acid and glutamic acid; and 7) sulfur-containing side chains: cysteine and methionine. Conservative amino acids substitution groups are: valine-leucine-isoleucine, phenylalanine-tyrosine, lysine-arginine, alanine-valine, glutamate-aspartate, and asparagine-glutamine.

[0123] Alternatively, a conservative replacement is any change having a positive value in the PAM250 log-likelihood matrix disclosed in Gonnet *et al.*, *Science* 256:1443-45 (1992), herein incorporated by reference. A “moderately conservative” replacement is any change having a nonnegative value in the PAM250 log-likelihood matrix.

[0124] Sequence identity for polypeptides, is typically measured using sequence analysis software. Protein analysis software matches sequences using measures of similarity assigned to various substitutions, deletions and other modifications, including conservative amino acid substitutions. For instance, GCG contains programs such as “Gap” and “Bestfit” which can be used with default parameters, as specified with the programs, to determine sequence homology or sequence identity between closely related polypeptides, such as homologous polypeptides from different species of organisms or between a wild type protein and a mutein thereof. See, e.g., GCG Version 6.1. Polypeptide sequences also can be compared using FASTA using default or recommended parameters, see GCG Version 6.1. (University of Wisconsin WI) FASTA (e.g., FASTA2 and FASTA3) provides alignments and percent sequence identity of the regions of the best overlap between the query and search sequences (Pearson, *Methods Enzymol.* 183:63-98 (1990); Pearson, *Methods Mol. Biol.* 132:185-219 (2000)). Another preferred algorithm when comparing a sequence of the invention to a database containing a large number of sequences from different organisms is the computer program BLAST, especially blastp or tblastn, using default parameters, as supplied with the programs. See, e.g., Altschul *et al.*, *J. Mol. Biol.* 215:403-410 (1990); Altschul *et al.*, *Nucleic Acids Res.* 25:3389-402 (1997).

[0125] The length of polypeptide sequences compared for homology will generally be at least about 16 amino acid residues, usually at least about 20 residues, more usually at least about 24 residues, typically at least about 28 residues, and preferably more than about 35 residues. When searching a database containing sequences from a large number of different organisms, it is preferable to compare amino acid sequences.

[0126] As used herein, the terms “label” or “labeled” refers to incorporation of another molecule in the antibody. In one embodiment, the label is a detectable

- marker, e.g., incorporation of a radiolabeled amino acid or attachment to a polypeptide of biotinyl moieties that can be detected by marked avidin (e.g., streptavidin containing a fluorescent marker or enzymatic activity that can be detected by optical or colorimetric methods). In another embodiment, the label or
- 5 marker can be therapeutic, e.g., a drug conjugate or toxin. Various methods of labeling polypeptides and glycoproteins are known in the art and may be used. Examples of labels for polypeptides include, but are not limited to, the following: radioisotopes or radionuclides (e.g., ^3H , ^{14}C , ^{15}N , ^{35}S , ^{90}Y , ^{99}Tc , ^{111}In , ^{125}I , ^{131}I), fluorescent labels (e.g., FITC, rhodamine, lanthanide phosphors), enzymatic labels
- 10 (e.g., horseradish peroxidase, β -galactosidase, luciferase, alkaline phosphatase), chemiluminescent markers, biotinyl groups, predetermined polypeptide epitopes recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags), magnetic agents, such as gadolinium chelates, toxins such as pertussis toxin, taxol,
- 15 cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof. In some
- 20 embodiments, labels are attached by spacer arms of various lengths to reduce potential steric hindrance.

- [0127] Throughout this specification and claims, the word "comprise," or variations such as "comprises" or "comprising," will be understood to imply the inclusion of a stated integer or group of integers but not the exclusion of any other
- 25 integer or group of integers.

Human Anti-M-CSF Antibodies and Characterization Thereof

- [0128] In one embodiment, the invention provides humanized anti-M-CSF antibodies. In another embodiment, the invention provides human anti-M-CSF antibodies. In some embodiments, human anti-M-CSF antibodies are produced by
- 30 immunizing a non-human transgenic animal, e.g., a rodent, whose genome comprises human immunoglobulin genes so that the rodent produces human antibodies.

[0129] An anti-M-CSF antibody of the invention can comprise a human kappa or a human lamda light chain or an amino acid sequence derived therefrom. In some embodiments comprising a kappa light chain, the light chain variable domain (V_L) is encoded in part by a human V_KO12, V_KL2, V_KL5, V_KA27 or V_KB3 gene and a
5 J_K1, J_K2, J_K3, or J_K4 gene. In particular embodiments of the invention, the light chain variable domain is encoded by V_KO12/J_K3, V_KL2/J_K3, V_KL5/J_K3, V_KL5/J_K4, V_KA27/J_K4 or V_KB3/J_K1 gene.

[0130] In some embodiments, the V_L of the M-CSF antibody comprises one or more amino acid substitutions relative to the germline amino acid sequence. In
10 some embodiments, the V_L of the anti-M-CSF antibody comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 amino acid substitutions relative to the germline amino acid sequence. In some embodiments, one or more of those substitutions from germline is in the CDR regions of the light chain. In some embodiments, the amino acid substitutions relative to germline are at one or more of the same positions as the
15 substitutions relative to germline in any one or more of the V_L of antibodies 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 or 9.14.4G1. For example, the V_L of the anti-M-CSF antibody may contain one or
20 more amino acid substitutions compared to germline found in the V_L of antibody 88, and other amino acid substitutions compared to germline found in the V_L of antibody 252 which utilizes the same V_K gene as antibody 88. In some embodiments, the amino acid changes are at one or more of the same positions but involve a different mutation than in the reference antibody.

[0131] In some embodiments, amino acid changes relative to germline occur at one or more of the same positions as in any of the V_L of antibodies 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 or 9.14.4G1, but
25 the changes may represent conservative amino acid substitutions at such position(s) relative to the amino acid in the reference antibody. For example, if a particular position in one of these antibodies is changed relative to germline and is
30

glutamate, one may substitute aspartate at that position. Similarly, if an amino acid substitution compared to germline is serine, one may substitute threonine for serine at that position. Conservative amino acid substitutions are discussed *supra*.

- [0132] In some embodiments, the light chain of the human anti-M-CSF antibody comprises the amino acid sequence that is the same as the amino acid sequence of the V_L of antibody 252 (SEQ ID NO: 4), 88 (SEQ ID NO: 8), 100 (SEQ ID NO: 12), 3.8.3 (SEQ ID NO: 16), 2.7.3 (SEQ ID NO: 20), 1.120.1 (SEQ ID NO: 24), 9.14.4I (SEQ ID NO: 28), 8.10.3F (SEQ ID NO: 32), 9.7.2IF (SEQ ID NO: 36), 9.14.4 (SEQ ID NO: 28), 8.10.3 (SEQ ID NO: 44), 9.7.2 (SEQ ID NO: 48), 9.7.2C-Ser (SEQ ID NO: 52), 9.14.4C-Ser (SEQ ID NO: 56), 8.10.3C-Ser (SEQ ID NO: 60), 8.10.3-CG2 (SEQ ID NO: 60), 9.7.2-CG2 (SEQ ID NO: 52), 9.7.2-CG4 (SEQ ID NO: 52), 9.14.4-CG2 (SEQ ID NO: 56), 9.14.4-CG4 (SEQ ID NO: 56), 9.14.4-Ser (SEQ ID NO: 28), 9.7.2-Ser (SEQ ID NO: 48), 8.10.3-Ser (SEQ ID NO: 44), 8.10.3-CG4 (SEQ ID NO: 60) 8.10.3FG1 (SEQ ID NO: 32) or 9.14.4G1 (SEQ ID NO: 28), or said amino acid sequence having up to 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 conservative amino acid substitutions and/or a total of up to 3 non-conservative amino acid substitutions. In some embodiments, the light chain comprises the amino acid sequence from the beginning of the CDR1 to the end of the CDR3 of any one of the foregoing antibodies.
- [0133] In some embodiments, the light chain of the anti-M-CSF antibody comprises at least the light chain CDR1, CDR2 or CDR3 of a germline or antibody sequence, as described herein. In another embodiment, the light chain may comprise a CDR1, CDR2 or CDR3 regions of an antibody independently selected from 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 or 9.14.4G1, or CDR regions each having less than 4 or less than 3 conservative amino acid substitutions and/or a total of three or fewer non-conservative amino acid substitutions. In other embodiments, the light chain of the anti-M-CSF antibody comprises the light chain CDR1, CDR2 or CDR3, each of which are independently selected from the CDR1, CDR2 and CDR3 regions of an antibody having a light chain variable region comprising the amino acid sequence

- of the V_L region selected from SEQ ID NOS: 4, 8, 12, 16, 20, 24, 28, 32, 36, 44, 48, 52, 56 or 60, or encoded by a nucleic acid molecule encoding the V_L region selected from SEQ ID NOS: 3, 7, 11, 27, 31, 35, 43 or 47. The light chain of the anti-M-CSF antibody may comprise the CDR1, CDR2 and CDR3 regions of an antibody comprising the amino acid sequence of the V_L region selected from 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 or 9.14.4G1 or SEQ ID NOS: 4, 8, 12, 16, 20, 24, 28, 32, 36, 44, 48, 52, 56 or 60.
- 10 [0134] In some embodiments, the light chain comprises the CDR1, CDR2 and CDR3 regions of antibody 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 or 9.14.4G1, or said CDR regions each having less than 4
- 15 or less than 3 conservative amino acid substitutions and/or a total of three or fewer non-conservative amino acid substitutions.
- [0135] With regard to the heavy chain, in some embodiments, the variable region of the heavy chain amino acid sequence is encoded in part by a human V_H3-11, V_H3-23, V_H3-7, V_H1-18, V_H3-33, V_H3-48 gene and a J_H4, J_H6, J_H4b, or J_H6b gene.
- 20 In a particular embodiment of the invention, the heavy chain variable region is encoded by V_H3-11/D_H7-27/J_H6, V_H3-7/D_H6-13/J_H4, V_H3-23/D_H1-26/J_H4, V_H3-11/D_H7-27/J_H4, V_H3-33/D_H1-26/J_H4, V_H1-18/D_H4-23/J_H4, V_H3-11/D_H7-27/J_H4b, V_H3-48/D_H1-26/J_H4b, V_H3-11/D_H6-13/J_H6b, V_H3-11/D_H7-27/J_H4b, V_H3-48/D_H1-6/J_H4b, or V_H3-11/D_H6-13/J_H6b gene. In some embodiments, the V_H of the anti-
- 25 M-CSF antibody contains one or more amino acid substitutions, deletions or insertions (additions) relative to the germline amino acid sequence. In some embodiments, the variable domain of the heavy chain comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, or 18 mutations from the germline amino acid sequence. In some embodiments, the mutation(s) are non-conservative
- 30 substitutions compared to the germline amino acid sequence. In some embodiments, the mutations are in the CDR regions of the heavy chain. In some embodiments, the amino acid changes are made at one or more of the same

positions as the mutations from germline in any one or more of the V_H of antibodies 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 or 9.14.4G1. In other embodiments, the amino acid changes are at one or more of the same positions but involve a different mutation than in the reference antibody.

[0136] In some embodiments, the heavy chain comprises an amino acid sequence of the variable domain (V_H) of antibody 252 (SEQ ID NO: 2), 88 (SEQ ID NO: 6), 100 (SEQ ID NO: 10), 3.8.3 (SEQ ID NO: 14), 2.7.3 (SEQ ID NO: 18), 1.120.1 (SEQ ID NO: 22), 9.14.4I (SEQ ID NO: 26), 8.10.3F (SEQ ID NO: 30), 9.7.2IF (SEQ ID NO: 34), 9.14.4 (SEQ ID NO: 38), 8.10.3 (SEQ ID NO: 30), 9.7.2 (SEQ ID NO: 46), 9.7.2C-Ser (SEQ ID NO: 50), 9.14.4C-Ser (SEQ ID NO: 54), 8.10.3C-Ser (SEQ ID NO: 58), 8.10.3-CG2 (SEQ ID NO: 62), 9.7.2-CG2 (SEQ ID NO: 66), 9.7.2-CG4 (SEQ ID NO: 70), 9.14.4-CG2 (SEQ ID NO: 74), 9.14.4-CG4 (SEQ ID NO: 78), 9.14.4-Ser (SEQ ID NO: 82), 9.7.2-Ser (SEQ ID NO: 86), 8.10.3-Ser (SEQ ID NO: 90), 8.10.3-CG4 (SEQ ID NO: 94), 8.10.3FG1 (SEQ ID NO: 98) or 9.14.4G1 (SEQ ID NO: 102), or said amino acid sequence having up to 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 conservative amino acid substitutions and/or a total of up to 3 non-conservative amino acid substitutions. In some embodiments, the heavy chain comprises the amino acid sequence from the beginning of the CDR1 to the end of the CDR3 of any one of the foregoing antibodies.

[0137] In some embodiments, the heavy chain comprises the heavy chain CDR1, CDR2 and CDR3 regions of antibody 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 or 9.14.4G1, or said CDR regions each having less than 8, less than 6, less than 4, or less than 3 conservative amino acid substitutions and/or a total of three or fewer non-conservative amino acid substitutions.

[0138] In some embodiments, the heavy chain comprises a germline or antibody CDR3, as described above, of an antibody sequence as described herein, and may

- also comprise the CDR1 and CDR2 regions of a germline sequence, or may comprise a CDR1 and CDR2 of an antibody sequence, each of which are independently selected from an antibody comprising a heavy chain of an antibody selected from 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4,
- 5 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 or 9.14.4G1. In another embodiment, the heavy chain comprises a CDR3 of an antibody sequence as described herein, and may also comprise the CDR1 and CDR2 regions, each of which are independently selected from a CDR1
- 10 and CDR2 region of a heavy chain variable region comprising an amino acid sequence of the V_H region selected from SEQ ID NOS: 2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 46, 50, 54, 58, 62, 66, 70, 74, 78, 82, 86, 90, 94, 98 or 102, or encoded by a nucleic acid sequence encoding the V_H region selected from SEQ ID NOS: 1, 5, 9, 25, 29, 33, 37, 45, 97 or 101. In another embodiment, the antibody comprises a
- 15 light chain as disclosed above and a heavy chain as disclosed above.
- [0139] One type of amino acid substitution that may be made is to change one or more cysteines in the antibody, which may be chemically reactive, to another residue, such as, without limitation, alanine or serine. In one embodiment, there is a substitution of a non-canonical cysteine. The substitution can be in a framework
- 20 region of a variable domain or in the constant domain of an antibody. In another embodiment, the cysteine is in a non-canonical region of the antibody.
- [0140] Another type of amino acid substitution that may be made is to remove any potential proteolytic sites in the antibody, particularly those that are in a CDR or framework region of a variable domain or in the constant domain of an
- 25 antibody. Substitution of cysteine residues and removal of proteolytic sites may decrease the risk of any heterogeneity in the antibody product and thus increase its homogeneity. Another type of amino acid substitution is elimination of asparagine-glycine pairs, which form potential deamidation sites, by altering one or both of the residues.
- 30 [0141] In some embodiments, the C-terminal lysine of the heavy chain of the anti-M-CSF antibody of the invention is not present (Lewis D.A., *et al.*, *Anal. Chem.*, 66(5): 585-95 (1994)). In various embodiments of the invention, the heavy

and light chains of the anti-M-CSF antibodies may optionally include a signal sequence.

- [0142] In one aspect, the invention relates to inhibiting human anti-M-CSF monoclonal antibodies and the cell lines engineered to produce them. Table 1 lists
- 5 the sequence identifiers (SEQ ID NOS) of the nucleic acids that encode the variable region of the heavy and light chains and the corresponding predicted amino acid sequences for the monoclonal antibodies: 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3 and 9.7.2. Additional variant antibodies 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-
- 10 CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4 8.10.3FG1 or 9.14.4G1 could be made by methods known to one skilled in the art.

Table 1

HUMAN ANTI-M-CSF ANTIBODIES				
MAb	SEQUENCE IDENTIFIER (SEQ ID NOS:)			
	Full Length			
	Heavy		Light	
	DNA	Protein	DNA	Protein
252	1	2	3	4
88	5	6	7	8
100	9	10	11	12
3.8.3		14		16
2.7.3		18		20
1.120.1		22		24
9.14.4I	25	26	27	28
9.14.4	37	38	27	28
9.14.4C-Ser		54		56
9.14.4-CG2		74		56
9.14.4-CG4		78		56
9.14.4-Ser		82	27	28
9.14.4-G1	101	102	27	28
8.10.3F	29	30	31	32
8.10.3	29	30	43	44
8.10.3C-Ser		58		60
8.10.3-CG2		62		60
8.10.3-Ser		90	43	44
8.10.3-CG4		94		60
8.10.3FG1	97	98	31	32
9.7.2IF	33	34	35	36
9.7.2	45	46	47	48
9.7.2C-Ser		50		52
9.7.2-CG2		66		52
9.7.2-CG4		70		52
9.7.2-Ser		86	47	48

Class and Subclass of Anti-M-CSF Antibodies

[0143] The class and subclass of anti-M-CSF antibodies may be determined by any method known in the art. In general, the class and subclass of an antibody
5 may be determined using antibodies that are specific for a particular class and subclass of antibody. Such antibodies are commercially available. The class and subclass can be determined by ELISA, or Western Blot as well as other techniques. Alternatively, the class and subclass may be determined by sequencing all or a portion of the constant domains of the heavy and/or light chains of the antibodies,
10 comparing their amino acid sequences to the known amino acid sequences of various class and subclasses of immunoglobulins, and determining the class and subclass of the antibodies.

[0144] In some embodiments, the anti-M-CSF antibody is a monoclonal antibody. The anti-M-CSF antibody can be an IgG, an IgM, an IgE, an IgA, or an
15 IgD molecule. In preferred embodiments, the anti-M-CSF antibody is an IgG and is an IgG1, IgG2, IgG3 or IgG4 subclass. In other preferred embodiments, the antibody is subclass IgG2 or IgG4. In another preferred embodiment, the antibody is subclass IgG1.

Species and Molecular Selectivity

20 [0145] In another aspect of the invention, the anti-M-CSF antibodies demonstrate both species and molecule selectivity. In some embodiments, the anti-M-CSF antibody binds to human, cynomolgus monkey and mouse M-CSF. Following the teachings of the specification, one may determine the species selectivity for the anti-M-CSF antibody using methods well known in the art. For instance, one may
25 determine the species selectivity using Western blot, FACS, ELISA, RIA, a cell proliferation assay, or a M-CSF receptor binding assay. In a preferred embodiment, one may determine the species selectivity using a cell proliferation assay or ELISA.

[0146] In another embodiment, the anti-M-CSF antibody has a selectivity for M-CSF that is at least 100 times greater than its selectivity for GM-/G-CSF. In some
30 embodiments, the anti-M-CSF antibody does not exhibit any appreciable specific

binding to any other protein other than M-CSF. One can determine the selectivity of the anti-M-CSF antibody for M-CSF using methods well known in the art following the teachings of the specification. For instance one can determine the selectivity using Western blot, FACS, ELISA, or RIA.

5 Identification of M-CSF Epitopes Recognized by Anti- M-CSF Antibodies

[0147] The invention provides a human anti-M-CSF monoclonal antibody that binds to M-CSF and competes with, cross-competes with and/or binds the same epitope and/or binds to M-CSF with the same K_D as (a) an antibody selected from 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 10 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 or 9.14.4G1; (b) an antibody that comprises a heavy chain variable region having an amino acid sequence of SEQ ID NO: 2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 46, 50, 54, 58, 62, 66, 70, 74, 78, 82, 86, 90, 94, 98 or 102; (c) an antibody that comprises a 15 light chain variable region having an amino acid sequence of SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 36, 44, 48, 52, 56 or 60; (d) an antibody that comprises both a heavy chain variable region as defined in (b) and a light chain variable region as defined in (c).

[0148] One can determine whether an antibody binds to the same epitope, 20 competes for binding with, cross competes for binding with or has the same K_D an anti-M-CSF antibody by using methods known in the art. In one embodiment, one allows the anti-M-CSF antibody of the invention to bind to M-CSF under saturating conditions and then measures the ability of the test antibody to bind to M-CSF. If the test antibody is able to bind to M-CSF at the same time as the anti- 25 M-CSF antibody, then the test antibody binds to a different epitope as the anti-M-CSF antibody. However, if the test antibody is not able to bind to M-CSF at the same time, then the test antibody binds to the same epitope, an overlapping epitope, or an epitope that is in close proximity to the epitope bound by the human anti-M-CSF antibody. This experiment can be performed using ELISA, RIA, or 30 FACS. In a preferred embodiment, the experiment is performed using BIACORE™.

Binding Affinity of Anti-M-CSF Antibodies to M-CSF

[0149] In some embodiments of the invention, the anti-M-CSF antibodies bind to M-CSF with high affinity. In some embodiments, the anti-M-CSF antibody binds to M-CSF with a K_D of 1×10^{-7} M or less. In other preferred embodiments, the antibody binds to M-CSF with a K_D of 1×10^{-8} M, 1×10^{-9} M, 1×10^{-10} M, 1×10^{-11} M, 1×10^{-12} M or less. In certain embodiments, the K_D is 1 pM to 500 pM. In other embodiments, the K_D is between 500 pM to 1 μ M. In other embodiments, the K_D is between 1 μ M to 100 nM. In other embodiments, the K_D is between 100 mM to 10 nM. In an even more preferred embodiment, the antibody binds to M-CSF with substantially the same K_D as an antibody selected from 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 or 9.14.4G1. In another preferred embodiment, the antibody binds to M-CSF with substantially the same K_D as an antibody that comprises a CDR2 of a light chain, and/or a CDR3 of a heavy chain from an antibody selected from 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 or 9.14.4G1. In still another preferred embodiment, the antibody binds to M-CSF with substantially the same K_D as an antibody that comprises a heavy chain variable region having an amino acid sequence of SEQ ID NO: 2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 46, 50, 54, 58, 62, 66, 70, 74, 78, 82, 86, 90, 94, 98 or 102, or that comprises a light chain variable region having an amino acid sequence of SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 36, 44, 48, 52, 56 or 60. In another preferred embodiment, the antibody binds to M-CSF with substantially the same K_D as an antibody that comprises a CDR2, and may optionally comprise a CDR1 and/or CDR3, of a light chain variable region having an amino acid sequence of the V_L region of SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 36, 44, 48, 52, 56 or 60, or that comprises a CDR3, and may optionally comprise a CDR1 and/or CDR2, of a heavy chain variable region having an amino acid sequence of the V_H region of SEQ ID NO: 2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 46, 50, 54, 58, 62, 66, 70, 74, 78, 82, 86, 90, 94, 98 or 102.

- [0150] In some embodiments, the anti-M-CSF antibody has a low dissociation rate. In some embodiments, the anti-M-CSF antibody has an k_{off} of $2.0 \times 10^{-4} \text{ s}^{-1}$ or lower. In other preferred embodiments, the antibody binds to M-CSF with a k_{off} of 2.0×10^{-5} or a k_{off} of $2.0 \times 10^{-6} \text{ s}^{-1}$ or lower. In some embodiments, the k_{off} is
- 5 substantially the same as an antibody described herein, such as an antibody selected from 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 or 9.14.4G1. In some embodiments, the antibody binds to M-CSF with
- 10 substantially the same k_{off} as an antibody that comprises (a) a CDR3, and may optionally comprise a CDR1 and/or CDR2, of a heavy chain of an antibody selected from 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 or 9.14.4G1; or (b) a CDR2, and may optionally comprise a CDR1
- 15 and/or CDR3, of a light chain from an antibody selected from 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 or 9.14.4G1. In some
- 20 embodiments, the antibody binds to M-CSF with substantially the same k_{off} as an antibody that comprises a heavy chain variable region having an amino acid sequence of SEQ ID NO: 2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 46, 50, 54, 58, 62, 66, 70, 74, 78, 82, 86, 90, 94, 98 or 102; or that comprises a light chain variable region having an amino acid sequence of SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 36, 44,
- 25 48, 52, 56 or 60; In another preferred embodiment, the antibody binds to M-CSF with substantially the same k_{off} as an antibody that comprises a CDR2, and may optionally comprise a CDR1 and/or CDR3, of a light chain variable region having an amino acid sequence of SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 36, 44, 48, 52, 56 or 60; or a CDR3, and may optionally comprise a CDR1 and/or CDR2, of a
- 30 heavy chain variable region having an amino acid sequence of SEQ ID NO: 2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 46, 50, 54, 58, 62, 66, 70, 74, 78, 82, 86, 90, 94, 98 or 102.

[0151] The binding affinity and dissociation rate of an anti-M-CSF antibody to a M-CSF can be determined by methods known in the art. The binding affinity can be measured by competitive ELISAs, RIAs, surface plasmon resonance (e.g., by using BIACORE™ technology). The dissociation rate can be measured by surface plasmon resonance. Preferably, the binding affinity and dissociation rate is measured by surface plasmon resonance. More preferably, the binding affinity and dissociation rate are measured using BIACORE™ technology. Example VI exemplifies a method for determining affinity constants of anti-M-CSF monoclonal antibodies by BIACORE™ technology.

10 Inhibition of M-CSF Activity by Anti-M-CSF Antibody

Inhibition of M-CSF binding to c-fms

[0152] In another embodiment, the invention provides an anti-M-CSF antibody that inhibits the binding of a M-CSF to *c-fms* receptor and blocks or prevents activation of *c-fms*. In an preferred embodiment, the M-CSF is human. In another preferred embodiment, the anti-M-CSF antibody is a human antibody. The IC₅₀ can be measured by ELISA, RIA, and cell based assays such as a cell proliferation assay, a whole blood monocyte shape change assay, or a receptor binding inhibition assay. In one embodiment, the antibody or portion thereof inhibits cell proliferation with an IC₅₀ of no more than 8.0×10^{-7} M, preferably no more than 3×10^{-7} M, or more preferably no more than 8×10^{-8} M as measured by a cell proliferation assay. In another embodiment, the IC₅₀ as measured by a monocyte shape change assay is no more than 2×10^{-6} M, preferably no more than 9.0×10^{-7} M, or more preferably no more than 9×10^{-8} M. In another preferred embodiment, the IC₅₀ as measured by a receptor binding assay is no more than 2×10^{-6} M, preferably no more than 8.0×10^{-7} M, or more preferably no more than 7.0×10^{-8} M. Examples III, IV, and V exemplify various types of assays.

[0153] In another aspect anti-M-CSF antibodies of the invention inhibit monocyte/macrophage cell proliferation in response to a M-CSF by at least 20%, more preferably 40%, 45%, 50%, 55%, 60%, 65%, 70%, 80%, 85%, 90%, 95% or 100% compared to the proliferation of cell in the absence of antibody.

Methods of Producing Antibodies and Antibody Producing Cell Lines

Immunization

[0154] In some embodiments, human antibodies are produced by immunizing a non-human animal comprising in its genome some or all of human immunoglobulin heavy chain and light chain loci with a M-CSF antigen. In a preferred
5 globulin heavy chain and light chain loci with a M-CSF antigen. In a preferred embodiment, the non-human animal is a XENOMOUSE™ animal (Abgenix Inc., Fremont, CA). Another non-human animal that may be used is a transgenic mouse produced by Medarex (Medarex, Inc., Princeton, NJ).

[0155] XENOMOUSE™ mice are engineered mouse strains that comprise large
10 fragments of human immunoglobulin heavy chain and light chain loci and are deficient in mouse antibody production. See, e.g., Green *et al.*, *Nature Genetics* 7:13-21 (1994) and U.S. Patents 5,916,771, 5,939,598, 5,985,615, 5,998,209, 6,075,181, 6,091,001, 6,114,598, 6,130,364, 6,162,963 and 6,150,584. See also WO 91/10741, WO 94/02602, WO 96/34096, WO 96/33735, WO 98/16654,
15 WO 98/24893, WO 98/50433, WO 99/45031, WO 99/53049, WO 00/09560, and WO 00/037504.

[0156] In another aspect, the invention provides a method for making anti-M-CSF antibodies from non-human, non-mouse animals by immunizing non-human transgenic animals that comprise human immunoglobulin loci with a M-CSF
20 antigen. One can produce such animals using the methods described in the above-cited documents. The methods disclosed in these documents can be modified as described in U.S. Patent 5,994,619. U.S. Patent 5,994,619 describes methods for producing novel cultural inner cell mass (CICM) cells and cell lines, derived from pigs and cows, and transgenic CICM cells into which heterologous DNA has been
25 inserted. CICM transgenic cells can be used to produce cloned transgenic embryos, fetuses, and offspring. The '619 patent also describes the methods of producing the transgenic animals, that are capable of transmitting the heterologous DNA to their progeny. In preferred embodiments, the non-human animals are rats, sheep, pigs, goats, cattle or horses.

[0157] XENOMOUSE™ mice produce an adult-like human repertoire of fully
30 human antibodies and generate antigen-specific human antibodies. In some embodiments, the XENOMOUSE™ mice contain approximately 80% of the

human antibody V gene repertoire through introduction of megabase sized, germline configuration yeast artificial chromosome (YAC) fragments of the human heavy chain loci and kappa light chain loci. In other embodiments, XENOMOUSE™ mice further contain approximately all of the lambda light chain locus. See Mendez *et al.*, *Nature Genetics* 15:146-156 (1997), Green and Jakobovits, *J. Exp. Med.* 188:483-495 (1998), and WO 98/24893, the disclosures of which are hereby incorporated by reference.

[0158] In some embodiments, the non-human animal comprising human immunoglobulin genes are animals that have a human immunoglobulin “minilocus”. In the minilocus approach, an exogenous Ig locus is mimicked through the inclusion of individual genes from the Ig locus. Thus, one or more V_H genes, one or more D_H genes, one or more J_H genes, a mu constant domain, and a second constant domain (preferably a gamma constant domain) are formed into a construct for insertion into an animal. This approach is described, *inter alia*, in U.S. Patent Nos. 5,545,807, 5,545,806, 5,569,825, 5,625,126, 5,633,425, 5,661,016, 5,770,429, 5,789,650, 5,814,318, 5,591,669, 5,612,205, 5,721,367, 5,789,215, and 5,643,763, hereby incorporated by reference.

[0159] In another aspect, the invention provides a method for making humanized anti-M-CSF antibodies. In some embodiments, non-human animals are immunized with a M-CSF antigen as described below under conditions that permit antibody production. Antibody-producing cells are isolated from the animals, fused with myelomas to produce hybridomas, and nucleic acids encoding the heavy and light chains of an anti-M-CSF antibody of interest are isolated. These nucleic acids are subsequently engineered using techniques known to those of skill in the art and as described further below to reduce the amount of non-human sequence, i.e., to humanize the antibody to reduce the immune response in humans

[0160] In some embodiments, the M-CSF antigen is isolated and/or purified M-CSF. In a preferred embodiment, the M-CSF antigen is human M-CSF. In some embodiments, the M-CSF antigen is a fragment of M-CSF. In some embodiments, the M-CSF fragment is the extracellular domain of M-CSF. In some embodiments, the M-CSF fragment comprises at least one epitope of M-CSF. In other embodiments, the M-CSF antigen is a cell that expresses or overexpresses M-

CSF or an immunogenic fragment thereof on its surface. In some embodiments, the M-CSF antigen is a M-CSF fusion protein. M-CSF can be purified from natural sources using known techniques. Recombinant M-CSF is commercially available.

- 5 [0161] Immunization of animals can be by any method known in the art. See, e.g., Harlow and Lane, *Antibodies: A Laboratory Manual*, New York: Cold Spring Harbor Press, 1990. Methods for immunizing non-human animals such as mice, rats, sheep, goats, pigs, cattle and horses are well known in the art. See, e.g., Harlow and Lane, *supra*, and U.S. Patent 5,994,619. In a preferred embodiment,
- 10 the M-CSF antigen is administered with an adjuvant to stimulate the immune response. Exemplary adjuvants include complete or incomplete Freund's adjuvant, RIBI (muramyl dipeptides) or ISCOM (immunostimulating complexes). Such adjuvants may protect the polypeptide from rapid dispersal by sequestering it in a local deposit, or they may contain substances that stimulate the host to secrete
- 15 factors that are chemotactic for macrophages and other components of the immune system. Preferably, if a polypeptide is being administered, the immunization schedule will involve two or more administrations of the polypeptide, spread out over several weeks. Example I exemplifies a method for producing anti-M-CSF monoclonal antibodies in XENOMOUSE™ mice.

20 *Production of Antibodies and Antibody-Producing Cell Lines*

- [0162] After immunization of an animal with a M-CSF antigen, antibodies and/or antibody-producing cells can be obtained from the animal. In some embodiments, anti-M-CSF antibody-containing serum is obtained from the animal by bleeding or sacrificing the animal. The serum may be used as it is obtained from the animal,
- 25 an immunoglobulin fraction may be obtained from the serum, or the anti-M-CSF antibodies may be purified from the serum.

- [0163] In some embodiments, antibody-producing immortalized cell lines are prepared from cells isolated from the immunized animal. After immunization, the animal is sacrificed and lymph node and/or splenic B cells are immortalized.
- 30 Methods of immortalizing cells include, but are not limited to, transfecting them with oncogenes, infecting them with an oncogenic virus, cultivating them under conditions that select for immortalized cells, subjecting them to carcinogenic or

- mutating compounds, fusing them with an immortalized cell, e.g., a myeloma cell, and inactivating a tumor suppressor gene. See, e.g., Harlow and Lane, *supra*. If fusion with myeloma cells is used, the myeloma cells preferably do not secrete immunoglobulin polypeptides (a non-secretory cell line). Immortalized cells are
- 5 screened using M-CSF, a portion thereof, or a cell expressing M-CSF. In a preferred embodiment, the initial screening is performed using an enzyme-linked immunoassay (ELISA) or a radioimmunoassay. An example of ELISA screening is provided in WO 00/37504, incorporated herein by reference.
- [0164] Anti-M-CSF antibody-producing cells, e.g., hybridomas, are selected,
- 10 cloned and further screened for desirable characteristics, including robust growth, high antibody production and desirable antibody characteristics, as discussed further below. Hybridomas can be expanded *in vivo* in syngeneic animals, in animals that lack an immune system, e.g., nude mice, or in cell culture *in vitro*. Methods of selecting, cloning and expanding hybridomas are well known to those
- 15 of ordinary skill in the art.
- [0165] In a preferred embodiment, the immunized animal is a non-human animal that expresses human immunoglobulin genes and the splenic B cells are fused to a myeloma cell line from the same species as the non-human animal. In a more preferred embodiment, the immunized animal is a XENOMOUSE™ animal and
- 20 the myeloma cell line is a non-secretory mouse myeloma. In an even more preferred embodiment, the myeloma cell line is P3-X63-AG8-653. See, e.g., Example I.
- [0166] Thus, in one embodiment, the invention provides methods of producing a cell line that produces a human monoclonal antibody or a fragment thereof directed
- 25 to M-CSF comprising (a) immunizing a non-human transgenic animal described herein with M-CSF, a portion of M-CSF or a cell or tissue expressing M-CSF; (b) allowing the transgenic animal to mount an immune response to M-CSF; (c) isolating B lymphocytes from a transgenic animal; (d) immortalizing the B lymphocytes; (e) creating individual monoclonal populations of the immortalized
- 30 B lymphocytes; and (f) screening the immortalized B lymphocytes to identify an antibody directed to M-CSF.

- [0167] In another aspect, the invention provides hybridomas that produce an human anti-M-CSF antibody. In a preferred embodiment, the hybridomas are mouse hybridomas, as described above. In other embodiments, the hybridomas are produced in a non-human, non-mouse species such as rats, sheep, pigs, goats, cattle or horses. In another embodiment, the hybridomas are human hybridomas.
- [0168] In another preferred embodiment, a transgenic animal is immunized with M-CSF, primary cells, e.g., spleen or peripheral blood cells, are isolated from an immunized transgenic animal and individual cells producing antibodies specific for the desired antigen are identified. Polyadenylated mRNA from each individual cell is isolated and reverse transcription polymerase chain reaction (RT-PCR) is performed using sense primers that anneal to variable region sequences, e.g., degenerate primers that recognize most or all of the FR1 regions of human heavy and light chain variable region genes and antisense primers that anneal to constant or joining region sequences. cDNAs of the heavy and light chain variable regions are then cloned and expressed in any suitable host cell, e.g., a myeloma cell, as chimeric antibodies with respective immunoglobulin constant regions, such as the heavy chain and κ or λ constant domains. See Babcook, J.S. *et al.*, *Proc. Natl. Acad. Sci. USA* 93:7843-48, 1996, herein incorporated by reference. Anti M-CSF antibodies may then be identified and isolated as described herein.
- [0169] In another embodiment, phage display techniques can be used to provide libraries containing a repertoire of antibodies with varying affinities for M-CSF. For production of such repertoires, it is unnecessary to immortalize the B cells from the immunized animal. Rather, the primary B cells can be used directly as a source of DNA. The mixture of cDNAs obtained from B cell, e.g., derived from spleens, is used to prepare an expression library, for example, a phage display library transfected into *E.coli*. The resulting cells are tested for immunoreactivity to M-CSF. Techniques for the identification of high affinity human antibodies from such libraries are described by Griffiths *et al.*, *EMBO J.*, 13:3245-3260 (1994); Nissim *et al.*, *ibid*, pp. 692-698 and by Griffiths *et al.*, *ibid*, 12:725-734.
- Ultimately, clones from the library are identified which produce binding affinities of a desired magnitude for the antigen and the DNA encoding the product responsible for such binding is recovered and manipulated for standard

recombinant expression. Phage display libraries may also be constructed using previously manipulated nucleotide sequences and screened in a similar fashion. In general, the cDNAs encoding heavy and light chains are independently supplied or linked to form Fv analogs for production in the phage library.

- 5 [0170] The phage library is then screened for the antibodies with the highest affinities for M-CSF and the genetic material recovered from the appropriate clone. Further rounds of screening can increase affinity of the original antibody isolated.

- [0171] In another aspect, the invention provides hybridomas that produce an human anti-M-CSF antibody. In a preferred embodiment, the hybridomas are
10 mouse hybridomas, as described above. In other embodiments, the hybridomas are produced in a non-human, non-mouse species such as rats, sheep, pigs, goats, cattle or horses. In another embodiment, the hybridomas are human hybridomas.

Nucleic Acids, Vectors, Host Cells, and
Recombinant Methods of Making Antibodies

- 15 *Nucleic Acids*

- [0172] The present invention also encompasses nucleic acid molecules encoding anti-M-CSF antibodies. In some embodiments, different nucleic acid molecules encode a heavy chain and a light chain of an anti-M-CSF immunoglobulin. In other embodiments, the same nucleic acid molecule encodes a heavy chain an a
20 light chain of an anti-M-CSF immunoglobulin. In one embodiment, the nucleic acid encodes a M-CSF antibody of the invention.

[0173] In some embodiments, the nucleic acid molecule encoding the variable domain of the light chain comprises a human V_L L5, O12, L2, B3, A27 gene and a $J\kappa 1$, $J\kappa 2$, $J\kappa 3$, or $J\kappa 4$ gene.

- 25 [0174] In some embodiments, the nucleic acid molecule encoding the light chain, encodes an amino acid sequence comprising 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 mutations from the germline amino acid sequence. In some embodiments, the nucleic acid molecule comprises a nucleotide sequence that encodes a V_L amino acid sequence comprising 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 non-conservative amino acid substitutions
30 and/or 1, 2, or 3 non-conservative substitutions compared to germline sequence. Substitutions may be in the CDR regions, the framework regions, or in the constant domain.

- [0175] In some embodiments, the nucleic acid molecule encodes a V_L amino acid sequence comprising one or more variants compared to germline sequence that are identical to the variations found in the V_L of one of the antibodies 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 or 9.14.4G1.
- [0176] In some embodiments, the nucleic acid molecule encodes at least three amino acid mutations compared to the germline sequence found in the V_L of one of the antibodies 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4, 8.10.3, or 9.7.2.
- [0177] In some embodiments, the nucleic acid molecule comprises a nucleotide sequence that encodes the V_L amino acid sequence of monoclonal antibody 252 (SEQ ID NO: 4), 88 (SEQ ID NO: 8), 100 (SEQ ID NO: 12), 3.8.3 (SEQ ID NO: 16), 2.7.3 (SEQ ID NO: 20), 1.120.1 (SEQ ID NO: 24), 9.14.4I (SEQ ID NO: 28), 8.10.3F (SEQ ID NO: 32), 9.7.2IF (SEQ ID NO: 36), 9.14.4 (SEQ ID NO: 28), 8.10.3 (SEQ ID NO: 44), 9.7.2 (SEQ ID NO: 48), 9.7.2C-Ser (SEQ ID NO: 52), 9.14.4C-Ser (SEQ ID NO: 56), 8.10.3C-Ser (SEQ ID NO: 60), 8.10.3-CG2 (SEQ ID NO: 60), 9.7.2-CG2 (SEQ ID NO: 52), 9.7.2-CG4 (SEQ ID NO: 52), 9.14.4-CG2 (SEQ ID NO: 56), 9.14.4-CG4 (SEQ ID NO: 56), 9.14.4-Ser (SEQ ID NO: 28), 9.7.2-Ser (SEQ ID NO: 48), 8.10.3-Ser (SEQ ID NO: 44), 8.10.3-CG4 (SEQ ID NO: 60) 8.10.3FG1 (SEQ ID NO: 32) or 9.14.4G1 (SEQ ID NO: 28), or a portion thereof. In some embodiments, said portion comprises at least the CDR2 region. In some embodiments, the nucleic acid encodes the amino acid sequence of the light chain CDRs of said antibody. In some embodiments, said portion is a contiguous portion comprising CDR1-CDR3.
- [0178] In some embodiments, the nucleic acid molecule comprises a nucleotide sequence that encodes the light chain amino acid sequence of one of SEQ ID NOS: 4, 8, 12, 16, 20, 24, 28, 32, 36, 44, 48, 52, 56 or 60. In some preferred embodiments, the nucleic acid molecule comprises the light chain nucleotide sequence of SEQ ID NOS: 3, 7, 11, 27, 31, 35, 43 or 47, or a portion thereof.
- [0179] In some embodiments, the nucleic acid molecule encodes a V_L amino acid sequence that is at least 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98% or 99% identical to a V_L amino acid sequence shown in Figure 1 or to a V_L amino acid

- sequences of any one of antibodies 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 or 9.14.4G1, or an amino acid sequence of
- 5 any one of SEQ ID NOS: 4, 8, 12, 16, 20, 24, 28, 32, 36, 44, 48, 52, 56 or 60.
- Nucleic acid molecules of the invention include nucleic acids that hybridize under highly stringent conditions, such as those described above, to a nucleic acid sequence encoding the light chain amino acid sequence of SEQ ID NOS: 4, 8, 12, 16, 20, 24, 28, 32, 36, 44, 48, 52, 56 or 60, or that has the light chain nucleic acid
- 10 sequence of SEQ ID NOS: 3, 7, 11, 27, 31, 35, 43 or 47.
- [0180] In another embodiment, the nucleic acid encodes a full-length light chain of an antibody selected from 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser,
- 15 8.10.3-CG4, 8.10.3FG1 or 9.14.4G1, or a light chain comprising the amino acid sequence of SEQ ID NOS: 4, 8, 12, 16, 20, 24, 28, 32, 36, 44, 48, 52, 56 or 60 and a constant region of a light chain, or a light chain comprising a mutation. Further, the nucleic acid may comprise the light chain nucleotide sequence of SEQ ID NOS: 3, 7, 11, 27, 31, 35, 43 or 47 and the nucleotide sequence encoding a
- 20 constant region of a light chain, or a nucleic acid molecule encoding a light chain comprise a mutation.
- [0181] In another preferred embodiment, the nucleic acid molecule encodes the variable domain of the heavy chain (V_H) that comprises a human V_H 1-18, 3-33, 3-11, 3-23, 3-48, or 3-7 gene sequence or a sequence derived therefrom. In various
- 25 embodiments, the nucleic acid molecule comprises a human V_H 1-18 gene, a D_H 4-23 gene and a human J_H 4 gene; a human V_H 3-33 gene, a human D_H 1-26 gene and a human J_H 4 gene; a human V_H 3-11 gene, a human D_H 7-27 gene and a human J_H 6 gene; a human V_H 3-11 gene, a human D_H 7-27 gene and a human J_H 6 gene; a human V_H 3-23 gene, a human D_H 1-26 gene and a human J_H 4 gene; a human V_H
- 30 3-7 gene, a human D_H 6-13 gene and a human J_H 4 gene; a human V_H 3-11 gene, a human D_H 7-27 gene, and a human J_H 4b gene; a human V_H 3-48 gene, a human

D_H1-26 gene, and a human J_H4b gene; a human V_H3-11 gene, a human D_H6-13 gene, and a human J_H6b gene, or a sequence derived from the human genes.

[0182] In some embodiments, the nucleic acid molecule encodes an amino acid sequence comprising 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17 or 18

- 5 mutations compared to the germline amino acid sequence of the human V, D or J genes. In some embodiments, said mutations are in the V_H region. In some embodiments, said mutations are in the CDR regions.

- [0183] In some embodiments, the nucleic acid molecule encodes one or more amino acid mutations compared to the germline sequence that are identical to
- 10 amino acid mutations found in the V_H of monoclonal antibody 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 or 9.14.4G1. In some
- 15 embodiments, the nucleic acid encodes at least three amino acid mutations compared to the germline sequences that are identical to at least three amino acid mutations found in one of the above-listed monoclonal antibodies.

- [0184] In some embodiments, the nucleic acid molecule comprises a nucleotide sequence that encodes at least a portion of the V_H amino acid sequence of antibody
- 20 252 (SEQ ID NO: 4), 88 (SEQ ID NO: 8), 100 (SEQ ID NO: 12), 3.8.3 (SEQ ID NO: 16), 2.7.3 (SEQ ID NO: 20), 1.120.1 (SEQ ID NO: 24), 9.14.4I (SEQ ID NO: 28), 8.10.3F (SEQ ID NO: 32), 9.7.2IF (SEQ ID NO: 36), 9.14.4 (SEQ ID NO: 28), 8.10.3 (SEQ ID NO: 44), 9.7.2 (SEQ ID NO: 48), 9.7.2C-Ser (SEQ ID NO: 52), 9.14.4C-Ser (SEQ ID NO: 56), 8.10.3C-Ser (SEQ ID NO: 60), 8.10.3-CG2 (SEQ ID NO: 60), 9.7.2-CG2 (SEQ ID NO: 52), 9.7.2-CG4 (SEQ ID NO: 52),
- 25 9.14.4-CG2 (SEQ ID NO: 56), 9.14.4-CG4 (SEQ ID NO: 56), 9.14.4-Ser (SEQ ID NO: 28), 9.7.2-Ser (SEQ ID NO: 48), 8.10.3-Ser (SEQ ID NO: 44), 8.10.3-CG4 (SEQ ID NO: 60) 8.10.3FG1 (SEQ ID NO: 32) or 9.14.4G1 (SEQ ID NO: 28), or said sequence having conservative amino acid mutations and/or a total of three or fewer non-conservative amino acid substitutions. In various embodiments the
- 30 sequence encodes one or more CDR regions, preferably a CDR3 region, all three CDR regions, a contiguous portion including CDR1-CDR3, or the entire V_H region.

[0185] In some embodiments, the nucleic acid molecule comprises a heavy chain nucleotide sequence that encodes the amino acid sequence of one of SEQ ID NOS: 2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 46, 50, 54, 58, 62, 66, 70, 74, 78, 82, 86, 90, 94, 98 or 102. In some preferred embodiments, the nucleic acid molecule comprises at least a portion of the heavy chain nucleotide sequence of SEQ ID NO: 1, 5, 9, 25, 29, 33, 37, 45, 97 or 101. In some embodiments, said portion encodes the V_H region, a CDR3 region, all three CDR regions, or a contiguous region including CDR1-CDR3.

[0186] In some embodiments, the nucleic acid molecule encodes a V_H amino acid sequence that is at least 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98% or 99% identical to the V_H amino acid sequences shown in Figure 4 or to a V_H amino acid sequence of any one of SEQ ID NOS: 2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 46, 50, 54, 58, 62, 66, 70, 74, 78, 82, 86, 90, 94, 98 or 102. Nucleic acid molecules of the invention include nucleic acids that hybridize under highly stringent conditions, such as those described above, to a nucleotide sequence encoding the heavy chain amino acid sequence of SEQ ID NOS: 2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 46, 50, 54, 58, 62, 66, 70, 74, 78, 82, 86, 90, 94, 98 or 102 or that has the nucleotide sequence of SEQ ID NOS: 1, 5, 9, 25, 29, 33, 37, 45, 97 or 101.

[0187] In another embodiment, the nucleic acid encodes a full-length heavy chain of an antibody selected from 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 or 9.14.4G1, or a heavy chain having the amino acid sequence of SEQ ID NOS: 2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 46, 50, 54, 58, 62, 66, 70, 74, 78, 82, 86, 90, 94, 98 or 102 and a constant region of a heavy chain, or a heavy chain comprising a mutation. Further, the nucleic acid may comprise the heavy chain nucleotide sequence of SEQ ID NOS: 1, 5, 9, 25, 29, 33, 37, 45, 97 or 101 and a nucleotide sequence encoding a constant region of a light chain, or a nucleic acid molecule encoding a heavy chain comprising a mutation.

[0188] A nucleic acid molecule encoding the heavy or entire light chain of an anti-M-CSF antibody or portions thereof can be isolated from any source that

produces such antibody. In various embodiments, the nucleic acid molecules are isolated from a B cell isolated from an animal immunized with M-CSF or from an immortalized cell derived from such a B cell that expresses an anti-M-CSF antibody. Methods of isolating mRNA encoding an antibody are well-known in the art. See, e.g., Sambrook *et al.* The mRNA may be used to produce cDNA for use in the polymerase chain reaction (PCR) or cDNA cloning of antibody genes.

In a preferred embodiment, the nucleic acid molecule is isolated from a hybridoma that has as one of its fusion partners a human immunoglobulin-producing cell from a non-human transgenic animal. In an even more preferred embodiment, the human immunoglobulin producing cell is isolated from a XENOMOUSETM animal. In another embodiment, the human immunoglobulin-producing cell is from a non-human, non-mouse transgenic animal, as described above. In another embodiment, the nucleic acid is isolated from a non-human, non-transgenic animal. The nucleic acid molecules isolated from a non-human, non-transgenic animal may be used, e.g., for humanized antibodies.

[0189] In some embodiments, a nucleic acid encoding a heavy chain of an anti-M-CSF antibody of the invention can comprise a nucleotide sequence encoding a V_H domain of the invention joined in-frame to a nucleotide sequence encoding a heavy chain constant domain from any source. Similarly, a nucleic acid molecule encoding a light chain of an anti-M-CSF antibody of the invention can comprise a nucleotide sequence encoding a V_L domain of the invention joined in-frame to a nucleotide sequence encoding a light chain constant domain from any source.

[0190] In a further aspect of the invention, nucleic acid molecules encoding the variable domain of the heavy (V_H) and light (V_L) chains are "converted" to full-length antibody genes. In one embodiment, nucleic acid molecules encoding the V_H or V_L domains are converted to full-length antibody genes by insertion into an expression vector already encoding heavy chain constant (C_H) or light chain (C_L) constant domains, respectively, such that the V_H segment is operatively linked to the C_H segment(s) within the vector, and the V_L segment is operatively linked to the C_L segment within the vector. In another embodiment, nucleic acid molecules encoding the V_H and/or V_L domains are converted into full-length antibody genes by linking, e.g., ligating, a nucleic acid molecule encoding a V_H and/or V_L domains

to a nucleic acid molecule encoding a C_H and/or C_L domain using standard molecular biological techniques. Nucleic acid sequences of human heavy and light chain immunoglobulin constant domain genes are known in the art. See, e.g., Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed., NIH Publ.

- 5 No. 91-3242, 1991. Nucleic acid molecules encoding the full-length heavy and/or light chains may then be expressed from a cell into which they have been introduced and the anti-M-CSF antibody isolated.

[0191] The nucleic acid molecules may be used to recombinantly express large quantities of anti-M-CSF antibodies. The nucleic acid molecules also may be used
10 to produce chimeric antibodies, bispecific antibodies, single chain antibodies, immunoadhesins, diabodies, mutated antibodies and antibody derivatives, as described further below. If the nucleic acid molecules are derived from a non-human, non-transgenic animal, the nucleic acid molecules may be used for antibody humanization, also as described below.

- 15 [0192] In another embodiment, a nucleic acid molecule of the invention is used as a probe or PCR primer for a specific antibody sequence. For instance, the nucleic acid can be used as a probe in diagnostic methods or as a PCR primer to amplify regions of DNA that could be used, inter alia, to isolate additional nucleic acid molecules encoding variable domains of anti-M-CSF antibodies. In some
20 embodiments, the nucleic acid molecules are oligonucleotides. In some embodiments, the oligonucleotides are from highly variable regions of the heavy and light chains of the antibody of interest. In some embodiments, the oligonucleotides encode all or a part of one or more of the CDRs of antibody 252, 88, 100, 3.8.3, 2.7.3, or 1.120.1, or variants thereof described herein.

25

Vectors

- [0193] The invention provides vectors comprising nucleic acid molecules that encode the heavy chain of an anti-M-CSF antibody of the invention or an antigen-binding portion thereof. The invention also provides vectors comprising nucleic
30 acid molecules that encode the light chain of such antibodies or antigen-binding portion thereof. The invention further provides vectors comprising nucleic acid

molecules encoding fusion proteins, modified antibodies, antibody fragments, and probes thereof.

- [0194] In some embodiments, the anti-M-CSF antibodies, or antigen-binding portions of the invention are expressed by inserting DNAs encoding partial or full-length light and heavy chains, obtained as described above, into expression vectors such that the genes are operatively linked to necessary expression control sequences such as transcriptional and transnational control sequences. Expression vectors include plasmids, retroviruses, adenoviruses, adeno-associated viruses (AAV), plant viruses such as cauliflower mosaic virus, tobacco mosaic virus, cosmids, YACs, EBV derived episomes, and the like. The antibody gene is ligated into a vector such that transcriptional and transnational control sequences within the vector serve their intended function of regulating the transcription and translation of the antibody gene. The expression vector and expression control sequences are chosen to be compatible with the expression host cell used. The antibody light chain gene and the antibody heavy chain gene can be inserted into separate vectors. In a preferred embodiment, both genes are inserted into the same expression vector. The antibody genes are inserted into the expression vector by standard methods (e.g., ligation of complementary restriction sites on the antibody gene fragment and vector, or blunt end ligation if no restriction sites are present).
- [0195] A convenient vector is one that encodes a functionally complete human C_H or C_L immunoglobulin sequence, with appropriate restriction sites engineered so that any V_H or V_L sequence can easily be inserted and expressed, as described above. In such vectors, splicing usually occurs between the splice donor site in the inserted J region and the splice acceptor site preceding the human C domain, and also at the splice regions that occur within the human C_H exons. Polyadenylation and transcription termination occur at native chromosomal sites downstream of the coding regions. The recombinant expression vector also can encode a signal peptide that facilitates secretion of the antibody chain from a host cell. The antibody chain gene may be cloned into the vector such that the signal peptide is linked in-frame to the amino terminus of the immunoglobulin chain. The signal peptide can be an immunoglobulin signal peptide or a heterologous signal peptide (i.e., a signal peptide from a non-immunoglobulin protein).

- [0196] In addition to the antibody chain genes, the recombinant expression vectors of the invention carry regulatory sequences that control the expression of the antibody chain genes in a host cell. It will be appreciated by those skilled in the art that the design of the expression vector, including the selection of regulatory sequences may depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, etc. Preferred regulatory sequences for mammalian host cell expression include viral elements that direct high levels of protein expression in mammalian cells, such as promoters and/or enhancers derived from retroviral LTRs, cytomegalovirus (CMV) (such as the CMV promoter/enhancer), Simian Virus 40 (SV40) (such as the SV40 promoter/enhancer), adenovirus, (e.g., the adenovirus major late promoter (AdMLP)), polyoma and strong mammalian promoters such as native immunoglobulin and actin promoters. For further description of viral regulatory elements, and sequences thereof, see e.g., U.S. Patent No. 5,168,062, U.S. Patent No. 4,510,245 and U.S. Patent No. 4,968,615. Methods for expressing antibodies in plants, including a description of promoters and vectors, as well as transformation of plants is known in the art. See, e.g., United States Patents 6,517,529, herein incorporated by reference. Methods of expressing polypeptides in bacterial cells or fungal cells, e.g., yeast cells, are also well known in the art.
- [0197] In addition to the antibody chain genes and regulatory sequences, the recombinant expression vectors of the invention may carry additional sequences, such as sequences that regulate replication of the vector in host cells (e.g., origins of replication) and selectable marker genes. The selectable marker gene facilitates selection of host cells into which the vector has been introduced (see e.g., U.S. Patent Nos. 4,399,216, 4,634,665 and 5,179,017). For example, typically the selectable marker gene confers resistance to drugs, such as G418, hygromycin or methotrexate, on a host cell into which the vector has been introduced. Preferred selectable marker genes include the dihydrofolate reductase (DHFR) gene (for use in dhfr-host cells with methotrexate selection/amplification), the neomycin resistance gene (for G418 selection), and the glutamate synthetase gene.

- [0198] Nucleic acid molecules encoding anti-M-CSF antibodies and vectors comprising these nucleic acid molecules can be used for transfection of a suitable mammalian, plant, bacterial or yeast host cell. Transformation can be by any known method for introducing polynucleotides into a host cell. Methods for
- 5 introduction of heterologous polynucleotides into mammalian cells are well known in the art and include dextran-mediated transfection, calcium phosphate precipitation, polybrene-mediated transfection, protoplast fusion, electroporation, encapsulation of the polynucleotide(s) in liposomes, and direct microinjection of the DNA into nuclei. In addition, nucleic acid molecules may be introduced into
- 10 mammalian cells by viral vectors. Methods of transforming cells are well known in the art. See, e.g., U.S. Patent Nos. 4,399,216, 4,912,040, 4,740,461, and 4,959,455 (which patents are hereby incorporated herein by reference). Methods of transforming plant cells are well known in the art, including, e.g.,
- 15 *Agrobacterium*-mediated transformation, biolistic transformation, direct injection, electroporation and viral transformation. Methods of transforming bacterial and yeast cells are also well known in the art.
- [0199] Mammalian cell lines available as hosts for expression are well known in the art and include many immortalized cell lines available from the American Type Culture Collection (ATCC). These include, *inter alia*, Chinese hamster ovary
- 20 (CHO) cells, NSO, SP2 cells, HeLa cells, baby hamster kidney (BHK) cells, monkey kidney cells (COS), human hepatocellular carcinoma cells (e.g., Hep G2), A549 cells, and a number of other cell lines. Cell lines of particular preference are selected through determining which cell lines have high expression levels. Other cell lines that may be used are insect cell lines, such as Sf9 cells. When
- 25 recombinant expression vectors encoding antibody genes are introduced into mammalian host cells, the antibodies are produced by culturing the host cells for a period of time sufficient to allow for expression of the antibody in the host cells or, more preferably, secretion of the antibody into the culture medium in which the host cells are grown. Antibodies can be recovered from the culture medium using
- 30 standard protein purification methods. Plant host cells include, e.g., *Nicotiana*, *Arabidopsis*, duckweed, corn, wheat, potato, etc. Bacterial host cells include

E. coli and *Streptomyces* species. Yeast host cells include *Schizosaccharomyces pombe*, *Saccharomyces cerevisiae* and *Pichia pastoris*.

- [0200] Further, expression of antibodies of the invention (or other moieties therefrom) from production cell lines can be enhanced using a number of known techniques. For example, the glutamine synthetase gene expression system (the GS system) is a common approach for enhancing expression under certain conditions. The GS system is discussed in whole or part in connection with European Patent Nos. 0 216 846, 0 256 055, and 0 323 997 and European Patent Application No. 89303964.4.
- 10 [0201] It is possible that antibodies expressed by different cell lines or in transgenic animals will have different glycosylation from each other. However, all antibodies encoded by the nucleic acid molecules provided herein, or comprising the amino acid sequences provided herein are part of the instant invention, regardless of the glycosylation state or pattern or modification of the antibodies.

15

Transgenic Animals and Plants

- [0202] Anti-M-CSF antibodies of the invention also can be produced transgenically through the generation of a mammal or plant that is transgenic for the immunoglobulin heavy and light chain sequences of interest and production of the antibody in a recoverable form therefrom. In connection with the transgenic production in mammals, anti-M-CSF antibodies can be produced in, and recovered from, the milk of goats, cows, or other mammals. See, e.g., U.S. Patent Nos. 5,827,690, 5,756,687, 5,750,172, and 5,741,957. In some embodiments, non-human transgenic animals that comprise human immunoglobulin loci are
- 20 immunized with M-CSF or an immunogenic portion thereof, as described above. Methods for making antibodies in plants, yeast or fungi/algae are described, e.g., in US patents 6,046,037 and US 5,959,177.
- [0203] In some embodiments, non-human transgenic animals or plants are produced by introducing one or more nucleic acid molecules encoding an anti-M-
- 30 CSF antibody of the invention into the animal or plant by standard transgenic techniques. See Hogan and United States Patent 6,417,429, *supra*. The transgenic cells used for making the transgenic animal can be embryonic stem cells or somatic

- cells. The transgenic non-human organisms can be chimeric, nonchimeric heterozygotes, and nonchimeric homozygotes. See, e.g., Hogan *et al.*, *Manipulating the Mouse Embryo: A Laboratory Manual* 2ed., Cold Spring Harbor Press (1999); Jackson *et al.*, *Mouse Genetics and Transgenics: A Practical Approach*, Oxford University Press (2000); and Pinkert, *Transgenic Animal Technology: A Laboratory Handbook*, Academic Press (1999). In some embodiments, the transgenic non-human animals have a targeted disruption and replacement by a targeting construct that encodes a heavy chain and/or a light chain of interest. In a preferred embodiment, the transgenic animals comprise and
- 10 express nucleic acid molecules encoding heavy and light chains that specifically bind to M-CSF, preferably human M-CSF. In some embodiments, the transgenic animals comprise nucleic acid molecules encoding a modified antibody such as a single-chain antibody, a chimeric antibody or a humanized antibody. The anti-M-CSF antibodies may be made in any transgenic animal. In a preferred
- 15 embodiment, the non-human animals are mice, rats, sheep, pigs, goats, cattle or horses. The non-human transgenic animal expresses said encoded polypeptides in blood, milk, urine, saliva, tears, mucus and other bodily fluids.

Phage Display Libraries

- 20 [0204] The invention provides a method for producing an anti-M-CSF antibody or antigen-binding portion thereof comprising the steps of synthesizing a library of human antibodies on phage, screening the library with M-CSF or a portion thereof, isolating phage that bind M-CSF, and obtaining the antibody from the phage. By way of example, one method for preparing the library of antibodies for use in
- 25 phage display techniques comprises the steps of immunizing a non-human animal comprising human immunoglobulin loci with M-CSF or an antigenic portion thereof to create an immune response, extracting antibody producing cells from the immunized animal; isolating RNA from the extracted cells, reverse transcribing the RNA to produce cDNA, amplifying the cDNA using a primer, and inserting the
- 30 cDNA into a phage display vector such that antibodies are expressed on the phage. Recombinant anti-M-CSF antibodies of the invention may be obtained in this way.

- [0205] Recombinant anti-M-CSF human antibodies of the invention can be isolated by screening a recombinant combinatorial antibody library. Preferably the library is a scFv phage display library, generated using human V_L and V_H cDNAs prepared from mRNA isolated from B cells. Methodologies for preparing and screening such libraries are known in the art. There are commercially available kits for generating phage display libraries (e.g., the Pharmacia Recombinant Phage Antibody System, catalog no. 27-9400-01; and the Stratagene SurfZAP[™] phage display kit, catalog no. 240612). There also are other methods and reagents that can be used in generating and screening antibody display libraries (see, e.g., U.S. Patent No. 5,223,409; PCT Publication Nos. WO 92/18619, WO 91/17271, WO 92/20791, WO 92/15679, WO 93/01288, WO 92/01047, WO 92/09690; Fuchs *et al.*, *Bio/Technology* 9:1370-1372 (1991); Hay *et al.*, *Hum. Antibod. Hybridomas* 3:81-85 (1992); Huse *et al.*, *Science* 246:1275-1281 (1989); McCafferty *et al.*, *Nature* 348:552-554 (1990); Griffiths *et al.*, *EMBO J.* 12:725-734 (1993); Hawkins *et al.*, *J. Mol. Biol.* 226:889-896 (1992); Clackson *et al.*, *Nature* 352:624-628 (1991); Gram *et al.*, *Proc. Natl. Acad. Sci. USA* 89:3576-3580 (1992); Garrad *et al.*, *Bio/Technology* 9:1373-1377 (1991); Hoogenboom *et al.*, *Nuc. Acid Res.* 19:4133-4137 (1991); and Barbas *et al.*, *Proc. Natl. Acad. Sci. USA* 88:7978-7982 (1991).
- [0206] In one embodiment, to isolate a human anti-M-CSF antibodies with the desired characteristics, a human anti-M-CSF antibody as described herein is first used to select human heavy and light chain sequences having similar binding activity toward M-CSF, using the epitope imprinting methods described in PCT Publication No. WO 93/06213. The antibody libraries used in this method are preferably scFv libraries prepared and screened as described in PCT Publication No. WO 92/01047, McCafferty *et al.*, *Nature* 348:552-554 (1990); and Griffiths *et al.*, *EMBO J.* 12:725-734 (1993). The scFv antibody libraries preferably are screened using human M-CSF as the antigen.
- [0207] Once initial human V_L and V_H domains are selected, "mix and match" experiments are performed, in which different pairs of the initially selected V_L and V_H segments are screened for M-CSF binding to select preferred V_L/V_H pair combinations. Additionally, to further improve the quality of the antibody, the V_L

- and V_H segments of the preferred V_L/V_H pair(s) can be randomly mutated, preferably within the CDR3 region of V_H and/or V_L , in a process analogous to the *in vivo* somatic mutation process responsible for affinity maturation of antibodies during a natural immune response. This *in vitro* affinity maturation can be
- 5 accomplished by amplifying V_H and V_L domains using PCR primers complimentary to the V_H CDR3 or V_L CDR3, respectively, which primers have been "spiked" with a random mixture of the four nucleotide bases at certain positions such that the resultant PCR products encode V_H and V_L segments into which random mutations have been introduced into the V_H and/or V_L CDR3
- 10 regions. These randomly mutated V_H and V_L segments can be re-screened for binding to M-CSF.
- [0208] Following screening and isolation of an anti-M-CSF antibody of the invention from a recombinant immunoglobulin display library, nucleic acids encoding the selected antibody can be recovered from the display package (e.g.,
- 15 from the phage genome) and subcloned into other expression vectors by standard recombinant DNA techniques. If desired, the nucleic acid can further be manipulated to create other antibody forms of the invention, as described below. To express a recombinant human antibody isolated by screening of a combinatorial library, the DNA encoding the antibody is cloned into a recombinant expression
- 20 vector and introduced into a mammalian host cells, as described above.

Class switching

- [0209] Another aspect of the invention provides a method for converting the class or subclass of an anti-M-CSF antibody to another class or subclass. In some
- 25 embodiments, a nucleic acid molecule encoding a V_L or V_H that does not include any nucleic acid sequences encoding C_L or C_H is isolated using methods well-known in the art. The nucleic acid molecule then is operatively linked to a nucleic acid sequence encoding a C_L or C_H from a desired immunoglobulin class or subclass. This can be achieved using a vector or nucleic acid molecule that
- 30 comprises a C_L or C_H chain, as described above. For example, an anti-M-CSF antibody that was originally IgM can be class switched to an IgG. Further, the class switching may be used to convert one IgG subclass to another, e.g., from

- IgG1 to IgG2. Another method for producing an antibody of the invention comprising a desired isotype comprises the steps of isolating a nucleic acid encoding a heavy chain of an anti-M-CSF antibody and a nucleic acid encoding a light chain of an anti-M-CSF antibody, isolating the sequence encoding the V_H region, ligating the V_H sequence to a sequence encoding a heavy chain constant domain of the desired isotype, expressing the light chain gene and the heavy chain construct in a cell, and collecting the anti-M-CSF antibody with the desired isotype.
- [0210] In some embodiments, anti-M-CSF antibodies of the invention have the serine at position 228 (according to the EU-numbering convention) of the heavy chain changed to a proline. Accordingly, the CPSC sub-sequence in the F_C region of IgG4 becomes CPPC, which is the sub-sequence in IgG1. (Aalberse, R.C. and Schuurman, J., *Immunology*, 105:9-19 (2002)). For example, the serine at residue 243 SEQ ID NO: 46 (which corresponds to residue 228 in the EU-numbering convention) would become proline. Similarly, the serine at residue 242 of SEQ ID NO: 38 (which corresponds to residue 228 in the EU-numbering convention) would become proline. In some embodiments, the framework region of the IgG4 antibody can be back-mutated to the germline framework sequence. Some embodiments comprise both the back-mutates framework region and the serine to proline change in the F_C region. See, e.g., SEQ ID NO: 54 (antibody 9.14.4C-Ser) and SEQ ID NO: 58 (antibody 8.10.3C-Ser) in Table 1.

Deimmunized Antibodies

- [0211] Another way of producing antibodies with reduced immunogenicity is the deimmunization of antibodies. In another aspect of the invention, the antibody may be deimmunized using the techniques described in, e.g., PCT Publication Nos. WO98/52976 and WO00/34317 (which incorporated herein by reference in their entirety).

Mutated Antibodies

- [0212] In another embodiment, the nucleic acid molecules, vectors and host cells may be used to make mutated anti-M-CSF antibodies. The antibodies may be

mutated in the variable domains of the heavy and/or light chains, e.g., to alter a binding property of the antibody. For example, a mutation may be made in one or more of the CDR regions to increase or decrease the K_D of the antibody for M-CSF, to increase or decrease k_{off} , or to alter the binding specificity of the

5 antibody. Techniques in site-directed mutagenesis are well-known in the art. See, e.g., Sambrook *et al.* and Ausubel *et al.*, *supra*. In a preferred embodiment, mutations are made at an amino acid residue that is known to be changed compared to germline in a variable domain of an anti-M-CSF antibody. In another embodiment, one or more mutations are made at an amino acid residue that is

10 known to be changed compared to the germline in a CDR region or framework region of a variable domain, or in a constant domain of a monoclonal antibody 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 or

15 9.14.4G1. In another embodiment, one or more mutations are made at an amino acid residue that is known to be changed compared to the germline in a CDR region or framework region of a variable domain of a heavy chain amino acid sequence selected from SEQ ID NOS: 2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 46, 50, 54, 58, 62, 66, 70, 74, 78, 82, 86, 90, 94, 98 or 102, or whose heavy chain

20 nucleotide sequence is presented in SEQ ID NOS: 1, 5, 9, 25, 29, 33, 37, 45, 97 or 101. In another embodiment, one or more mutations are made at an amino acid residue that is known to be changed compared to the germline in a CDR region or framework region of a variable domain of a light chain amino acid sequence selected from SEQ ID NOS: 4, 8, 12, 16, 20, 24, 28, 32, 36, 44, 48, 52, 56 or 60, or

25 whose light chain nucleotide sequence is presented in SEQ ID NOS: 3, 7, 11, 27, 31, 35, 43 or 47.

[0213] In one embodiment, the framework region is mutated so that the resulting framework region(s) have the amino acid sequence of the corresponding germline gene. A mutation may be made in a framework region or constant domain to

30 increase the half-life of the anti-M-CSF antibody. See, e.g., PCT Publication No. WO 00/09560, herein incorporated by reference. A mutation in a framework region or constant domain also can be made to alter the immunogenicity of the

antibody, to provide a site for covalent or non-covalent binding to another molecule, or to alter such properties as complement fixation, FcR binding and antibody-dependent cell-mediated cytotoxicity (ADCC). According to the invention, a single antibody may have mutations in any one or more of the CDRs or framework regions of the variable domain or in the constant domain.

- 5 [0214] In some embodiments, there are from 1 to 8 including any number in between, amino acid mutations in either the V_H or V_L domains of the mutated anti-M-CSF antibody compared to the anti-M-CSF antibody prior to mutation. In any of the above, the mutations may occur in one or more CDR regions. Further, any
- 10 of the mutations can be conservative amino acid substitutions. In some embodiments, there are no more than 5, 4, 3, 2, or 1 amino acid changes in the constant domains.

Modified Antibodies

- 15 [0215] In another embodiment, a fusion antibody or immunoadhesin may be made that comprises all or a portion of an anti-M-CSF antibody of the invention linked to another polypeptide. In a preferred embodiment, only the variable domains of the anti-M-CSF antibody are linked to the polypeptide. In another preferred embodiment, the V_H domain of an anti-M-CSF antibody is linked to a
- 20 first polypeptide, while the V_L domain of an anti-M-CSF antibody is linked to a second polypeptide that associates with the first polypeptide in a manner such that the V_H and V_L domains can interact with one another to form an antibody binding site. In another preferred embodiment, the V_H domain is separated from the V_L domain by a linker such that the V_H and V_L domains can interact with one another
- 25 (see below under Single Chain Antibodies). The V_H-linker-V_L antibody is then linked to the polypeptide of interest. The fusion antibody is useful for directing a polypeptide to a M-CSF-expressing cell or tissue. The polypeptide may be a therapeutic agent, such as a toxin, growth factor or other regulatory protein, or may be a diagnostic agent, such as an enzyme that may be easily visualized, such as
- 30 horseradish peroxidase. In addition, fusion antibodies can be created in which two (or more) single-chain antibodies are linked to one another. This is useful if one

wants to create a divalent or polyvalent antibody on a single polypeptide chain, or if one wants to create a bispecific antibody.

[0216] To create a single chain antibody, (scFv) the V_H - and V_L -encoding DNA fragments are operatively linked to another fragment encoding a flexible linker, e.g., encoding the amino acid sequence (Gly₄-Ser)₃, such that the V_H and V_L sequences can be expressed as a contiguous single-chain protein, with the V_L and V_H domains joined by the flexible linker. See, e.g., Bird *et al.*, *Science* 242:423-426 (1988); Huston *et al.*, *Proc. Natl. Acad. Sci. USA* 85:5879-5883 (1988); McCafferty *et al.*, *Nature* 348:552-554 (1990). The single chain antibody may be monovalent, if only a single V_H and V_L are used, bivalent, if two V_H and V_L are used, or polyvalent, if more than two V_H and V_L are used. Bispecific or polyvalent antibodies may be generated that bind specifically to M-CSF and to another molecule.

[0217] In other embodiments, other modified antibodies may be prepared using anti-M-CSF antibody-encoding nucleic acid molecules. For instance, "Kappa bodies" (Ill *et al.*, *Protein Eng.* 10: 949-57 (1997)), "Minibodies" (Martin *et al.*, *EMBO J.* 13: 5303-9 (1994)), "Diabodies" (Holliger *et al.*, *Proc. Natl. Acad. Sci. USA* 90: 6444-6448 (1993)), or "Janusins" (Traunecker *et al.*, *EMBO J.* 10:3655-3659 (1991) and Traunecker *et al.*, *Int. J. Cancer* (Suppl.) 7:51-52 (1992)) may be prepared using standard molecular biological techniques following the teachings of the specification.

[0218] Bispecific antibodies or antigen-binding fragments can be produced by a variety of methods including fusion of hybridomas or linking of Fab' fragments. See, e.g., Songsivilai & Lachmann, *Clin. Exp. Immunol.* 79: 315-321 (1990), Kostelny *et al.*, *J. Immunol.* 148:1547-1553 (1992). In addition, bispecific antibodies may be formed as "diabodies" or "Janusins." In some embodiments, the bispecific antibody binds to two different epitopes of M-CSF. In some embodiments, the bispecific antibody has a first heavy chain and a first light chain from monoclonal antibody 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, or 9.7.2 and an additional antibody heavy chain and light chain. In some embodiments, the additional light chain and heavy chain also are

from one of the above-identified monoclonal antibodies, but are different from the first heavy and light chains.

- [0219] In some embodiments, the modified antibodies described above are prepared using one or more of the variable domains or CDR regions from a human anti-M-CSF monoclonal antibody provided herein, from an amino acid sequence of said monoclonal antibody, or from a heavy chain or light chain encoded by a nucleic acid sequence encoding said monoclonal antibody.

Derivatized and Labeled Antibodies

- [0220] An anti-M-CSF antibody or antigen-binding portion of the invention can be derivatized or linked to another molecule (e.g., another peptide or protein). In general, the antibodies or portion thereof is derivatized such that the M-CSF binding is not affected adversely by the derivatization or labeling. Accordingly, the antibodies and antibody portions of the invention are intended to include both intact and modified forms of the human anti-M-CSF antibodies described herein. For example, an antibody or antibody portion of the invention can be functionally linked (by chemical coupling, genetic fusion, noncovalent association or otherwise) to one or more other molecular entities, such as another antibody (e.g., a bispecific antibody or a diabody), a detection agent, a cytotoxic agent, a pharmaceutical agent, and/or a protein or peptide that can mediate association of the antibody or antibody portion with another molecule (such as a streptavidin core region or a polyhistidine tag).

- [0221] One type of derivatized antibody is produced by crosslinking two or more antibodies (of the same type or of different types, e.g., to create bispecific antibodies). Suitable crosslinkers include those that are heterobifunctional, having two distinctly reactive groups separated by an appropriate spacer (e.g., m-maleimidobenzoyl-N-hydroxysuccinimide ester) or homobifunctional (e.g., disuccinimidyl suberate). Such linkers are available from Pierce Chemical Company, Rockford, Ill.

- [0222] Another type of derivatized antibody is a labeled antibody. Useful detection agents with which an antibody or antigen-binding portion of the invention may be derivatized include fluorescent compounds, including

- fluorescein, fluorescein isothiocyanate, rhodamine, 5-dimethylamine-1-naphthalenesulfonyl chloride, phycoerythrin, lanthanide phosphors and the like. An antibody can also be labeled with enzymes that are useful for detection, such as horseradish peroxidase, β -galactosidase, luciferase, alkaline phosphatase, glucose oxidase and the like. When an antibody is labeled with a detectable enzyme, it is detected by adding additional reagents that the enzyme uses to produce a reaction product that can be discerned. For example, when the agent horseradish peroxidase is present, the addition of hydrogen peroxide and diaminobenzidine leads to a colored reaction product, which is detectable. An antibody can also be labeled with biotin, and detected through indirect measurement of avidin or streptavidin binding. An antibody can also be labeled with a predetermined polypeptide epitope recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags). In some embodiments, labels are attached by spacer arms of various lengths to reduce potential steric hindrance.

- [0223] An anti-M-CSF antibody can also be labeled with a radiolabeled amino acid. The radiolabeled anti-M-CSF antibody can be used for both diagnostic and therapeutic purposes. For instance, the radiolabeled anti-M-CSF antibody can be used to detect M-CSF-expressing tumors by x-ray or other diagnostic techniques. Further, the radiolabeled anti-M-CSF antibody can be used therapeutically as a toxin for cancerous cells or tumors. Examples of labels for polypeptides include, but are not limited to, the following radioisotopes or radionuclides – ^3H , ^{14}C , ^{15}N , ^{35}S , ^{90}Y , ^{99}Tc , ^{111}In , ^{125}I , and ^{131}I .

- [0224] An anti-M-CSF antibody can also be derivatized with a chemical group such as polyethylene glycol (PEG), a methyl or ethyl group, or a carbohydrate group. These groups are useful to improve the biological characteristics of the antibody, e.g., to increase serum half-life or to increase tissue binding.

Pharmaceutical Compositions and Kits

- [0225] The invention also relates to compositions comprising a human anti-M-CSF antagonist antibody for the treatment of subjects in need of treatment for rheumatoid arthritis, osteoporosis, or atherosclerosis. In some embodiments, the subject of treatment is a human. In other embodiments, the subject is a veterinary

subject. Hyperproliferative disorders where monocytes play a role that may be treated by an antagonist anti-M-CSF antibody of the invention can involve any tissue or organ and include but are not limited to brain, lung, squamous cell, bladder, gastric, pancreatic, breast, head, neck, liver, renal, ovarian, prostate, colorectal, esophageal, gynecological, nasopharynx, or thyroid cancers, melanomas, lymphomas, leukemias or multiple myelomas. In particular, human antagonist anti-M-CSF antibodies of the invention are useful to treat or prevent carcinomas of the breast, prostate, colon and lung.

- [0226] This invention also encompasses compositions for the treatment of a condition selected from the group consisting of arthritis, psoriatic arthritis, Reiter's syndrome, gout, traumatic arthritis, rubella arthritis and acute synovitis, rheumatoid arthritis, rheumatoid spondylitis, ankylosing spondylitis, osteoarthritis, gouty arthritis and other arthritic conditions, sepsis, septic shock, endotoxic shock, gram negative sepsis, toxic shock syndrome, Alzheimer's disease, stroke, neurotrauma, asthma, adult respiratory distress syndrome, cerebral malaria, chronic pulmonary inflammatory disease, silicosis, pulmonary sarcoidosis, bone resorption disease, osteoporosis, restenosis, cardiac and renal reperfusion injury, thrombosis, glomerulonephritis, diabetes, graft vs. host reaction, allograft rejection, inflammatory bowel disease, Crohn's disease, ulcerative colitis, multiple sclerosis, muscle degeneration, eczema, contact dermatitis, psoriasis, sunburn, or conjunctivitis shock in a mammal, including a human, comprising an amount of a human anti-M-CSF monoclonal antibody of the invention effective in such treatment and a pharmaceutically acceptable carrier.

- [0227] Treatment may involve administration of one or more antagonist anti-M-CSF monoclonal antibodies of the invention, or antigen-binding fragments thereof, alone or with a pharmaceutically acceptable carrier. As used herein, "pharmaceutically acceptable carrier" means any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like that are physiologically compatible. Some examples of pharmaceutically acceptable carriers are water, saline, phosphate buffered saline, dextrose, glycerol, ethanol and the like, as well as combinations thereof. In many cases, it will be preferable to include isotonic agents, for example, sugars,

polyalcohols such as mannitol, sorbitol, or sodium chloride in the composition. Additional examples of pharmaceutically acceptable substances are wetting agents or minor amounts of auxiliary substances such as wetting or emulsifying agents, preservatives or buffers, which enhance the shelf life or effectiveness of the antibody.

[0228] Anti-M-CSF antibodies of the invention and compositions comprising them, can be administered in combination with one or more other therapeutic, diagnostic or prophylactic agents. Additional therapeutic agents include other anti-neoplastic, anti-tumor, anti-angiogenic or chemotherapeutic agents. Such additional agents may be included in the same composition or administered separately. In some embodiments, one or more inhibitory anti-M-CSF antibodies of the invention can be used as a vaccine or as adjuvants to a vaccine.

[0229] The compositions of this invention may be in a variety of forms, for example, liquid, semi-solid and solid dosage forms, such as liquid solutions (e.g., injectable and infusible solutions), dispersions or suspensions, tablets, pills, powders, liposomes and suppositories. The preferred form depends on the intended mode of administration and therapeutic application. Typical preferred compositions are in the form of injectable or infusible solutions, such as compositions similar to those used for passive immunization of humans. The preferred mode of administration is parenteral (e.g., intravenous, subcutaneous, intraperitoneal, intramuscular). In a preferred embodiment, the antibody is administered by intravenous infusion or injection. In another preferred embodiment, the antibody is administered by intramuscular or subcutaneous injection. In another embodiment, the invention includes a method of treating a subject in need thereof with an antibody or an antigen-binding portion thereof that specifically binds to M-CSF comprising the steps of: (a) administering an effective amount of an isolated nucleic acid molecule encoding the heavy chain or the antigen-binding portion thereof, an isolated nucleic acid molecule encoding the light chain or the antigen-binding portion thereof, or both the nucleic acid molecules encoding the light chain and the heavy chain or antigen-binding portions thereof; and (b) expressing the nucleic acid molecule.

- [0230] Therapeutic compositions typically must be sterile and stable under the conditions of manufacture and storage. The composition can be formulated as a solution, microemulsion, dispersion, liposome, or other ordered structure suitable to high drug concentration. Sterile injectable solutions can be prepared by
- 5 incorporating the anti-M-CSF antibody in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated
- 10 above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying that yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof. The proper fluidity of a solution can be maintained, for example, by the use of a coating such
- 15 as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prolonged absorption of injectable compositions can be brought about by including in the composition an agent that delays absorption, for example, monostearate salts and gelatin.
- [0231] The antibodies of the present invention can be administered by a variety
- 20 of methods known in the art, although for many therapeutic applications, the preferred route/mode of administration is subcutaneous, intramuscular, or intravenous infusion. As will be appreciated by the skilled artisan, the route and/or mode of administration will vary depending upon the desired results.
- [0232] In certain embodiments, the antibody compositions active compound may
- 25 be prepared with a carrier that will protect the antibody against rapid release, such as a controlled release formulation, including implants, transdermal patches, and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Many methods for the preparation
- 30 of such formulations are patented or generally known to those skilled in the art. See, e.g., *Sustained and Controlled Release Drug Delivery Systems* (J. R. Robinson, ed., Marcel Dekker, Inc., New York, 1978).

[0233] In certain embodiments, an anti-M-CSF antibody of the invention can be orally administered, for example, with an inert diluent or an assimilable edible carrier. The compound (and other ingredients, if desired) can also be enclosed in a hard or soft shell gelatin capsule, compressed into tablets, or incorporated directly into the subject's diet. For oral therapeutic administration, the anti-M-CSF antibodies can be incorporated with excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. To administer a compound of the invention by other than parenteral administration, it may be necessary to coat the compound with, or co-administer the compound with, a material to prevent its inactivation.

[0234] Additional active compounds also can be incorporated into the compositions. In certain embodiments, an anti-M-CSF antibody of the invention is co-formulated with and/or co-administered with one or more additional therapeutic agents. These agents include antibodies that bind other targets, antineoplastic agents, antitumor agents, chemotherapeutic agents, peptide analogues that inhibit M-CSF, soluble *c-fms* that can bind M-CSF, one or more chemical agents that inhibit M-CSF, anti-inflammatory agents, anti-coagulants, agents that lower blood pressure (i.e., angiotensin-converting enzyme (ACE) inhibitors). Such combination therapies may require lower dosages of the anti-M-CSF antibody as well as the co-administered agents, thus avoiding possible toxicities or complications associated with the various monotherapies.

[0235] Inhibitory anti-M-CSF antibodies of the invention and compositions comprising them also may be administered in combination with other therapeutic regimens, in particular in combination with radiation treatment for cancer. The compounds of the present invention may also be used in combination with anticancer agents such as endostatin and angiostatin or cytotoxic drugs such as adriamycin, daunomycin, cis-platinum, etoposide, taxol, taxotere and alkaloids, such as vincristine, farnesyl transferase inhibitors, VEGF inhibitors, and antimetabolites such as methotrexate.

[0236] The compounds of the invention may also be used in combination with antiviral agents such as Viracept, AZT, aciclovir and famciclovir, and antiseptic compounds such as Valant.

[0237] The compositions of the invention may include a “therapeutically effective amount” or a “prophylactically effective amount” of an antibody or antigen-binding portion of the invention. A “therapeutically effective amount” refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired therapeutic result. A therapeutically effective amount of the antibody or antibody portion may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the antibody or antibody portion to elicit a desired response in the individual. A therapeutically effective amount is also one in which any toxic or detrimental effects of the antibody or antibody portion are outweighed by the therapeutically beneficial effects. A “prophylactically effective amount” refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired prophylactic result. Typically, since a prophylactic dose is used in subjects prior to or at an earlier stage of disease, the prophylactically effective amount will be less than the therapeutically effective amount.

[0238] Dosage regimens can be adjusted to provide the optimum desired response (e.g., a therapeutic or prophylactic response). For example, a single bolus can be administered, several divided doses can be administered over time or the dose can be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the mammalian subjects to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on (a) the unique characteristics of the anti-M-CSF antibody or portion and the particular therapeutic or prophylactic effect to be achieved, and (b) the limitations inherent in the art of compounding such an antibody for the treatment of sensitivity in individuals.

[0239] An exemplary, non-limiting range for a therapeutically or prophylactically effective amount of an antibody or antibody portion of the

invention is 0.025 to 50 mg/kg, more preferably 0.1 to 50 mg/kg, more preferably 0.1-25, 0.1 to 10 or 0.1 to 3 mg/kg. It is to be noted that dosage values may vary with the type and severity of the condition to be alleviated. It is to be further understood that for any particular subject, specific dosage regimens should be
5 adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions, and that dosage ranges set forth herein are exemplary only and are not intended to limit the scope or practice of the claimed composition.

[0240] Another aspect of the present invention provides kits comprising an anti-
10 M-CSF antibody or antigen-binding portion of the invention or a composition comprising such an antibody or portion. A kit may include, in addition to the antibody or composition, diagnostic or therapeutic agents. A kit also can include instructions for use in a diagnostic or therapeutic method. In a preferred embodiment, the kit includes the antibody or a composition comprising it and a
15 diagnostic agent that can be used in a method described below. In another preferred embodiment, the kit includes the antibody or a composition comprising it and one or more therapeutic agents that can be used in a method described below. One embodiment of the invention is a kit comprising a container, instructions on the administration of an anti-M-CSF antibody to a human suffering from an
20 inflammatory disease, or instructions for measuring the number of CD14⁺CD16⁺ monocytes in a biological sample and an anti-M-CSF antibody.

[0241] This invention also relates to compositions for inhibiting abnormal cell growth in a mammal comprising an amount of an antibody of the invention in combination with an amount of a chemotherapeutic agent, wherein the amounts of
25 the compound, salt, solvate, or prodrug, and of the chemotherapeutic agent are together effective in inhibiting abnormal cell growth. Many chemotherapeutic agents are known in the art. In some embodiments, the chemotherapeutic agent is selected from the group consisting of mitotic inhibitors, alkylating agents, anti-metabolites, intercalating antibiotics, growth factor inhibitors, cell cycle inhibitors,
30 enzymes, topoisomerase inhibitors, biological response modifiers, anti-hormones, e.g. anti-androgens, and anti-angiogenesis agents.

- [0242] Anti-angiogenic agents, such as MMP-2 (matrix-metalloproteinase 2) inhibitors, MMP-9 (matrix-metalloproteinase 9) inhibitors, and COX-II (cyclooxygenase II) inhibitors, can be used in conjunction with an anti-M-CSF antibody of the invention. Examples of useful COX-II inhibitors include
- 5 CELEBREXTM (celecoxib), valdecoxib, and rofecoxib. Examples of useful matrix metalloproteinase inhibitors are described in WO 96/33172 (published October 24, 1996), WO 96/27583 (published March 7, 1996), European Patent Application No. 97304971.1 (filed July 8, 1997), European Patent Application No. 99308617.2 (filed October 29, 1999), WO 98/07697 (published February 26, 1998), WO
- 10 98/03516 (published January 29, 1998), WO 98/34918 (published August 13, 1998), WO 98/34915 (published August 13, 1998), WO 98/33768 (published August 6, 1998), WO 98/30566 (published July 16, 1998), European Patent Publication 606,046 (published July 13, 1994), European Patent Publication 931,788 (published July 28, 1999), WO 90/05719 (published May 31, 1990), WO
- 15 99/52910 (published October 21, 1999), WO 99/52889 (published October 21, 1999), WO 99/29667 (published June 17, 1999), PCT International Application No. PCT/IB98/01113 (filed July 21, 1998), European Patent Application No. 99302232.1 (filed March 25, 1999), Great Britain patent application number 9912961.1 (filed June 3, 1999), U.S. Provisional Application No. 60/148,464 (filed
- 20 August 12, 1999), U.S. Patent 5,863,949 (issued January 26, 1999), U.S. Patent 5,861,510 (issued January 19, 1999), and European Patent Publication 780,386 (published June 25, 1997), all of which are incorporated herein in their entireties by reference. Preferred MMP inhibitors are those that do not demonstrate arthralgia. More preferred, are those that selectively inhibit MMP-2 and/or MMP-
- 25 9 relative to the other matrix-metalloproteinases (i.e. MMP-1, MMP-3, MMP-4, MMP-5, MMP-6, MMP-7, MMP-8, MMP-10, MMP-11, MMP-12, and MMP-13). Some specific examples of MMP inhibitors useful in the present invention are AG-3340, RO 32-3555, RS 13-0830, and the compounds recited in the following list:
- 30 3-[[4-(4-fluoro-phenoxy)-benzenesulfonyl]-(1-hydroxycarbonyl-cyclopentyl)-amino]-propionic acid; 3-exo-3-[4-(4-fluoro-phenoxy)-benzenesulfonylamino]-8-oxa-bicyclo[3.2.1]octane-3-carboxylic acid hydroxyamide; (2R, 3R) 1-[4-(2-chloro-4-fluoro-benzoyloxy)-benzenesulfonyl]-3-hydroxy-3-methyl-piperidine-2-

- carboxylic acid hydroxyamide; 4-[4-(4-fluoro-phenoxy)-benzenesulfonylamino]-tetrahydro-pyran-4-carboxylic acid hydroxyamide; 3-[[4-(4-fluoro-phenoxy)-benzenesulfonyl]-(1-hydroxycarbamoyl-cyclobutyl)-amino]-propionic acid; 4-[4-(4-chloro-phenoxy)-benzenesulfonylamino]-tetrahydro-pyran-4-carboxylic acid hydroxyamide; (R) 3-[4-(4-chloro-phenoxy)-benzenesulfonylamino]-tetrahydro-pyran-3-carboxylic acid hydroxyamide; (2R, 3R) 1-[4-(4-fluoro-2-methyl-benzyloxy)-benzenesulfonyl]-3-hydroxy-3-methyl-piperidine-2-carboxylic acid hydroxyamide; 3-[[4-(4-fluoro-phenoxy)-benzenesulfonyl]-(1-hydroxycarbamoyl-1-methyl-ethyl)-amino]-propionic acid; 3-[[4-(4-fluoro-phenoxy)-benzenesulfonyl]-(4-hydroxycarbamoyl-tetrahydro-pyran-4-yl)-amino]-propionic acid; 3-exo-3-[4-(4-chloro-phenoxy)-benzenesulfonylamino]-8-oxa-bicyclo[3.2.1]octane-3-carboxylic acid hydroxyamide; 3-endo-3-[4-(4-fluoro-phenoxy)-benzenesulfonylamino]-8-oxa-bicyclo[3.2.1]octane-3-carboxylic acid hydroxyamide; and (R) 3-[4-(4-fluoro-phenoxy)-benzenesulfonylamino]-tetrahydro-furan-3-carboxylic acid hydroxyamide; and pharmaceutically acceptable salts and solvates of said compounds.

- [0243] A compound comprising a human anti-M-CSF monoclonal antibody of the invention can also be used with signal transduction inhibitors, such as agents that can inhibit EGF-R (epidermal growth factor receptor) responses, such as EGF-R antibodies, EGF antibodies, and molecules that are EGF-R inhibitors; VEGF (vascular endothelial growth factor) inhibitors, such as VEGF receptors and molecules that can inhibit VEGF; and erbB2 receptor inhibitors, such as organic molecules or antibodies that bind to the erbB2 receptor, for example, HERCEPTINTM (Genentech, Inc.). EGF-R inhibitors are described in, for example
- in WO 95/19970 (published July 27, 1995), WO 98/14451 (published April 9, 1998), WO 98/02434 (published January 22, 1998), and United States Patent 5,747,498 (issued May 5, 1998), and such substances can be used in the present invention as described herein. EGFR-inhibiting agents include, but are not limited to, the monoclonal antibodies C225 and anti-EGFR 22Mab (ImClone Systems Incorporated), ABX-EGF (Abgenix/Cell Genesys), EMD-7200 (Merck KgaA), EMD-5590 (Merck KgaA), MDX-447/H-477 (Medarex Inc. and Merck KgaA), and the compounds ZD-1834, ZD-1838 and ZD-1839 (AstraZeneca), PKI-166

- (Novartis), PKI-166/CGP-75166 (Novartis), PTK 787 (Novartis), CP 701 (Cephalon), leflunomide (Pharmacia/Sugen), CI-1033 (Warner Lambert Parke Davis), CI-1033/PD 183,805 (Warner Lambert Parke Davis), CL-387,785 (Wyeth-Ayerst), BBR-1611 (Boehringer Mannheim GmbH/Roche), Naamidine A (Bristol Myers Squibb), RC-3940-II (Pharmacia), BIBX-1382 (Boehringer Ingelheim), OLX-103 (Merck & Co.), VRCTC-310 (Ventech Research), EGF fusion toxin (Seragen Inc.), DAB-389 (Seragen/Lilgand), ZM-252808 (Imperial Cancer Research Fund), RG-50864 (INSERM), LFM-A12 (Parker Hughes Cancer Center), WHI-P97 (Parker Hughes Cancer Center), GW-282974 (Glaxo), KT-8391 (Kyowa Hakko) and EGF-R Vaccine (York Medical/Centro de Immunologia Molecular (CIM)). These and other EGF-R-inhibiting agents can be used in the present invention.
- [0244] VEGF inhibitors, for example SU-5416 and SU-6668 (Sugen Inc.), AVASTIN™ (Genentech), SH-268 (Schering), and NX-1838 (NeXstar) can also be combined with the compound of the present invention. VEGF inhibitors are described in, for example in WO 99/24440 (published May 20, 1999), PCT International Application PCT/IB99/00797 (filed May 3, 1999), in WO 95/21613 (published August 17, 1995), WO 99/61422 (published December 2, 1999), United States Patent 5,834,504 (issued November 10, 1998), WO 98/50356 (published November 12, 1998), United States Patent 5,883,113 (issued March 16, 1999), United States Patent 5,886,020 (issued March 23, 1999), United States Patent 5,792,783 (issued August 11, 1998), WO 99/10349 (published March 4, 1999), WO 97/32856 (published September 12, 1997), WO 97/22596 (published June 26, 1997), WO 98/54093 (published December 3, 1998), WO 98/02438 (published January 22, 1998), WO 99/16755 (published April 8, 1999), and WO 98/02437 (published January 22, 1998), all of which are incorporated herein in their entireties by reference. Other examples of some specific VEGF inhibitors useful in the present invention are IM862 (Cytran Inc.); anti-VEGF monoclonal antibody of Genentech, Inc.; and angiozyme, a synthetic ribozyme from Ribozyme and Chiron.
- These and other VEGF inhibitors can be used in the present invention as described herein. ErbB2 receptor inhibitors, such as GW-282974 (Glaxo Wellcome plc), and the monoclonal antibodies AR-209 (Aronex Pharmaceuticals Inc.) and 2B-1

(Chiron), can furthermore be combined with the compound of the invention, for example those indicated in WO 98/02434 (published January 22, 1998), WO 99/35146 (published July 15, 1999), WO 99/35132 (published July 15, 1999), WO 98/02437 (published January 22, 1998), WO 97/13760 (published April 17, 1997),
5 WO 95/19970 (published July 27, 1995), United States Patent 5,587,458 (issued December 24, 1996), and United States Patent 5,877,305 (issued March 2, 1999), which are all hereby incorporated herein in their entireties by reference. ErbB2 receptor inhibitors useful in the present invention are also described in United States Patent 6,465,449 (issued October 15, 2002), and in United States Patent
10 6,284,764 (issued September 4, 2001), both of which are incorporated in their entireties herein by reference. The erbB2 receptor inhibitor compounds and substance described in the aforementioned PCT applications, U.S. patents, and U.S. provisional applications, as well as other compounds and substances that inhibit the erbB2 receptor, can be used with the compound of the present invention
15 in accordance with the present invention.

[0245] Anti-survival agents include anti-IGF-IR antibodies and anti-integrin agents, such as anti-integrin antibodies.

[0246] Anti-inflammatory agents can be used in conjunction with an anti-M-CSF antibody of the invention. For the treatment of rheumatoid arthritis, the human
20 anti-M-CSF antibodies of the invention may be combined with agents such as TNF- α inhibitors such as TNF drugs (such as REMICADE™, CDP-870 and HUMIRA™) and TNF receptor immunoglobulin molecules (such as ENBRELE™), IL-1 inhibitors, receptor antagonists or soluble IL-1ra (e.g. Kineret or ICE inhibitors), COX-2 inhibitors (such as celecoxib, rofecoxib, valdecoxib and etoricoxib), metalloprotease inhibitors (preferably MMP-13 selective inhibitors),
25 p2X7 inhibitors, $\alpha 2\delta$ ligands (such as NEUROTIN™ AND PREGABALIN™), low dose methotrexate, leflunomide, hydroxychloroquine, d-penicillamine, auranofin or parenteral or oral gold. The compounds of the invention can also be used in combination with existing therapeutic agents for the treatment of
30 osteoarthritis. Suitable agents to be used in combination include standard non-steroidal anti-inflammatory agents (hereinafter NSAID's) such as piroxicam, diclofenac, propionic acids such as naproxen, flurbiprofen, fenoprofen, ketoprofen

and ibuprofen, fenamates such as mefenamic acid, indomethacin, sulindac, apazone, pyrazolones such as phenylbutazone, salicylates such as aspirin, COX-2 inhibitors such as celecoxib, valdecoxib, rofecoxib and etoricoxib, analgesics and intraarticular therapies such as corticosteroids and hyaluronic acids such as

5 hyalgan and synvisc.

[0247] Anti-coagulant agents can be used in conjunction with an anti-M-CSF antibody of the invention. Examples of anti-coagulant agents include, but are not limited to, warfarin (COUMADIN™), heparin, and enoxaparin (LOVENOX™).

[0248] The human anti-M-CSF antibodies of the present invention may also be
10 used in combination with cardiovascular agents such as calcium channel blockers, lipid lowering agents such as statins, fibrates, beta-blockers, Ace inhibitors, Angiotensin-2 receptor antagonists and platelet aggregation inhibitors. The compounds of the present invention may also be used in combination with CNS agents such as antidepressants (such as sertraline), anti-Parkinsonian drugs (such as
15 deprenyl, L-dopa, REQUIP™, MIRAPEX™, MAOB inhibitors such as selegine and rasagiline, comP inhibitors such as Tasmar, A-2 inhibitors, dopamine reuptake inhibitors, NMDA antagonists, Nicotine agonists, Dopamine agonists and inhibitors of neuronal nitric oxide synthase), and anti-Alzheimer's drugs such as donepezil, tacrine, $\alpha 2\delta$ LIGANDS (such as NEUROTIN™ and PREGABALIN™)
20 inhibitors, COX-2 inhibitors, propentofylline or metryfonate.

[0249] The human anti-M-CSF antibodies of the present invention may also be used in combination with osteoporosis agents such as roloxifene, droloxifene, lasofoxifene or fosomax and immunosuppressant agents such as FK-506 and rapamycin.

25 Diagnostic Methods of Use

[0250] In another aspect, the invention provides diagnostic methods. The anti-M-CSF antibodies can be used to detect M-CSF in a biological sample *in vitro* or *in vivo*. In one embodiment, the invention provides a method for diagnosing the presence or location of a M-CSF-expressing tumor in a subject in need thereof,
30 comprising the steps of injecting the antibody into the subject, determining the expression of M-CSF in the subject by localizing where the antibody has bound,

comparing the expression in the subject with that of a normal reference subject or standard, and diagnosing the presence or location of the tumor.

[0251] The anti-M-CSF antibodies can be used in a conventional immunoassay, including, without limitation, an ELISA, an RIA, FACS, tissue

- 5 immunohistochemistry, Western blot or immunoprecipitation. The anti-M-CSF antibodies of the invention can be used to detect M-CSF from humans. In another embodiment, the anti-M-CSF antibodies can be used to detect M-CSF from primates such as cynomolgus monkey, rhesus monkeys, chimpanzees or apes. The invention provides a method for detecting M-CSF in a biological sample
- 10 comprising contacting a biological sample with an anti-M-CSF antibody of the invention and detecting the bound antibody. In one embodiment, the anti-M-CSF antibody is directly labeled with a detectable label. In another embodiment, the anti-M-CSF antibody (the first antibody) is unlabeled and a second antibody or other molecule that can bind the anti-M-CSF antibody is labeled. As is well
- 15 known to one of skill in the art, a second antibody is chosen that is able to specifically bind the particular species and class of the first antibody. For example, if the anti-M-CSF antibody is a human IgG, then the secondary antibody could be an anti-human-IgG. Other molecules that can bind to antibodies include, without limitation, Protein A and Protein G, both of which are available commercially, e.g.,
- 20 from Pierce Chemical Co.

- [0252] Suitable labels for the antibody or secondary antibody have been disclosed *supra*, and include various enzymes, prosthetic groups, fluorescent materials, luminescent materials and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or
- 25 acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S or ^3H .
- 30

[0253] In other embodiments, M-CSF can be assayed in a biological sample by a competition immunoassay utilizing M-CSF standards labeled with a detectable

substance and an unlabeled anti-M-CSF antibody. In this assay, the biological sample, the labeled M-CSF standards and the anti-M-CSF antibody are combined and the amount of labeled M-CSF standard bound to the unlabeled antibody is determined. The amount of M-CSF in the biological sample is inversely

- 5 proportional to the amount of labeled M-CSF standard bound to the anti-M-CSF antibody.

[0254] One can use the immunoassays disclosed above for a number of purposes. For example, the anti-M-CSF antibodies can be used to detect M-CSF in cells or on the surface of cells in cell culture, or secreted into the tissue culture medium.

- 10 The anti-M-CSF antibodies can be used to determine the amount of M-CSF on the surface of cells or secreted into the tissue culture medium that have been treated with various compounds. This method can be used to identify compounds that are useful to inhibit or activate M-CSF expression or secretion. According to this method, one sample of cells is treated with a test compound for a period of time
- 15 while another sample is left untreated. If the total level of M-CSF is to be measured, the cells are lysed and the total M-CSF level is measured using one of the immunoassays described above. The total level of M-CSF in the treated versus the untreated cells is compared to determine the effect of the test compound.

- [0255] An immunoassay for measuring total M-CSF levels is an ELISA or
- 20 Western blot. If the cell surface level of M-CSF is to be measured, the cells are not lysed, and the M-CSF cell surface levels can be measured using one of the immunoassays described above. An immunoassay for determining cell surface levels of M-CSF can include the steps of labeling the cell surface proteins with a detectable label, such as biotin or ^{125}I , immunoprecipitating the M-CSF with an
- 25 anti-M-CSF antibody and then detecting the labeled M-CSF. Another immunoassay for determining the localization of M-CSF, e.g., cell surface levels, can be immunohistochemistry. Methods such as ELISA, RIA, Western blot, immunohistochemistry, cell surface labeling of integral membrane proteins and immunoprecipitation are well known in the art. See, e.g., Harlow and Lane, *supra*.
- 30 In addition, the immunoassays can be scaled up for high throughput screening in order to test a large number of compounds for inhibition or activation of M-CSF.

- [0256] Another example of an immunoassay for measuring secreted M-CSF levels can be an antigen capture assay, ELISA, immunohistochemistry assay, Western blot and the like using antibodies of the invention. If secreted M-CSF is to be measured, cell culture media or body fluid, such as blood serum, urine, or synovial fluid, can be assayed for secreted M-CSF and/or cells can be lysed to release produced, but not yet secreted M-CSF. An immunoassay for determining secreted levels of M-CSF includes the steps of labeling the secreted proteins with a detectable label, such as biotin or ^{125}I , immunoprecipitating the M-CSF with an anti-M-CSF antibody and then detecting the labeled M-CSF. Another immunoassay for determining secreted levels of M-CSF can include the steps of (a) pre-binding anti-M-CSF antibodies to the surface of a microtiter plate; (b) adding tissue culture cell media or body fluid containing the secreted M-CSF to the wells of the microtiter plate to bind to the anti-M-CSF antibodies; (c) adding an antibody that will detect the anti-M-CSF antibody, e.g., anti-M-CSF labeled with digoxigenin that binds to an epitope of M-CSF different from the anti-M-CSF antibody of step (a); (d) adding an antibody to digoxigenin conjugated to peroxidase; and (e) adding a peroxidase substrate that will yield a colored reaction product that can be quantitated to determine the level of secreted M-CSF in tissue culture cell media or a body fluid sample. Methods such as ELISA, RIA, Western blot, immunohistochemistry, and antigen capture assay are well known in the art. See, e.g., Harlow and Lane, *supra*. In addition, the immunoassays can be scaled up for high throughput screening in order to test a large number of compounds for inhibition or activation of M-CSF.
- [0257] The anti-M-CSF antibodies of the invention can also be used to determine the levels of cell surface M-CSF in a tissue or in cells derived from the tissue. In some embodiments, the tissue is from a diseased tissue. In some embodiments, the tissue can be a tumor or a biopsy thereof. In some embodiments of the method, a tissue or a biopsy thereof can be excised from a patient. The tissue or biopsy can then be used in an immunoassay to determine, e.g., total M-CSF levels, cell surface levels of M-CSF, or localization of M-CSF by the methods discussed above.
- [0258] The method can comprise the steps of administering a detectably labeled anti-M-CSF antibody or a composition comprising them to a patient in need of

- such a diagnostic test and subjecting the patient to imaging analysis to determine the location of the M-CSF-expressing tissues. Imaging analysis is well known in the medical art, and includes, without limitation, x-ray analysis, magnetic resonance imaging (MRI) or computed tomography (CE). The antibody can be
- 5 labeled with any agent suitable for *in vivo* imaging, for example a contrast agent, such as barium, which can be used for x-ray analysis, or a magnetic contrast agent, such as a gadolinium chelate, which can be used for MRI or CE. Other labeling agents include, without limitation, radioisotopes, such as ⁹⁹Tc. In another embodiment, the anti-M-CSF antibody will be unlabeled and will be imaged by
- 10 administering a second antibody or other molecule that is detectable and that can bind the anti-M-CSF antibody. In an embodiment, a biopsy is obtained from the patient to determine whether the tissue of interest expresses M-CSF.
- [0259] The anti-M-CSF antibodies of the invention can also be used to determine the secreted levels of M-CSF in a body fluid such as blood serum, urine, or
- 15 synovial fluid derived from a tissue. In some embodiments, the body fluid is from a diseased tissue. In some embodiments, the body fluid is from a tumor or a biopsy thereof. In some embodiments of the method, body fluid is removed from a patient. The body fluid is then used in an immunoassay to determine secreted M-CSF levels by the methods discussed above. One embodiment of the invention is a
- 20 method of assaying for the activity of a M-CSF antagonist comprising: administering a M-CSF antagonist to a primate or human subject and measuring the number of CD14+CD16+ monocytes in a biological sample.

Therapeutic Methods of Use

- 25 [0260] In another embodiment, the invention provides a method for inhibiting M-CSF activity by administering an anti-M-CSF antibody to a patient in need thereof. Any of the types of antibodies described herein may be used therapeutically. In a preferred embodiment, the anti-M-CSF antibody is a human, chimeric or humanized antibody. In another preferred embodiment, the M-CSF is
- 30 human and the patient is a human patient. Alternatively, the patient may be a mammal that expresses a M-CSF that the anti-M-CSF antibody cross-reacts with. The antibody may be administered to a non-human mammal expressing a M-CSF

with which the antibody cross-reacts (i.e. a primate) for veterinary purposes or as an animal model of human disease. Such animal models may be useful for evaluating the therapeutic efficacy of antibodies of this invention.

[0261] As used herein, the term “a disorder in which M-CSF activity is detrimental” is intended to include diseases and other disorders in which the presence of high levels of M-CSF in a subject suffering from the disorder has been shown to be or is suspected of being either responsible for the pathophysiology of the disorder or a factor that contributes to a worsening of the disorder. Such disorders may be evidenced, for example, by an increase in the levels of M-CSF secreted and/or on the cell surface or increased tyrosine autophosphorylation of *c-fms* in the affected cells or tissues of a subject suffering from the disorder. The increase in M-CSF levels may be detected, for example, using an anti-M-CSF antibody as described above.

[0262] In one embodiment, an anti-M-CSF antibody may be administered to a patient who has a *c-fms*-expressing tumor or a tumor that secretes M-CSF and/or that expresses M-CSF on its cell surface. Preferably, the tumor expresses a level of *c-fms* or M-CSF that is higher than a normal tissue. The tumor may be a solid tumor or may be a non-solid tumor, such as a lymphoma. In a more preferred embodiment, an anti-M-CSF antibody may be administered to a patient who has a *c-fms*-expressing tumor, a M-CSF-expressing tumor, or a tumor that secretes M-CSF that is cancerous. Further, the tumor may be cancerous. In an even more preferred embodiment, the tumor is a cancer of lung, breast, prostate or colon. In another preferred embodiment, the anti-M-CSF antibody administered to a patient results in M-CSF no longer bound to the *c-fms* receptor. In a highly preferred embodiment, the method causes the tumor not to increase in weight or volume or to decrease in weight or volume. In another embodiment, the method causes *c-fms* on tumor cells to not be bound by M-CSF. In another embodiment, the method causes M-CSF on tumor cells to not be bound to *c-fms*. In another embodiment, the method causes secreted M-CSF of the tumor cells to not be bound to *c-fms*. In a preferred embodiment, the antibody is selected from 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-

Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 or 9.14.4G1, or comprises a heavy chain, light chain or antigen binding region thereof.

- [0263] In another preferred embodiment, an anti-M-CSF antibody may be administered to a patient who expresses inappropriately high levels of M-CSF. It
- 5 is known in the art that high-level expression of M-CSF can lead to a variety of common cancers. In one embodiment, said method relates to the treatment of cancer such as brain, squamous cell, bladder, gastric, pancreatic, breast, head, neck, esophageal, prostate, colorectal, lung, renal, kidney, ovarian, gynecological or thyroid cancer. Patients that can be treated with a compounds of the invention
- 10 according to the methods of this invention include, for example, patients that have been diagnosed as having lung cancer, bone cancer, pancreatic cancer, skin cancer, cancer of the head and neck, cutaneous or intraocular melanoma, uterine cancer, ovarian cancer, rectal cancer, cancer of the anal region, stomach cancer, colon cancer, breast cancer, gynecologic tumors (e.g., uterine sarcomas, carcinoma of the fallopian tubes, carcinoma of the endometrium, carcinoma of the cervix, carcinoma of the vagina or carcinoma of the vulva), Hodgkin's disease, cancer of the esophagus, cancer of the small intestine, cancer of the endocrine system (e.g., cancer of the thyroid, parathyroid or adrenal glands), sarcomas of soft tissues, cancer of the urethra, cancer of the penis, prostate cancer, chronic or acute
- 20 leukemia, solid tumors (e.g., sarcomas, carcinomas or lymphomas that are cancers of body tissues other than blood, bone marrow or the lymphatic system), solid tumors of childhood, lymphocytic lymphomas, cancer of the bladder, cancer of the kidney or ureter (e.g., renal cell carcinoma, carcinoma of the renal pelvis), or neoplasms of the central nervous system (e.g., primary CNS lymphoma, spinal axis tumors, brain stem gliomas or pituitary adenomas). In a more preferred embodiment, the anti-M-CSF antibody is administered to a patient with breast cancer, prostate cancer, lung cancer or colon cancer. In an even more preferred embodiment, the method causes the cancer to stop proliferating abnormally, or not to increase in weight or volume or to decrease in weight or volume.
- 30 [0264] The antibody may be administered once, but more preferably is administered multiple times. For example, the antibody may be administered from three times daily to once every six months or longer. The administering may be on

- a schedule such as three times daily, twice daily, once daily, once every two days, once every three days, once weekly, once every two weeks, once every month, once every two months, once every three months and once every six months. The antibody may also be administered continuously via a minipump. The antibody
- 5 may be administered via an oral, mucosal, buccal, intranasal, inhalable, intravenous, subcutaneous, intramuscular, parenteral, intratumor or topical route. The antibody may be administered at the site of the tumor or inflamed body part, into the tumor or inflamed body part, or at a site distant from the site of the tumor or inflamed body part. The antibody may be administered once, at least twice or
- 10 for at least the period of time until the condition is treated, palliated or cured. The antibody generally will be administered for as long as the tumor is present provided that the antibody causes the tumor or cancer to stop growing or to decrease in weight or volume or until the inflamed body part is healed. The antibody will generally be administered as part of a pharmaceutical composition as
- 15 described *supra*. The dosage of antibody will generally be in the range of 0.1-100 mg/kg, more preferably 0.5-50 mg/kg, more preferably 1-20 mg/kg, and even more preferably 1-10 mg/kg. The serum concentration of the antibody may be measured by any method known in the art.
- [0265] In another aspect, the anti-M-CSF antibody may be co-administered with
- 20 other therapeutic agents, such as anti-inflammatory agents, anti-coagulant agents, agents that will lower or reduce blood pressure, anti-neoplastic drugs or molecules, to a patient who has a hyperproliferative disorder, such as cancer or a tumor. In one aspect, the invention relates to a method for the treatment of the hyperproliferative disorder in a mammal comprising administering to said mammal
- 25 a therapeutically effective amount of a compound of the invention in combination with an anti-tumor agent selected from the group consisting of, but not limited to, mitotic inhibitors, alkylating agents, anti-metabolites, intercalating agents, growth factor inhibitors, cell cycle inhibitors, enzymes, topoisomerase inhibitors, biological response modifiers, anti-hormones, kinase inhibitors, matrix
- 30 metalloprotease inhibitors, genetic therapeutics and anti-androgens. In a more preferred embodiment, the antibody may be administered with an antineoplastic agent, such as adriamycin or taxol. In another preferred embodiment, the antibody

or combination therapy is administered along with radiotherapy, chemotherapy, photodynamic therapy, surgery or other immunotherapy. In yet another preferred embodiment, the antibody will be administered with another antibody. For example, the anti-M-CSF antibody may be administered with an antibody or other agent that is known to inhibit tumor or cancer cell proliferation, e.g., an antibody or agent that inhibits erbB2 receptor, EGF-R, CD20 or VEGF.

[0266] Co-administration of the antibody with an additional therapeutic agent (combination therapy) encompasses administering a pharmaceutical composition comprising the anti-M-CSF antibody and the additional therapeutic agent and administering two or more separate pharmaceutical compositions, one comprising the anti-M-CSF antibody and the other(s) comprising the additional therapeutic agent(s). Further, although co-administration or combination therapy generally means that the antibody and additional therapeutic agents are administered at the same time as one another, it also encompasses instances in which the antibody and additional therapeutic agents are administered at different times. For instance, the antibody may be administered once every three days, while the additional therapeutic agent is administered once daily. Alternatively, the antibody may be administered prior to or subsequent to treatment of the disorder with the additional therapeutic agent. Similarly, administration of the anti-M-CSF antibody may be administered prior to or subsequent to other therapy, such as radiotherapy, chemotherapy, photodynamic therapy, surgery or other immunotherapy

[0267] The antibody and one or more additional therapeutic agents (the combination therapy) may be administered once, twice or at least the period of time until the condition is treated, palliated or cured. Preferably, the combination therapy is administered multiple times. The combination therapy may be administered from three times daily to once every six months. The administering may be on a schedule such as three times daily, twice daily, once daily, once every two days, once every three days, once weekly, once every two weeks, once every month, once every two months, once every three months and once every six months, or may be administered continuously via a minipump. The combination therapy may be administered via an oral, mucosal, buccal, intranasal, inhalable, intravenous, subcutaneous, intramuscular, parenteral, intratumor or topical route.

The combination therapy may be administered at a site distant from the site of the tumor. The combination therapy generally will be administered for as long as the tumor is present provided that the antibody causes the tumor or cancer to stop growing or to decrease in weight or volume.

- 5 [0268] In a still further embodiment, the anti-M-CSF antibody is labeled with a radiolabel, an immunotoxin or a toxin, or is a fusion protein comprising a toxic peptide. The anti-M-CSF antibody or anti-M-CSF antibody fusion protein directs the radiolabel, immunotoxin, toxin or toxic peptide to the M-CSF-expressing cell. In a preferred embodiment, the radiolabel, immunotoxin, toxin or toxic peptide is
- 10 internalized after the anti-M-CSF antibody binds to the M-CSF on the surface of the target cell.
- [0269] In another aspect, the anti-M-CSF antibody may be used to treat noncancerous states in which high levels of M-CSF and/or M-CSF have been associated with the noncancerous state or disease. In one embodiment, the method
- 15 comprises the step of administering an anti-M-CSF antibody to a patient who has a noncancerous pathological state caused or exacerbated by high levels of M-CSF and/or M-CSF levels or activity. In a more preferred embodiment, the anti-M-CSF antibody slows the progress of the noncancerous pathological state. In a more preferred embodiment, the anti-M-CSF antibody stops or reverses, at least in part,
- 20 the noncancerous pathological state.

Gene Therapy

- [0270] The nucleic acid molecules of the instant invention can be administered to a patient in need thereof via gene therapy. The therapy may be either *in vivo* or *ex vivo*. In a preferred embodiment, nucleic acid molecules encoding both a heavy
- 25 chain and a light chain are administered to a patient. In a more preferred embodiment, the nucleic acid molecules are administered such that they are stably integrated into chromosomes of B cells because these cells are specialized for producing antibodies. In a preferred embodiment, precursor B cells are transfected or infected *ex vivo* and re-transplanted into a patient in need thereof. In another
- 30 embodiment, precursor B cells or other cells are infected *in vivo* using a virus known to infect the cell type of interest. Typical vectors used for gene therapy include liposomes, plasmids and viral vectors. Exemplary viral vectors are

retroviruses, adenoviruses and adeno-associated viruses. After infection either *in vivo* or *ex vivo*, levels of antibody expression can be monitored by taking a sample from the treated patient and using any immunoassay known in the art or discussed herein.

- 5 [0271] In a preferred embodiment, the gene therapy method comprises the steps of administering an isolated nucleic acid molecule encoding the heavy chain or an antigen-binding portion thereof of an anti-M-CSF antibody and expressing the nucleic acid molecule. In another embodiment, the gene therapy method comprises the steps of administering an isolated nucleic acid molecule encoding the light chain or an antigen-binding portion thereof of an anti-M-CSF antibody and expressing the nucleic acid molecule. In a more preferred method, the gene therapy method comprises the steps of administering of an isolated nucleic acid molecule encoding the heavy chain or an antigen-binding portion thereof and an isolated nucleic acid molecule encoding the light chain or the antigen-binding portion thereof of an anti-M-CSF antibody of the invention and expressing the nucleic acid molecules. The gene therapy method may also comprise the step of administering another anti-cancer agent, such as taxol or adriamycin.
- 10 [0272] In order that this invention may be better understood, the following examples are set forth. These examples are for purposes of illustration only and are not to be construed as limiting the scope of the invention in any manner.
- 15
- 20

EXAMPLE I

Generation of Cell Lines Producing Anti-M-CSF Antibody

- [0273] Antibodies of the invention were prepared, selected, and assayed as follows:
- 25 *Immunization and hybridoma generation*
- Eight to ten week old XENOMOUSE™ mice were immunized intraperitoneally or in their hind footpads with human M-CSF (10 µg/dose/mouse). This dose was repeated five to seven times over a three to eight week period. Four days before fusion, the mice were given a final injection of human M-CSF in PBS. The spleen and lymph node lymphocytes from immunized mice were fused with the non-secretory myeloma P3-X63-Ag8.653 cell line, and the fused cells were subjected to HAT selection as previously described (Galfre and Milstein, *Methods Enzymol.*
- 30

- 73:3-46, 1981). A panel of hybridomas all secreting M-CSF specific human IgG2 and IgG4 antibodies was recovered. Antibodies also were generated using XENOMAXTM technology as described in Babcook, J.S. *et al.*, *Proc. Natl. Acad. Sci. USA* 93:7843-48, 1996. Nine cell lines engineered to produce antibodies of the invention were selected for further study and designated 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4, 8.10.3 and 9.7.2. The hybridomas were deposited under terms in accordance with the Budapest Treaty with the American Type Culture Collection (ATCC), 10801 University Blvd., Manassas, VA 20110-2209 on August 8, 2003. The hybridomas have been assigned the following accession numbers:
- | | |
|--------------------------------|----------|
| Hybridoma 3.8.3 (LN 15891) | PTA-5390 |
| Hybridoma 2.7.3 (LN 15892) | PTA-5391 |
| Hybridoma 1.120.1 (LN 15893) | PTA-5392 |
| Hybridoma 9.7.2 (LN 15894) | PTA-5393 |
| Hybridoma 9.14.4 (LN 15895) | PTA-5394 |
| Hybridoma 8.10.3 (LN 15896) | PTA-5395 |
| Hybridoma 88-gamma (UC 25489) | PTA-5396 |
| Hybridoma 88-kappa (UC 25490) | PTA-5397 |
| Hybridoma 100-gamma (UC 25491) | PTA-5398 |
| Hybridoma 100-kappa (UC 25492) | PTA-5399 |
| Hybridoma 252-gamma (UC 25493) | PTA-5400 |
| Hybridoma 252-kappa (UC 25494) | PTA-5401 |

EXAMPLE II

Gene Utilization Analysis

- [0274] DNA encoding the heavy and light chains of monoclonal antibodies 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4, 8.10.3 and 9.7.2 was cloned from the respective hybridoma cell lines and the DNA sequences were determined by methods known to one skilled in the art. Additionally, DNA from the hybridoma cell lines 9.14.4, 8.10.3 and 9.7.2 was mutated at specific framework regions in the variable domain and/or isotype-switched to obtain, for example, 9.14.4I, 8.10.3F, and 9.7.2IF, respectively. From nucleic acid sequence and predicted amino acid sequence of the antibodies, the identity of the gene usage for each antibody chain was determined ("VBASE"). Table 2 sets forth the gene utilization of selected antibodies in accordance with the invention:

Table 2
Heavy and Light Chain Gene Utilization

Clone	Heavy Chain				Kappa Light Chain		
	SEQ ID NO:	V _H	D _H	J _H	SEQ ID NO:	V _κ	J _κ
252	1, 2	3-11	7-27	6	3, 4	O12	3
88	5, 6	3-7	6-13	4	7, 8	O12	3
100	9, 10	3-23	1-26	4	11, 12	L2	3
3.8.3	14	3-11	7-27	4	16	L5	3
2.7.3	18	3-33	1-26	4	20	L5	4
1.120.1	22	1-18	4-23	4	24	B3	1
9.14.4I	25, 26	3-11	7-27	4b	27, 28	O12	3
8.10.3F	29, 30	3-48	1-26	4b	31, 32	A27	4
9.7.2IF	33, 34	3-11	6-13	6b	35, 36	O12	3
9.14.4	37, 38	3-11	7-27	4b	27, 28	O12	3
8.10.3	29, 30	3-48	1-26	4b	43, 44	A27	4
9.7.2	45, 46	3-11	6-13	6b	47, 48	O12	3
8.10.3FG1	97, 98	3-48	1-26	4b	31, 32	A27	4
9.14.4G1	101, 102	3-11	7-27	4b	27, 28	O12	3
9.14.4C-Ser	54	3-11	7-27	4b	56	O12	3
9.14.4-CG2	74	3-11	7-27	4b	56	O12	3
9.14.4-CG4	78	3-11	7-27	4b	56	O12	3
8.10.3C-Ser	58	3-48	1-26	4b	60	A27	4
8.10.3-CG2	62	3-48	1-26	4b	60	A27	4
8.10.3-CG4	94	3-48	1-26	4b	60	A27	4
8.10.3-Ser	90	3-48	1-26	4b	43, 44	A27	4
9.7.2C-Ser	50	3-11	6-13	6b	52	O12	3
9.7.2-CG2	66	3-11	6-13	6b	52	O12	3
9.7.2-CG4	70	3-11	6-13	6b	52	O12	3
9.7.2-Ser	86	3-11	6-13	6b	47, 48	O12	3
9.14.4-Ser	82	3-11	7-27	4b	27, 28	O12	3

- [0275] Mutagenesis of specific residues of the heavy and light chains was carried out by designing primers and using the QuickChange Site Directed Mutagenesis Kit from Stratagene, according to the manufacturer's instructions. Mutations were confirmed by automated sequencing, and mutagenized inserts were subcloned into expression vectors. The expression vectors were transfected into HEK293 cells to produce enough of the antibodies for characterization.

EXAMPLE III

M-CSF Mouse Monocytic Cell Proliferation Assay

- [0276] *In vitro* assays were conducted to measure M-CSF-dependent mouse monocytic cell proliferation in the presence of anti-M-CSF antibodies to determine the degree of inhibition by anti-M-CSF antibodies.
- [0277] Mouse monocytic cells, M-NFS-60 cells, from American Type Culture Collection (ATCC) (Manassas, VA), were obtained and maintained in RPMI-1640 medium containing 2 mM L-glutamine (ATCC), 10% heat inactivated fetal bovine serum (FBS) (Invitrogen, Carlsbad, CA), 0.05 mM 2-mercaptoethanol (Sigma, St. Louis MO) (assay medium), with 15 ng/ml human M-CSF. M-NFS-60 cells were split to 5×10^4 for next day use or to 2.5×10^4 for use in 2 days. Prior to use in the assay, the cells were washed three times with RPMI-1640, counted and the volume adjusted with assay medium to yield 2×10^5 cells/ml. All conditions were conducted in triplicate in 96-well treated tissue culture plates (Corning, Corning, NY). To each well 50 μ l of the washed cells, either 100 pM or 1000 pM M-CSF in a volume of 25 μ l and test or control antibody at various concentrations in a volume of 25 μ l in acetate buffer (140 mM sodium chloride, 20 mM sodium acetate, and 0.2 mg/ml polysorbate 80, pH 5.5) to a final volume of 100 μ l was added. Antibodies of the invention were tested alone and with human M-CFS. The plates were incubated for 24 hours (hrs) at 37°C with 5% CO₂.
- [0278] After 24 hrs, 10 μ l/well of 0.5 μ Ci ³H-thymidine (Amersham Bioseiences, Piscataway, NJ) was added and pulsed with the cells for 3 hrs. To detect the amount of incorporated thymidine, the cells were harvested onto pre-wet unifilter GF/C filterplates (Packard, Meriden, CT) and washed 10 times with water. The

plates were allowed to dry overnight. Bottom seals were added to the filterplates. Next, 45 μ l Microscint 20 (Packard, Meriden, CT) per well was added. After a top seal was added, the plates were counted in a Trilux microbeta counter (Wallac, Norton, OH).

- 5 [0279] These experiments demonstrate that anti-M-CSF antibodies of the invention inhibit mouse monocytic cell proliferation in response to M-CSF. Further, by using various concentrations of antibodies, the IC₅₀ for inhibition of mouse monocytic cell proliferation was determined for antibodies 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, and 9.7.2 (Cell
10 Proliferation Assay, Table 3a and Table 3b).

Table 3a

Antibody	252	88	100	3.8.3	2.7.3	1.120.1
M-CSF Mouse Monocytic Cell Proliferation Assay [IC ₅₀ , M]	1.86 x 10 ⁻¹⁰	2.31 x 10 ⁻¹⁰	7.44 x 10 ⁻¹⁰	7.3 x 10 ⁻¹¹	1.96 x 10 ⁻¹⁰	1.99 x 10 ⁻¹⁰
Human Whole Blood Monocyte Activation Assay [IC ₅₀ , M]	8.67 x 10 ⁻¹⁰	5.80 x 10 ⁻¹⁰	1.53 x 10 ⁻¹⁰	8.6 x 10 ⁻¹¹	7.15 x 10 ⁻¹⁰	8.85 x 10 ⁻¹⁰
Receptor Binding Inhibition Assay [IC ₅₀ , M]	7.47 x 10 ⁻¹⁰	4.45 x 10 ⁻¹⁰	1.252 x 10 ⁻⁹	7.0 x 10 ⁻¹¹	3.08 x 10 ⁻¹⁰	1.57 x 10 ⁻¹⁰

Table 3b

Antibody	9.14.4I	8.10.3F	9.7.2IF	9.14.4	8.10.3	9.7.2
M-CSF Mouse Monocytic Cell Proliferation Assay [IC ₅₀ , M]	2.02 x 10 ⁻¹⁰	4.13 x 10 ⁻¹⁰	7.37 x 10 ⁻¹⁰	2.02 x 10 ⁻¹⁰	4.13 x 10 ⁻¹⁰	7.37 x 10 ⁻¹⁰
Human Whole Blood Monocyte Activation Assay [IC ₅₀ , M]	2.49 x 10 ⁻¹⁰	4.46 x 10 ⁻¹⁰	1.125 x 10 ⁻⁹	6.48 x 10 ⁻¹⁰	2.8 x 10 ⁻¹⁰	1.98 x 10 ⁻¹⁰
Receptor Binding Inhibition Assay [IC ₅₀ , M]	2.97 x 10 ⁻¹⁰	9.8 x 10 ⁻¹¹	5.29 x 10 ⁻¹⁰	4.1 x 10 ⁻¹¹	1.5 x 10 ⁻⁹	6 x 10 ⁻¹²

5

EXAMPLE IVHuman Whole Blood Monocyte Activation Assay

[0280] *In vitro* assays were conducted to measure M-CSF dependent monocyte shape changes in the presence of anti-M-CSF antibodies to determine if the anti-M-CSF antibodies were capable of inhibiting whole blood monocyte activation and their degree of inhibition of monocyte shape changes.

[0281] In individual wells of a 96-well tissue culture plate, 6 μ l of 1.7 nM anti-M-CSF and 94 μ l of whole human blood for a final concentration of 102 pM anti-M-CSF antibody were mixed. The plates were incubated at 37°C in a CO₂ tissue

- culture incubator. Next, the plates were removed from the incubator. To each well, 100 μ l of a fixative solution (0.5% formalin in phosphate buffered saline without $MgCl_2$ or $CaCl_2$) was added and the plates were incubated for 10 minutes at room temperature. For each sample, 180 μ l from each well and 1 ml of Red Cell
- 5 Lysis Buffer were mixed. The tubes were vortexed for 2 seconds. Next, the samples were incubated at 37°C for 5 minutes in a shaking water bath to lyse the red blood cells, but to leave monocytes intact. Immediately following this incubation, the samples were read on a fluorescence-activated cell scanning (FACS) machine (BD Beckman FACS) and data was analyzed using FACS Station
- 10 Software Version 3.4.
- [0282] These experiments demonstrate that anti-M-CSF antibodies of the invention inhibit monocyte shape changes compared to control samples. Using the monocyte shape change assay, the IC_{50} was determined for antibodies 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, and 9.7.2
- 15 (Human Whole Blood Monocyte Activation, Table 3a and Table 3b).

EXAMPLE V

c-fms Receptor Binding Inhibition Assay

- [0283] *In vitro* assays were conducted to measure M-CSF binding to *c-fms* receptor in the presence of anti-M-CSF antibodies to determine if the anti-M-CSF antibodies were capable of inhibiting M-CSF binding to *c-fms* receptor and their degree of inhibition.
- [0284] NIH-3T3 cells transfected with human *c-fms* or M-NSF-60 cells maintained in Dulbecco's phosphate buffered saline without magnesium or
- 25 calcium were washed. NIH-3T3 cells were removed from tissue culture plates with 5 mM ethylene-diamine-tetra-acetate (EDTA), pH 7.4. The NIH-3T3 cells were returned to the tissue culture incubator for 1-2 minutes and the flask(s) were tapped to loosen the cells. The NIH-3T3 cells and the M-NSF-60 cells were transferred to 50 ml tubes and washed twice with reaction buffer (1x RPMI without
- 30 sodium bicarbonate containing 50 mM N-2-Hydroxyethylpiperazine-N'-2-ethanesulfonic acid (HEPES), pH 7.4). Next, the NIH-3T3 cells were resuspended

in reaction buffer for a final concentration of 1.5×10^5 cell/ml. The M-NSF-60 cells were resuspended in a reaction buffer for a final concentration of 2.5×10^6 cells/ml.

- [0285] For the assay, 9 μ l of a sterile 0.4 M sucrose solution, 100 μ l of 125 I-M-CSF (Amersham, IMQ7228v) at a final concentration of 200 pM in RPMI-1640 containing 50 mM HEPES (pH 7.4), 0.2% bovine serum albumin, and 100 μ l of unlabeled M-CSF at a final concentration of 200 nM were mixed in a binding tube. Next, 50 μ l/tube of increasing concentrations of a test antibody was added. In order to determine non-specific binding of the antibodies, we included samples to which we also added 200 nM M-CSF. To control tubes, we did not add antibody. Next, 15,000 NIH-3T3 cells or 250,000 M-NSF-60 cells were added per tube. All tubes were incubated at room temperature for 3 hrs and subjected to centrifugation at 10,000 rpm for 2 min. The tips of the tubes containing the cell pellets were cut off and the amount of M-CSF bound to the cells was determined using a Packard Cobra II Gamma counter. The specific binding was determined by subtracting non-specific binding from total binding. All assays were performed in duplicate. The binding data was analyzed using the computer program, Graph Pad Prism 2.01.

- [0286] These experiments demonstrate that anti-M-CSF antibodies of the invention inhibit the binding of M-CSF to *c-fms* receptor compared to control samples. Further, by using various concentrations of antibodies, the IC_{50} for inhibition of receptor binding was determined for antibodies 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, and 9.7.2 (Receptor Binding Inhibition Assay, Table 3a and Table 3b).

25

EXAMPLE VI

Determination of Affinity Constants (K_D) of Anti-M-CSF

Monoclonal Antibodies by BIAcore™

- [0287] Affinity measures of purified antibodies were performed by surface plasmon resonance using the BIAcore™ 3000 instrument, following the manufacturer's protocols.

30

- [0288] For antibodies 3.8.3, 2.7.3 and 1.120.1, the experiments were performed in a BIAcore™ 3000 instrument at 25°C in Dulbecco's phosphate buffered saline containing 0.0005% Tween-20. Protein concentrations were obtained from sedimentation velocity experiments or by measuring the wavelength of the sample at 280 nm using theoretical extinction coefficients derived from amino acid sequences. For experiments measuring the binding of antibody to immobilized antigens, M-CSF was immobilized on a B1 chip by standard direct amine coupling procedures. Antibody samples were prepared at 0.69 μ M for 3.8.3, 2.7.3 and 1.120.1. These samples were diluted 3-fold serially to 8.5 nM or 2.8 nM for roughly a 100-fold range in concentrations. For each concentration, the samples were injected in duplicate at 5 μ L/min flow for 4 min. The dissociation was monitored for 2000 seconds. The data were fit globally to a simple 1:1 binding model using BIAcore™ BIAevaluation software. In all cases, this method was used to obtain k_{off} and it was found that this data set compared well to data obtained from global fit of association and dissociation data.
- [0289] For antibodies 252, 88 and 100, the experiments were performed in a BIAcore™ 3000 instrument at 25°C in HBS-EP Buffer (0.01M HEPES, pH 7.4, 0.15 M NaCl, 3 mM EDTA, 0.005% Surfactant P20). For experiments measuring the binding of antibody to immobilized antigens, a M-CSF was immobilized on a CM5 Research Grade Sensor chip by standard direct amine coupling procedures. Antibody samples were prepared at 12.5 nM for antibodies 252 and 100 and at 25.0 nM for antibody 88. These samples were two-fold serially diluted to 0.78 nM for roughly a 15-30 fold range in concentrations. For each concentration, the samples were injected in duplicate in random order at 30 μ L/min flow for 3 min. The dissociation was monitored for 300 sec. The data were fit globally to a simple 1:1 binding model using BIAcore™ BIAevaluation software. In all cases, this method was used to obtain k_{off} and it was found that this data set compared well to data obtained from global fit of association and dissociation data.
- [0290] Table 4 shows results for antibodies 252, 88, 100, 3.8.3, 2.7.3 and 1.120.1.

Table 4

	252	88	100	3.8.3	2.7.3	1.120.1
K_D (M)	1.33×10^{-11}	1.33×10^{-9}	2.0×10^{-11}	4.0×10^{-10}	4.7×10^{-9}	5.4×10^{-9}
k_{off} (1/s)	1.03×10^{-6}	7.3×10^{-5}	1.7×10^{-5}			

EXAMPLE VIIProduction of 8.10.3 antibodies from 8.10.3 hybridoma cells

5 [0291] Antibody 8.10.3 was produced in 3L sparged spinners. The 3L sparged spinner flask is a glass vessel where cultures are mixed with an impeller controlled by a magnetic platform. The spinner is connected to gas lines to provide 5% CO₂ and air. 8.10.3 hybridoma cells were initially thawed into T-25 cell culture flasks. The cells were progressively expanded until there was a sufficient number of cells to seed the sparged spinners.

10 [0292] Two 3L sparged spinner flasks were seeded with 8.10.3 hybridoma cells in Hybridoma Serum-Free Medium with the additions noted on Table 5, for the two sparged flasks. The concentrations for Ultra low IgG serum (Gibco cat# 16250-078), L-glutamine (JRH Biosciences cat# 59202-500M), Non-Essential
 15 Amino Acids (Gibco cat# 11140-050), Peptone (Difco cat# 211693), glucose (In-house stock prepared from JT Baker cat# 1920-07), and Anti-foam C (Sigma cat.# A-8011) are given at their final concentrations in the media. The balance of the volume in each reactor is Hybridoma Serum-Free Medium.

20

Table 5. Conditions for Growing Hybridoma 8.10.3 in two 3L sparged spinners.

Conditions	Spinner 1	Spinner 2
Seeding density (1×10^6 cells/ml)	0.16 ml	0.16 ml
Hybridoma Serum-Free Medium (Gibco cat# 12045-076)	Balance	Balance
Ultra low IgG serum (Gibco cat# 16250-078)	5%	5%
L-glutamine (JRH Biosciences cat# 59202-500M)	8 mmol/L	8 mmol/L

Conditions	Spinner 1	Spinner 2
Non-Essential Amino Acids (Gibco cat# 11140-050)	1%	1%
Peptone (Difco cat# 211693)	1g/L	1g/L
2M glucose (In-house stock prepared from JT Baker cat# 1920-07)	8g/L	8g/L
Anti-foam C (Sigma cat.# A-8011)	1ml/L	1ml/L

[0293] The cultures were grown for 15 days and were harvested when the viability was below 20%. Viability was determined by trypan blue exclusion method with an automated cell counter (Cedex, Innovatis). Harvesting was accomplished by centrifugation and subsequent filtration. Clarified supernatant was obtained after centrifugation for 15 minutes at 7000 rpm and subsequent filtration with a sterile 0.22 μ m 4" Opticap Millipore filter (cat# KVSCO4HB3) into a 10L sterile TC-Tech bag (cat # P/N 12420 Bag Style CC-10-112420). The filtrate was then purified in the following example.

10

EXAMPLE VIII

Purification of an Anti-M-CSF Antibody

[0294] A Protein A column (Amersham Pharmacia) was prepped by washing with 3 column volumes of 8M Urea, followed by an equilibration wash with 20 mM Tris (pH 8). The final filtrate from Example VII was spiked with 2% v/v of 1M Tris pH 8.3 and 0.02% Na₂S₂O₃ before being loaded onto the Protein A column via gravity-drip mode. After load was complete, the resin was washed with 5 column volumes of 20 mM Tris (pH 8), followed by 5 column volumes of the elution buffer (0.1 M Glycine pH 3.0). Any precipitation was noted, and then a 10% v/v spike of 1M Tris pH 8.3 was added to the eluted antibody. The eluted protein was then dialyzed into 100 fold the volume amount of eluted material of dialysis buffer (140 mM NaCl/20mM Sodium Acetate pH 5.5). Following dialysis, the antibody was sterile filtered with a 0.22 μ m filter and stored until further use.

25

EXAMPLE IX

Monkey Treatment and Monocyte Counts

- [0295] One male and one female cynomolgus monkey per dosage group were intravenously administered vehicle or antibody 8.10.3 (produced as describe in
- 5 Examples VII and VIII) at 0, 0.1, 1, or 5 mg/kg in a dose volume of 3.79 mL/kg over an approximately 5 minute period. Blood samples for clinical laboratory analysis were collected at 24 and 72 hours postdose and weekly for 3 weeks. The monocyte counts were determined by light scatter using an Abbott Diagnostics Inc. Cell Dyn system (Abbott Park, Illinois).
- 10 [0296] A dose-related decrease (~25% to 85%) in total monocytes at all doses (Figures 1A and 1B) was observed. Monocyte counts at the 0.1 and 1 mg/kg appeared to rebound to near control levels by week 2, while monocyte counts at 5 mg/kg were still decreased at 3 weeks.

CD14+CD16+ monocyte subset analysis

- 15 [0297] Primate whole blood was drawn into Vacutainer tubes containing sodium heparin. 0.2 ml of each blood sample was added to a 15 ml conical polypropylene centrifuge tube containing 10 ml of red blood cell lysis buffer (Sigma), and incubated in a 37°C water bath for 15 minutes. The tubes were then centrifuged in a Sorvall RT7 centrifuge for 5 minutes at 1,200 rpm. The supernatant was
- 20 aspirated, the pellet resuspended in 10 ml of 4°C FACS buffer (Hanks' Balanced Salt Solution/2%FBS/0.02% sodium azide), and the tube centrifuged again for 5 minutes at 1,200 rpm. The supernatant was aspirated and the pellet resuspended in an antibody cocktail consisting of 80 µl 4°C FACS buffer, 10 µl FITC-conjugated anti-human CD14 monoclonal antibody (BD Biosciences, San Diego, CA), 0.5 µl
- 25 Cy5-PE-conjugated anti-human CD16 monoclonal antibody (BD Biosciences, San Diego, CA), and 10 µl PE-conjugated anti-human CD89 monoclonal antibody (BD Biosciences, San Diego, CA). The cell suspension was incubated on ice for 20 minutes, after which 10 ml of 4°C FACS buffer was added and the cells centrifuged as before. The supernatant was aspirated, and the cell pellet
- 30 resuspended in 400 µl FACS buffer and the cells analyzed on a FACSCaliber flow

cytometer (BD Biosciences, San Jose, CA). Data for 30,000 cells were collected from each sample.

- [0298] The monocyte population was identified by a combination of forward angle light scatter and orthogonal light scatter. Cells within the monocyte gate
- 5 were further analyzed for expression of CD14 and CD16. Two distinct population of monocytes were observed, one expressing high levels of CD14 with little or no CD16 expression (CD14++CD16-) and the other expressing lower levels of CD14, but high levels of CD16 (CD14+CD16+), similar to the two monocyte subsets previously described in human peripheral blood (Ziegler-Heitbrock H.W.,
- 10 *Immunology Today* 17:424-428 (1996)). For each primate tested, the percentage of monocytes within the CD14+CD16+ subset was determined after each blood draw, on days 1, 3, 7, 14, and 21 after 8.10.3 injection.
- [0299] In general, 8.10.3 treatment resulted in a reduction in the percentage of CD14+CD16+ monocytes (see Figures 2A and 2B). Monkeys not receiving 8.10.3
- 15 Antibody demonstrated relatively stable CD14+CD16+ monocyte levels. CD14+CD16+ monocytes have been termed "proinflammatory" because they produce higher levels of TNF- α and other inflammatory cytokines (Frankenberger, M.T., *et al.*, *Blood* 87:373-377 (1996)). It has also been reported that the differentiation of monocytes from the conventional CD14++CD16- phenotype to
- 20 the proinflammatory phenotype is dependent on M-CSF (Saleh M.N., *et al.*, *Blood* 85: 2910-2917 (1995)).

EXAMPLE X

Monkey Treatment and Monocyte Counts

- [0300] Three male cynomolgus monkeys per dosage group were intravenously
- 25 administered vehicle (20 mM Sodium acetate, pH 5.5, 140 mM NaCl), purified antibody 8.10.3F, or purified antibody 9.14.4I at 0, 1, or 5 mg/kg in a dose volume of 3.79 mL/kg over an approximately 5 minute period. The monkeys were 4 to 9 years of age and weighed 6 to 10 kg. Blood samples for clinical laboratory analysis were collected at 2, 4, 8, 15, 23, and 29 days. Monocyte counts were
- 30 determined by light scatter using an Abbott Diagnostics Inc. Cell Dyn system (Abbott Park, Illinois).

[0301] A decrease in the percentage change in total monocytes at all doses of antibody 8.10.3F and antibody 9.14.4I as compared to pre-test levels of monocytes (Figures 3A and 3B) was observed (see e.g., day 4, 8, 15, and 23 in Figures 3A and 3B).

- 5 [0302] All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference. Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be
- 10 readily apparent to those of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.

SEQUENCES

Key:

Signal peptide: underlined lower case

CDRs 1,2,3: underlined UPPERCASE

5 Variable domain: UPPER CASE

Constant domain: lower case

Mutations from germline in bold

SEQ ID NO: 1

10 252 Heavy Chain [Gamma chain] nucleotide sequence

atggagttggggctgtgctggatttctgttgctattataaaaggtgccagtgCAGGTGCAGCTGGTG
 GAGTCTGGGGGAGGCTTGGTCAAGCCTGGAGGGTCCCTGAGACTCTCC
 TGTGCAGCCTCTGGATTACCTTCAGTGACTACTACATGAGCTGGATCC
 GCCAGGCTCCAGGGAAGGGGCTGGAGTGGATTTTCATACATTAGTGGTA
 15 GTGGTAGTACCATATACTACGCAGACTCTGTGAAGGGCCGATTACCAT
CTCCAGGGACAACGCCAAGAACTCACTGTATCTGCAAATGAACAGCCT
GAGAGCCGAGGACACGGCCGTGTATCACTGTGCCAGAGCCCTGGGTGG
GATGGACGTCTGGGGCCAAGGGACCACGGTCACCGCTCTCCTCAGCTteca
 ccaaggcccatcgcttcccctggcgccctgctctagaagcactccgagagcacagcgccctggcgctgctg
 20 gtcaaggactactccccgaaccggtgacggtgtcgtggaactcaggcgctctgaccagcgcggtgcacacctccc
agctgtctacagctcctcaggactctactcctcagcagcgtggtgaccgtgccctccagcaacttcggcacccagac
 ctacacctgcaacgtatcacacaagccagcaacaccaaaggtggacaagacagttgagcgcaaatgtgtgtcgtgagt
 gccacccgtgcccagcaccacgtgtggcaggaacgtcagttcttcttcccccaaaacccaaggacacccctcatga
 tctcccgaacccctgaggtcacgtgcgtggtggtggacgtgagccacgaagaccccgaagtcacgttcaactgtgtac
 25 gtggacggcggtgaggtgcataatgccaagaacaagccacgggagggagcagtgtcaacagcagcttccgtgtggtca
gcgtctcctaccgttgtgcaccaggactggctgaacgcgaaggaggtacaagtgcaaggtcttccaacaaaggcctcca
gccccatcgagaaaaccatctccaaaaccaaaggcgagcccgagaaccacagggtgtacacctgccccatccc
gggagggagatgaccagaacaggtcagcctgacctgcctgctgtaaaggcttctatcccagcgacatcgccggtgga
gtgggagagcaatggcgagccggagaacaactacaagaccacacctccatgctggaactcgacggtcctctctt
 30 ctacagaaagctaccgtggacaagcaggtggcagcaggggaacgtctctctatgctccgtgatgcatgaggct
ctgcacaacctacacgcagaagcctctccctgtctccgggtaaa

SEQ ID NO: 2

252 Heavy Chain [Gamma chain] protein sequence

melglwiflvaitkgvqcQVQLVESGGGLVKPGGSLRLSCAASGFTESDYYMSWTR
 35 QAPGKGLEWISYISGSGSTIYADSVKGRFTISRDNAKNSLYLQMNSLRAE
DTAVYHCARALGGMDVWGQGTTVTVSSAstkgpsvflpapcsrstsestaalgclvkdyp
epvtvswnsgtsgvhtfpavlqsglyslssvvtvpsnfgtqtytcnvdhkpsntkvdktverccvecppep
appvagpsvflfppkpkdtlmisrtpevtcvvdvshedpevfqnvydgvevhnaktkpreeqfnstfrvvsv
 40 ltvhqdwlngleyckvsnklpapiektsktkgqprepqvytlppsreemtknqvsltclvgfypsdiave
wesngqpennykttppmldsgsflyskltvdksrwqgnvfscvmhealhnhytqkslslspgk

SEQ ID NO: 3

252 Light Chain [Kappa chain] nucleotide sequence

atgagggtccctgctcagctcctggggcctcctactctgctccgaggtgccagatgtGACATCCAGAT
 ATCCAGCTCTCCATCCTCCCTGTCTGTCATCTGTAGGAGACAGAGATCAAC
 5 ATCACTTGCCGGGCAAGTCAGAGCAATTAGCGGCTTTTAAATTTGGTATC
 AGCAGAAACCAGGGAAGGCCCTAAGCTCCTGATCTATGCTACATCCA
GTTTGCAAAGTGGGGTCCCATTCAGGTTCAAGTGGCAGTGGATCTGGGA
 CAGATTTCACTCTCAACATCAGCAGTCTGCAACCTGAAGATTTTGCAAC
 TTATTACTGTCAACAGAGTTACAGTGTCCCATTCACTTTCGGCCCTGGG
 10 ACCAAAGTGGATATCAAACGAactgtggetgcaccatctgtctatctccgccatctgatgagc
 agttgaatctgggaactgctagcgttgtgtgctctgtaataactctatcccagagagccaaagtacagtggaaggt
 ggataacgccctccaatcggtgaactcccagagaggtgtcacacagcaggacagcaaggacagcacctacagccctc
 agcagcaccctgacgctgagcaaaagcagactacgagaacacaaagtctacgctgcgaagtccatccatcagggcc
 tgaactgcgccgtcacaaagagcttcaacaggggagagtggt

15

SEQ ID NO: 4

252 Light Chain [Kappa chain] protein sequence

mrvpaqlglllwlrarcdIQMTQSPSSLASVGDVRVTITCRASQISGFLNWWYQQK
 PGKAPKLLIYATSSLOSGVPRFSGSGSDFTLTISSLQPEDFATYYCQQS
 20 YSVPFTFGPGTKVDIKRtaapsvtfppssdeqlksgtasvrlmhfpreakvqwkvdnlqsgns
 qesvteqskdstylssltflskadyekhkvyacevthqlsspvtksfirgce

SEQ ID NO: 5

88 Heavy Chain [Gamma chain] nucleotide sequence

atggaaattggcgtgctgctgggttttctctgtgctattttgaaggtgtccagtgGAGGTGCAGCTGGTG
 GAGTCTGGGGGAGGCTTGGTCCAGCCTGGGGGGTCCCTGAGACTCTCC
 25 TGTGCAAGCCTCTGGATTACCTTTAGTAGCTATTGGATGAGCTGGGTCC
 GCCAGGCTCCAGGGAAGGGGCTGGAGTGGGTGGCCAACATAAAGCAA
GATGGAAGTGAGAAATACTATGTGGACTCTGTGAAGGGCCGATTACAC
 ATCTCCAGAGACAACGCCAAGAACTCACTGTATCTGCAAAATGAACAGC
 30 CTGAGAGCCGAGGACACCGCTGTATTACTGTCTCCGGGTATAGCA
GCAGCTGGTAGGGCTACTTGGGGCCAGGGAACCTTGGTACCGTCTCC
 TCAAGTccacaagaggcccatccgtcttccctcctggcgccctgctctagaagcactccctgagagacacagcggc
 cctggctgctcgtgtcaaggactacttcccggaaccgtgacggtgtcgtggaactcaggcgtctgacacagcggc
 35 tgcacacctccagctgctctacagctcctcagcactctccctcagcagcgttggtgacgttgccctccagcaactc
 ggcaccca gacctacactgcaacgtatgctacaaagccca gcaacaaagggtgacaa gacagtgtagcgcaaat
 gttgtgtcagtgccaccgtgccaccgtgacccagcaccacctgtggcagggaccgtcagctctctctctcccccacaaacccaagg
 acacctcatgctctccggaccctgaggtcacgtgcgtgtgtgtgtgacgtgagccaaagaagaccocagggtgtcca
 40 gttcaactgtacgttgacggcgctgtggaggtgcataatgccaaagacaaggccacgggagaggacgttcaacagcacg
 ttccgtgtgtgtcagcgtctcaccgttggcacccaggactggtgacggcaaggtgacaaaggtgcaagggtctccaac
 aaaaggctccagccccctcagaaaaacatctccaaaaccaaaggcgcccgagcccgagaaaccaaggtgtacacc
 ctgcccccatcccgaggagatgacaaagaaccaggtgacgcttgacctgctggtcaaaaggctctaccctccagcg
 acatcgccgtggatgtgggagagcaatggcgacggcgagaaacactacaaagaccacacctccatgctggaactcg
 acggctcctctctctctacagcaagctcacogtggacaagagacaggtggcagcaggggaacgtctctcatgctccg
 45 tgatgatgaggctctgcacaaccactacacgcagaagacctctccctgctctcgggtaaa

SEO ID NO: 6

88 Heavy Chain [Gamma chain] protein sequence

```
mefglzfvvfaileqgevfQVLVESGGGLVPGGSLRLSQAASGFTSSYVMSWV
RQAPGKGLVWVANKVQKSGKYYVDSVKGKRTISDRDNAKNSLYQMNSI
RAEDTAVYYCAPGIAAAAGRAYWGQGLTIVSSAStkqpsvfpIapcprsrstsealgc
vldkfypcpvtswnsgalsvhtpafvqsgsllysvvsspsfghyqthvdkhpsktvdktrvck
ecppcpvpgvagsvffppkpldmlimstrvccvddshvqfwmvdyvdeghnaktcpreefms
tdrsvsvltvwhdglwlnkyckvksnslgapiaktictskqpprepytppsreemtnqlvsklqfygfp
sdvavsewngngpennqytktpmldsdgffsllyskvdkswqgqvfscsmhealhnlqtslslspk
```

SEO ID NO: 7

88 Light Chain [Kappa chain] nucleotide sequence

atgaggctccctgctcagcctctgggctctctgctactctggctcagagtgccagatgTGACATCCAGAT
GACCCAGTCTCCATCTCCCTCTCCCTGCTGCATCTGTTGGAGACAGAGTACC
ATCACTTCGCCGCCAAGTCAGGACATTTAGCAGTATTATTAATTTGGTATC
AGCAGAAACAGGGAAAGCCCTAAGCTCTGATCTATGCTGCATCCA
GTTTGCAAAGTGGGGTCCCATTAAGTTCTAGTGGCAGTGGATCTGGGA
CAGATTCTCACTCTCAACCTACAGCACTGTCAACCTGAAAGATTTTGCAAC
TTACTCTGTGCAACAGGATACACCCATTCACTTTCGGCCCTGGG
ACCAAAGTGGATATCAACAAGAActgtggtgcacactgtcttcattctccgcactcagtaggc
agttgaatactgaaactgtacgtgtgtgtgctgctgaataactttatccagagagcgaagactacagtgaagt
ggataacgccctccaatcgggttaactccagagagagtgtcacagagcaggacagcaagacacacctacagcctc
agcagcaacctgacgtgtgcagaagcagcactacgaaacacaaagtctacgctgcgaagtccaccatcaggccgc
taacgtcccccgtcacaagaagctctcaacagcagcagctgt

SEO ID NO: 8

88 Light Chain [Kappa chain] protein sequence

mrvpagllgllllwlgarcDIQMTQSPSSLSASVGDRVTITCRPSODISSYLNWYQQK
PGKAPKLLIYAAASSLSQGVLPRFSGSGSGTDFTLVLTISSLQPEDFATYYCQQS
YSTPTFFGPGTKVDIKRvalprrfifppsdqgksgtasvclnnfyprcakvqwkvdnalgsgns
yestvetdskdstvslstlltskadvekhkvvacevthqslsnvtsfirgec

SEQ ID NO: 9

100 Heavy Chain [Gamma chain] nucleotide sequence

atggagtttggcgctccgctggattttcttctgtgctatlttaaaaggtgccaagtgAGGTCGACGTGTTG
 5 GAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCC
 TGTGTCAGCCTCTGGATTACACCTTTAGCAGCTATGCCATGAGCTGGGTCC
 GCCAGGCTCCAGGGAAGGGGCTGGAATGGGTCTCAGCTATTAGTGGTC
GTGGTGGTAGGACATACTTCGCAGACTCCGTGAAGGGCCGGTTCACCA
 10 TCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAATGAACAGCC
 TGAGAGCCGAGGACACGGCCGTATATTTCTGTGCGGTAGAAGGCTATA
GTGGGCGCTACGGATTTTTGACTACTGGGGCCAGGGAACCTAGTCAC
 CGTCTCCTCAGCCTccaccaagggcccatcggtcttccctggcgccctgctctagaagcacctccgag
 agcacagcgccctgggctgcctgggtcaaggactacttcccgaaccggfagcggfctcgtgggaactcaggcgctct
 gaccagcgcggtgcacacctccagctgtctacagtcctcaggactctactcctcagcagcgtgtgaccgtgc
 15 ctccagcaacttggcacccagacctacacctgcaacgtatgacacaagccagcaacaccaaggtgacaagaca
 gtgagcgcgaatgttggcgtgagtgccaccgtgcccagcaccacctgtggcaggaocgtacgtctctcttctcccc
 caaaacccaaggacacctcatgctctccggaccctgaggtcacgtgcgtggtgtggtgacgtgagccacgaaga
 ccccgaggtccagttcaactgtacgtggacggcgtggaggtgcataatgccaaagacaagccacgggaggaaga
 gttcaacagcacgttccgtgtgtgacgtcctcaccgttggcaccagactggtgaacggcgaaggatagaagt
 20 caaggcttccaaaggtcctccagcccccacfcagaaaaacatctccaaaccaaaggcgacgccccgagaacc
 acaggtgtacacctgccccatccgggaggaagatgaccaagaaccaggtcagcctgacctgctggtcaaggc
 tcttaccoccgacatcgccgtggaagtgggaagagcaatggcgacgggaagaacaactacaagaccacacctccca
 tgcctgacctccagcggtcctctctctacagcaagctcaccgtggacagaagcaggtggcgacagggggaacgtc
 ttctcatgctccgtgatgcatgaggtcctgcacaaccactacacgcagaagagcctctcctgtctccgggtaaa
 25

SEQ ID NO: 10

100 Heavy Chain [Gamma chain] protein sequence

mefelrwiflvailkqvqcEVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVR
 30 QAPGKGLEWVSAISGRGRITYFADSVKGRFTISRDN SKNTLYLQMNSLR
 EDTAVYFCAVEGYSGRYGFFDYWGQGLTVTVSSAstkpsvfpfapcserstsestaal
 gclvldkyfpepvvtvswngaltsgvhtfpavlqssgylssvvtpssnfgtqtytcnvdhkpsntkvdktervk
 cveccppcpappvagsvflfppkpkdltlmsrtpvtcvvvdvshedpevqfnwvydgvvevhnaktkpreeqf
 nstfrvsvltvvhqdwlngkeyckvsnkglpapietkisktgqprepvtylppsreemtknqvsitclvkqf
 35 ypsdiawewesngspemnyktppmldsdgsfflyskltvdksrwqqgnvfcsvmhleahlnhytqkslslspg
 k

SEQ ID NO: 11

100 Light Chain [Kappa chain] nucleotide sequence

- 5 atggaagcccccagctcagcttctctctctgctactctggctcccagataccactggaGAAATAGTGATG
ACGCAGTCTCCAGGCCACCCTGTCTGTGTCTCCAGGGGAAGAGGCCACC
CTCTCTGCAAGGGCCAGTCAGAGTGTAGCAGCAACTTAGCCTGGTACC
AGCAGAAACCTGGCCAGGCTCCCAGGCTCCTCATCTATGGTGCATCCAC
10 CAGGGCCAGTGGTATCCAGACAGGATCAGTGGCAGTGGGTCTGGAAC
AGAGTTCACTCTCATCATCAGCAGCCTGCAGTCTGAAGATTTGCAGTT
TATTACTGTCAGCAGTCTAATAACTGGCCATTCACTTTCGGCCCTGGGA
CCAAAGTGGATATCAAACGAactgtgctgcaccatctgtctcattctccgccatctgatgagca
gttgaaatctggaactgctagcgtgtgtgctgctgaataactctatcccagagaggccaaaglacagtggaggtg
gataacgccctcaatcgggtaactccaggaagtgatcacagagcagagacagcaaggacagcacctacagcctca
gcagcaccctgacgtgagcaagcagactacgagaacacaaagctacgcctgcgaagtcacccatcagggcct
15 gagctcgcctgtcacaagagcttcaacaggggagagtgt

SEQ ID NO: 12

100 Light Chain [Kappa chain] protein sequence

- 20 meapaqlflflwlpdtteEIVMTQSPATLSVSPGERATLSCRASQSVSSNLAWYQQ
KPGQAPRLLIYGASTRASGIPDRISGSGSTEFTLIISLQSEDFAVYYCQQS
NNWPFTFPGTKVDIKRtvaapsvfifppsideqlksgtasvcllnnfypreakvqwkvdnalqsgn
sgsvteqdsksdstylssstltliskadyekhlkvacevthqglsspvtksfirgce

SEQ ID NO: 14

- 25 3.8.3 Heavy Chain [Gamma chain] protein sequence

- mefglsvvflvaiikvqgcQVQLVESGGGLVKPGGSLRLSCAASGFTFSDYYMSWI
RQAPGKGLEWFSYISSSGSTIYYADSVKGRFTISRDNAKNSLSLQMNLSRLA
EDTAVYYCARGLTGDYWGQGLTVTVSSAstkgpsvflapcsrcstestaaigclvkdyfpe
pvtvswngaltsgvhtfpavlqssglyslssvvtvpssnfgtqytencvdkhpsntkvdktverkkcvecppepa
30 pvvagpsvflfppkpkdltlmisrtpevtcvvdshedpeqvfwyvdgvevhnaktipreeqfnstfrvsvlt
vvhqdwlnqkeykckvsnkglpapietkisktgprepqvyltppsreemtknqvslclvkgfypsdiavew
esngqpennyktppmldsdgsfllyskltvdksrwqqgnvfscsvmhealnhhytqkslslspgk

SEQ ID NO: 16

- 35 3.8.3 Light Chain [Kappa chain] protein sequence

- mdmrvaqllglflwfpgrcDIQMTQSPSSVSASVGDRTISCRASQDISGWLAWY
QOKPGKAPKLLISATTSSLHSGVPSRFSGSGSGTDFTLTIISLQPEDFATYYC
QQTNSFPFTFGPGTKVDIKRtvaapsvfifppsideqlksgtasvcllnnfypreakvqwkvdnalq
sgnsqsvteqdsksdstylssstltliskadyekhlkvacevthqglsspvtksfirgce

SEQ ID NO: 18

2.7.3 Heavy Chain [Gamma chain] protein sequence

5 mefglswvflvallrceqQVQLVESGGGVVQPGRSRLRLSCAASGFTFSSYGMHWV
 RQAPGKGLEWVAFIWYDGSNKYYADSVKGRFTISRDNSKNTLYLQMNSL
 RAEDTAVYYCARGYRVYFDYWGQGLTVTVSSAstkgpsvfplapcsrcstseaa
 lgcldkdyfpepvtvswngaltsgvhtfpavlgssglyslssvvtvpssslgtktytcnvdhkp
 sntkvdkrveskygp
 pcpcspapeflggpsvflfppkpkdtlmisrtpetvcvvdvsqedpevqfniwyvdgvev
 hnaktkpreeqfns
 10 tyrvsvltvlhqdwlngkeykckvsnkglpssiektiskakgqprepqvytippscqe
 mtknqvsltlcvkgfy
 psdiavewesngqpennykttppvldsdgsfflysrltvdksrwqegnvfscsvmh
 ealhnhytqkslspsgk

SEQ ID NO: 20

2.7.3 Light Chain [Kappa chain] protein sequence

15 mdmrvpaaqlglgllwfpgsrCDIQTQSPSSVSASVGDRTITCRASQDISSWLAWY
 QRKPGKAPKLQIYAASLESQVPSRFNGSGSGTDFTLSSISLQPEDFATYYC
 QQTINSEPLTFGGGGTKVEIKRtvaapsvfi fppsdeqlksgtasvcllnfy
 preakvqwkvdnal
 qsgnsqesvteqdskdstyslsstltlskadyekhkvyacevthqglsspvtksfur
 gec

SEQ ID NO: 22

20 1.120.1 Heavy Chain [Gamma chain] protein sequence

mewtwsflfvaaatgahsQVQLVQSGAEVKKPGASVKVSCASGYTFTSYGISWV
 RQAPGQGLEWMGWISAYNGNTNYAQKLQDRVTMTTDTSTTTAYMELRS
 LRSDDTAVYYCARRAYGANFFDYWGQGLTVTVSSAstkgpsvfplapcsrcstseaa
 lgcldkdyfpepvtvswngaltsgvhtfpavlgssglyslssvvtvpssnfgtqtytcnvdhkp
 sntkvdktverk
 25 ccvecppcpappvagsvflfppkpkdtlmisrtpetvcvvdvshdepevqfniwyvdgvev
 hnaktkpree
 qfnstfrvsvltvlhqdwlngkeykckvsnkglpapiektisktkgqprepqvytipp
 sreemtknqvsltlcvk
 gfy psdiavewesngqpennykttppmldsdgsfflyskltvdksrwqqgnvfscsvmh
 ealhnhytqksls
 pgk

30

SEQ ID NO: 24

1.120.1 Light Chain [Kappa chain] protein sequence

mvlgatqvflslllwisgavGDIVMTQSPDSLAVSLGERATINCKSSQSILFFSNKNYL
 AWYRQKPGQPPNLLIYWASTRESQVPSRFNGSGSGTDFTLTISSLQAEQVA
 35 VYYCQQYYSSPWTFGGQGTKVEIKRtvaapsvfi fppsdeqlksgtasvcllnfy
 preakvq
 wkvdnalqsgnsqesvteqdskdstyslsstltlskadyekhkvyacevthqglsspvtksfur
 gec

SEQ ID NO: 25

9.14.4I Heavy Chain [Gamma Chain] nucleotide sequence

- atggagtttggcctgagctgggttttcttgttgcattataaaaggtgCCAGTGTCAAGTGCAGCTG
- 5 GTGGAGTCTGGGGGAGGCTTGGTCAAGCCTGGAGGGTCCCTGAGACTC
 TCCTGTGCAGCCTCTGGATTACCTTCACTGACTACTATATGAGCTGGA
 TCCGCCAGGCTCCAGGGAAGGACTGGAGTGGGTTCATACATTAGTA
GTAGTGGTAGTACCATATACTACGCAGACTCTGTGAAGGGCCGATTCA
- 10 CCATCTCCAGGGACAACGCCAAGAACTCACTGTATCTGCAAATGAACA
 GCCTGAGAGCCGAGGACACGGCCGTGTATTACTGTGCCGAGAGGCCTAA
CTGGGGACTACTGGGGCCAGGGAACCCTGGTCAACCGTCTCCTCAGCTacc
 accaaggggccatccgtcttcccttggcgccctgctctagaagcacctccgagagcagcagcgccctgggctgct
 ggtcaaggactacttccccgaaccggf gacggf gtcgt ggaactcagcgcgtctgaccagcggcgtgcacacctcc
 cagctgtctctacagtctcaggaactctactccctcagcagcgt ggtgacogtgcctccagcaactcggcaccaccaga
- 15 cctacacctgcaacgtatgacacaagcccagcaacacaaaggf gacaagacagf gacgcgcaaatgttgtc gag
 tgccccacgtgccagaccacctgtggcaggacgcgtcagctcttcttccccccaaaacccaaggacacctcatg
 atctccccggaccctgaggtcacgtgcgtggf ggtggacgtgagccacgaagaccocagggtccaggtcaactggta
 cgtggacggcgtggaaggtgcataatgccaaagacaagccacgggagagcaggttcaacagcagcttccgtgtggtc
 agcgtctctaccggtgtgtgcaccaggactggctgaacggcaaggagtaacaagf gcaaggtctcacaagaaggcctccc
- 20 agcccccatcgagaaaaccatctccaaaaccaaaggcgacgccccgagaaccacaggtgtacacctgcccccattcc
 cggggagagatgaccaagaaccagggtcagcctgacctgcctgttcaaaagcttctaccccagcgacatcgccgttg
 agtgggagagcaatggcgaccggagaacaactacaagaccacacctccatgctggacctccgacggctccttctc
 ctctacagcaagctcaccgtggacaaagacaggtggcagcaggggaaacgtcttctcatgtctcgtgcatgcatgaggc
 tctgcacaaccactacacgcagaagacgtctcctctctccgggtaaa

25

SEQ ID NO: 26

9.14.4I Heavy Chain [Gamma Chain] protein sequence

- mefglswvflvaiikgvacQVQLVESGGGLVKPGGSLRLSCAASGFTFSDYYMSWI
 RQAPGKGLEWVSYISSSGSTTYADSVKGRFTISRDNAKNSLYLQMNSLRA
- 30 EDTAVYYCARGLTGDYWGQGLTVTVSSAstkpsvflpaplcsrstsastaalgcflvkdyfpe
 pvtvswngaltsghvhtfpavlqssglyslssvvtvpsnftgtyctcnvdhkpnsntkvdtkverkecevepcppa
 pvtvagsvflfpkpdkdlmistrtpcvttcvvdshedpevqfwnvyvdgvevhnaktkpreeqfmsftfrvsvlt
 vvhqdwlngkeykckvsnkglpapiektiskdkgprepqvyltppsreemtknqvsltelvkglfypsdiavew
 esngqpennyktpmldsdgsfflyskltvdksrwqqgnvfscsvmhealnhnytklsislspgk

35

SEQ ID NO: 27

9.14.4, 9.14.4I, 9.14.4-Ser and 9.14.4-G1 Light Chain [Kappa Chain] nucleotide sequence

- 5 atggacatgagggtccccgctcagctcctgggctcctgctactctggctccgaggtccagatgTGACATCC
 AGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTCGGAGACAGAGT
 CACCATCACTTGCCGGCCAAAGTCAGATCATTAGCAGTTTATTAAATTGG
 TATCAGCAGAAACCAGGGAAGCCCCCTAAGCTCCTGATCCATGCTGCA
 10 TCCAGTTTGCAAAGTGGGGTCCCATCAAGGTTCAGTGGCAGTGGATCTG
 GGACAGATTTCACTCTCACCATCAGTAGTCTGCAACCTGAAGATTTTGC
 AACTTACTACTGTCAACAGAGTTACAGTACCCCATTCACTTTCGGCCCT
 GGGACCAAAGTGGATATCAAACGAactgtggctgcaccatctgtctcatcttcccgccatctga
 15 tgagcagttgaaatctggaactgcctctgtgtgtgcctgctgaataacttctatcccagagagggccaaagtacagtgg
 aggtggataacgccctccaatcgggtaactccagagagagtgacagagcaggacagcaagagacgacacctaca
 gcctcagcagcaccctgacgtgagcaaaagcagactacgagaacacaaagtctacgcctgcgaagtcacccatca
 gggcctgagctgcgccgtcacaaagagcttcaacaggggagagtg

SEQ ID NO: 28

9.14.4, 9.14.4I, 9.14.4-Ser and 9.14.4-G1 Light Chain [Kappa Chain] protein sequence

- 20 mdmrvpagllglllwlrgarcDIQMTQSPSSLSASVGDRTTITCRPSQIISSLLNWYQ
 QKPGKAPKLLIHAASSLQSGVPSRFSGSGSGTDFTLTISSLPEDFATYYCQ
QSYSTPFTFGPGTKVDIKRtaapsvfifpssdeqlksgtasvvelnnfybreakvqwkvdnalqs
 gnsqesvteqdskdstysstltlskadyekhkvyacevthqglsspvtksfirgec

25

SEQ ID NO: 37

9.14.4 Heavy Chain [Gamma Chain] nucleotide sequence

- atggaagtttggcgctgagctgggttttcttgttgcattataaaaggtatCCAGTGTCAAGGTGCAGCTG
 GTGGAGTCTGGGGGAGGCTTGGTCAAGCCTGGAGGGTCCCTGAGACTC
 5 TCCTGTGACAGCCTCTGGATTACCTTCAGTGACTACTATATGAGCTGGA
TCCGCCAGGCTCCAGGGAAGGGACTGGAGTGGGTTTCATACATTAGTA
GTATGGTGTAGTACCATATACTACGCAGACTCTGTGAAGGGCCGATTCA
CCATCTCCAGGGACAAAGCCCAAGAACTCACTGTATCTGCAAAATGAAACA
GCCTGAGAGCCGAGGACACGGCCGTGTATTACTGTGCGAGAGGGCCTAA
 10 CTGGGGACTACTGGGGCCAGGGAACCCCTGGTCAACCGTCTCTCAGCTTcc
accaaggggcccatcogtcttccocctggcgccctgctctagaagcacctccgagagcacagcggccctgggctgcct
ggtaaggaactacttcccgaaccggtgacgggtgctgfggaactcaggcgctctgaccagcggcgtgcacacctcc
cagctgtctcactagttcctcaggaactctactccctcagcagcgtggtgaccgtgccctccagcagcttgggcacgaaga
octacacctgcacgtagatcacaaagccagcaacaccaaaggtggacaagagagttagtccaaatatggtcccca
 15 tgcocatcatgccagcactgagttcctggggggaccatcagttcctctgttccccccaaaacccaaggacactctca
tgatctccggaccctgaggctcagctgcgtgtggtggagctgagccaggaagaccccgaggctcagttcaactgg
tacgtggatggcgtggaggtgcataatgccaaagacaaagccgggaggaagcagttcaacagcacgactacogtctggtg
tcagctctctacccgtctgcaccaggaactggctgaacggcaaggagtagacaagtgcaaggctctcaacaagggcctc
ccgtctccatcgagaaacatctccaaagccaaaggcagccccagagaccacaggtgtacacacctgcccccac
 20 cccaggaaggagatgacacaagaaccaggtcagcctgacctgcctgtcaaaagcttctaccaccagcgacatccgctg
ggagtgaggagacaatgggcagccggagaacaactacaagaccacgctcccgctgctggaactccgacgctccttc
ttctctacagcagctaacctggacaagacaggtgacagagggggaatgtctctcatgctccgtgatcgatgag
gctctgcacaaccactacacagagaagcctctccctgctctccgggtaaa

25 SEQ ID NO: 38

9.14.4 Heavy Chain [Gamma Chain] protein sequence

- mefglswvflvaiikgvqcQVQLVESGGGLVKPGGSLRLSCAASGFTFSDYYMSWI
 RQAPGKGLEWVVSYSISGSIYYADSVKGRFTISRDNAKNSLYLQMNSLRA
 EDTAVYYCARGLTGDYWGQGTLLVTVSSAstkgpsvfplapcrrstestaalgclvkdyfpe
 30 pvtvswngaltsghvthfpavlqssglyslssvvtvpssslgtktytncndvhkpsntkvdkrveskygppcppepa
peflggpsvflfppkpkdltlmisrtpvctvvdvsqdepvqfinwvvdgvevhnaktkpreeqfinstyrvsvl
tvllhqdwlnqkeykckvsnkglpssiektiskakgqprepqvyltppsgeemtknqvsltlcvkgfypsdiaev
wesngqpennytktppvldsdgsfflysrllvdkrswqegnvfscsvmhealhnhytqkslslspgk

35 SEQ ID NO: 54

9.14.4C-Ser Heavy Chain [Gamma chain] protein sequence

- mefglswvflvaiikgvqcQVQLVESGGGLVKPGGSLRLSCAASGFTFSDYYMSWI
 RQAPGKGLEWVVSYSISGSIYYADSVKGRFTISRDNAKNSLYLQMNSLRA
 EDTAVYYCARGLTGDYWGQGTLLVTVSSAstkgpsvfplapcrrstestaalgclvkdyfpe
 40 pvtvswngaltsghvthfpavlqssglyslssvvtvpssslgtktytncndvhkpsntkvdkrveskygppcppepa
peflggpsvflfppkpkdltlmisrtpvctvvdvsqdepvqfinwvvdgvevhnaktkpreeqfinstyrvsvl
tvllhqdwlnqkeykckvsnkglpssiektiskakgqprepqvyltppsgeemtknqvsltlcvkgfypsdiaev
wesngqpennytktppvldsdgsfflysrllvdkrswqegnvfscsvmhealhnhytqkslslspgk

SEQ ID NO: 56

9.14.4C-Ser, 9.14.4-CG2 and 9.14.4-CG4 Light Chain [Kappa chain] protein sequence

- 5 mdmrvpagllgllllwlrgrarcDIQMTQSPSSLSASVGDRTTTCRPSQIHSSLLNWFYQ
QKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFLTITSSLQPEDFATYYCQ
QSYSTPFTFGPGTKVDIKRtvaapsvfifppsdeqlksgtasvvcllnnfypreakvqwkvdnalqs
gnsqesvteqdsdktstyslsstflskadyekhkvayacevthqglsspvtksfmrgec

10 SEQ ID NO: 74

9.14.4-CG2 Heavy Chain [Gamma chain] protein sequence

- mefglswvflvaikgvqcQVQLVESGGGLVKPGGSLRLSCAASGFTFSDYYMSWI
RQAPGKGLEWVSYISSSGSTIYYADSVKGRFTISRDNNAKNSLYLQMNSLRA
EDTAVYYCARGLTGDYWGQGTLVTVSSAstkgpsvfplapcsrcstseaalglvkdypfe
 15 pvtvswngaltsgvhtfpavlqssgylsylvtpssngtqytencvdhkpntkvdtkverkcceppcpa
 ppvagpsvflfppkpkdtlmsirtpevtcvvdvshedpevqfhwvydgvgevhnaaktpreeqfnstfrvsvlt
 vvhqdwlngkeykckvsnkglpapiektisktkgqprepqvytlppsreemtknqvsitclvkgfypsdiavew
 esngqpennnyktppmldsdgsfflysrftvdksrwqegnvfscsvmhleahhnytkqskslspgk

20 SEQ ID NO: 78

9.14.4-CG4 Heavy Chain [Gamma chain] protein sequence

- mefglswvflvaikgvqcQVQLVESGGGLVKPGGSLRLSCAASGFTFSDYYMSWI
RQAPGKGLEWVSYISSSGSTIYYADSVKGRFTISRDNNAKNSLYLQMNSLRA
EDTAVYYCARGLTGDYWGQGTLVTVSSAstkgpsvfplapcsrcstseaalglvkdypfe
 25 pvtvswngaltsgvhtfpavlqssgylsylvtpssngtqytencvdhkpntkvdtkveskygppcpcpa
 peflggpsvflfppkpkdtlmsirtpevtcvvdvsqdedpevqfhwvydgvgevhnaaktpreeqfnstfrvsvlt
 tvlhqdwlngkeykckvsnkglpssiektiskakgqprepqvytlppsgeemtknqvslclvkgfypsdiave
 wesngqpennnyktppvldsdgsfflysrftvdksrwqegnvfscsvmhleahhnytkqskslspgk

30 SEQ ID NO: 82

9.14.4-Ser Heavy Chain [Gamma chain] protein sequence

- mefglswvflvaikgvqcQVQLVESGGGLVKPGGSLRLSCAASGFTFSDYYMSWI
RQAPGKGLEWVSYISSSGSTIYYADSVKGRFTISRDNNAKNSLYLQMNSLRA
EDTAVYYCARGLTGDYWGQGTLVTVSSAstkgpsvfplapcsrcstseaalglvkdypfe
 35 pvtvswngaltsgvhtfpavlqssgylsylvtpssngtqytencvdhkpntkvdtkveskygppcpcpa
 peflggpsvflfppkpkdtlmsirtpevtcvvdvsqdedpevqfhwvydgvgevhnaaktpreeqfnstfrvsvlt
 tvlhqdwlngkeykckvsnkglpssiektiskakgqprepqvytlppsgeemtknqvslclvkgfypsdiave
 wesngqpennnyktppvldsdgsfflysrftvdksrwqegnvfscsvmhleahhnytkqskslspgk

40

SEQ ID NO. 101

9.14.4G1 Heavy chain (gamma chain) nucleotide sequence

atggagtttgggctgagctgggtttccctgtgtgctattataaaagggtgccagtgtCAGGTGCAGCTGGTG
 GAGTCTGGGGGAGGCTTGGTCAAGCCTGGAGGGTCCCTGAGACTCTCC
 5 TGTGCAGCCTCTGGATTACCTTCACTGACTACTATATGAGCTGGATCC
 GCCAGGCTCCAGGGAAGGGACTGGAGTGGGTTTCATACATTAGTAGTA
 GTGGTAGTACCATATACTACGCAGACTCTGTGAAGGGCCGATTACCAT
 CTCCAGGGACAACGCCAAGAACTCACTGTATCTGCAAAATGAACAGCCT
 GAGAGCCGAGGACACGGCCGTGTATTACTGTGCGAGAGGCCTAACTGG
 10 GGACTACTGGGGCCAGGGAACCTGGTCACCGTCTCCTCAGCTTccccaag
 ggcccatcggtcttcccoctggcacctctcccaagagcacctctgggggcacagcggccctgggctgctgtc
 ggactactccccgaaccggtgacggtgtcgtggaactcaggcgccctgaccagcggcgtgcacacctccccggctg
 tctcacagtctcaggactctactccctcagcagcgtggtgaccgtgcccctccagcagctgggacccagacactcat
 ctgcaacgtggaatcacaagcccaagcaacaccaaggtggacaagaaagttagcccaaatctgtgacaaaactaca
 15 catgccaccgtgccagcacctgaactcctggggggaccgtcagctctctcttcccccaaaacccaaaggacacc
 ctcatgatctcccgacccctgaggtcacatgcgtgtgtgtggagctgagccacgaagacctgagggtcaagttaa
 ctgtgtacgtggagcggcgtggaggtgcataatgccaaagacaagccgcgggaggaagcagtacaacagcacgtaccg
 tgtgtgcagcgtctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgcgaaggtctccaacaaag
 cctctccagcccccacgcagaaaaccatccaaagccaaagggcagccccgagaaccacaggtgtacacctgcc
 20 cccatcccggaatgagctgaccaagaaccaggtcagcctgacctgctgtcgaaggtctctatccagcagacatcg
 ccgtggagtgaggagacgaatgggcagccggagaacaactacaagaccacgcctccgtgctgactccgacggct
 ccctctctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtctctcatgctcctgatgc
 atgaggctctgcacaaccactacacgcagaagcctctcctgctcctcggttaaatag

25 SEQ ID NO 102

9.14.4G1 Heavy chain (gamma chain) protein sequence

mefglswvflvaiikgvqcQVQLVESGGGLVKPGGSLRLSCAASGFTFSDYYMSWI
 RQAPGKGLEWVSYISSSGSTIYYADSVKGRFTISRDNAKNSLYLQMNLSRA
 EDTAVYYCARGLTGDYWGQGLTVTVSSAstkgpsvflapssktsstgtaalgclvkdyfp
 30 epvtvswnsgaltsgvhtfpavlqssglyslssvvtvpssslgtqtyicnvnhkpsntkvdkkvepksedkthtcpp
 cpapelggpsvflfppkpkdtlmsirtpvctvvdvshdepvkfnwyvdgvevhnaktkpreeqnystyrv
 vsvlthqdwlngkeyckvsnkalpapiektiskakgqprepqvyltppsrdeltknqslctivkgfypsda
 vewesngqpennyktppvldsgsfilyskltvdksrwqgnvfscsvmhhealnhytqkslspsgk

SEQ ID NO: 29

8.10.3 and 8.10.3F Heavy Chain [Gamma chain] nucleotide sequence

- 5 atggagtggggctgtcctgggtttctctgttctattttagaaggtgtccagttgGAGGTGCAGCTGGTG
GAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCC
TGTGCAGCCTCTGGATTCACTTCAGTAGTTTATGATATGACCTGGGTCC
GCCAGGCTCCAGGAAAGGGCTGGAGTGGGTTTCATACATTAGTAGTA
GAAGTAGTACCATATCTCTACGCAGACTCTGTGAAGGGCCGATTTCACCA
TCTCCAGAGACAATGCCAAGAACTCACTGTATCTGCAAATGAACAGCC
10 TGAGAGACGAGGACACGGCTGTGTATTACTGTGCGAGAGATCCTCTCTCT
AGCGGGAGCTACCTTCTTTGACTACTGGGGCCAGGGAACCCCTGGTCA
CGTCTCCTCAGCCtccaccaaggggcccatcggtcttcccoctggcgccctgctccaggagcacctccgag
agcacagcggccctgggtcgtcgtgcaaggactacttccccgaaccggfagcggfgtcgttggaaactcaggcgtct
gaccagcggcgtgcaaccttccagctgtcttacagtcctcaggaacttactccctcagcagcgtggtgacctgcc
15 ctccagcaacttcggcacccagacctacactgcaacgtatgatacaagccagcaacaccaaggtggacaagaca
gttgagcgcaaatgtgtgtcgtgagtgccaccgtgccacgacctgtgagcaggacctgactctctcttcccc
caaaacccaaggacacctcatgatctcccgaacctgaggtcacgtgcgtgtgtgtgagcgtgagccacgaaga
ccccgaggtccagttcaactgtgtacgtggacggcgtgaggtgcataatgccaaagacaaagccagggaggagca
gttcaacagcagcttccgtgtgtgtgacgtctcaccgtgtgtgacaccagactggctgaacggcgaaggagtacaagt
20 caaggtcttccaacaaggcctcccgcccccatcgagaaaacacatctccaaaccaaaggcgagcccgagaaacc
acaggtgtacacctgcccccatcccgggaggagatgaccaagaaccaggtcagcctgacctgctgtgtcaaaaggc
tttaccccaagcacatcgccgtggagtgaggagaacatgggcagccggagaacaactacaagaccacacctccca
tgtcggactccgagcgtctcttctctctacagcaagctcaccgtggacaagagcaggtggcagcagggggaacgtc
ttctatgtctcgtgatgatgagcgtctgcacaaccactacacgcagaagagcctctcctgtctccgggtaaa

25

SEQ ID NO: 30

8.10.3 and 8.10.3F Heavy Chain [Gamma chain] protein sequence

- melglwvflvallegvqeEVQLVESGGGLVQPGGSLRLSCAASGFTFSFSMTWV
RQAPGKGLEWVSYISSRSSTISYADSVKGRFTISRDNAKNSLYLQMNSLRD
30 EDTAVYYCARDPLLAGATTFDYWGQGLTVTVSSAstkqpsvflpapsrstsestaalg
clvkdyfpeptvswngaltsgvhtfpavllqssglyslssvvtvpssnftgitytcnvdhkpsntkvdktverkkc
vecppeppavpagsvflfppkpdltlmsirtpvctcvvdvshdpevqfnwydvgevhnaaktikpreeqfn
stfrvsvltvhvhdwlngkeyckvsnkglpapiektisktkgqprepqvtylpsreemtknqvslitckvkgy
psdiawewesngppennyktppmldsdgsfflyskltvdksrwqqgnvfscsvmhealhnhytkslslspgk

35

SEQ ID NO: 31

8.10.3FG1 and 8.10.3F Light Chain [Kappa chain] nucleotide sequence

- 5 atggaaacccagcgcagcttctctctcctgctactctggctccagataaccaccggaGAATTGTGTGTG
ACGCAGTCTCCAGGCACCCTGTCTTTGTCTCCAGGGGAAAAGAGCCACCC
TCTCTGCAAGGGCCAGTCAGAGTGTTAGCAGCAGTTACTTAGCCTGGTA
CCAGCAGAAACCTGGCCAGGCTCCCAGGCTCCTCATCTATGGTGCATCC
10 AGCAGGGCCCACTGGCATCCCAGACAGGTTCACTGGCAGTGGGTCTGGG
ACAGACTTCACTCTCACCATCAGCAGACTGGAGCCTGAAGATTTTGCAG
TGTATTACTGTCAGCAGTATGGTAGCTCACCTCTCACTTTCGGCGGAGG
GACCAAGGTGGAGATCAAACGAactgtggctgcaccatctgtttcatcttccgccatctgatga
gcagttgaaatctggaactgcctctgtttgtgtgcctgctgaataacttctatccagagagccaaagtacagtggaa
gtggataacgccctccaatcgggtaactccaggagagtgacacagagcaggacagcaaggacagcacciacagcc
15 tcagcagcacctgacgtgagcaaacgacactacgagaaacacaaagtctacgcctgcgaagtcacccatcaggg
cctgagctgccctgtcacaaagagcttcacaggggagagtgt

SEQ ID NO: 32

8.10.3FG1 and 8.10.3F Light Chain [Kappa chain] protein sequence

- 20 metpaqlflfllllwlpdttgEFVLTQSPGTLSPGERATLSCRASQSVSSSYLAWYQQ
KPGQAPRLLIYGAASSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAVYYCQQ
YGSSPLTFGGGKVEIKRtvaapsvfifppdsdeqlksqtasvvcllnnfyreakvqwkvdnalqsg
nsqesvtqcdskdstylsstlltllskadyekhkvyacevthqglsspvtksfnrgec

SEQ ID NO: 43

8.10.3 and 8.10.3-Ser Light Chain [Kappa chain] nucleotide sequence

- 25 atggaaacccagcgcagcttctctctcctgctactctggctccagataaccaccggaGAATTGTGTGTG
ACGCAGTCTCCAGGCACCCTGTCTTTGTCTCCAGGGGAAAAGAGCCACCC
TCTCTGCAAGGGCCAGTCAGAGTGTTAGCAGCAGTTACTTAGCCTGGTA
CCAGCAGAAAACCTGGCCAGGCTCCCAGGCTCCTCATCTATGGTGCATCC
30 AGCAGGGCCCACTGGCATCCCAGACAGGTTCACTGGCAGTGGGTCTGGG
ACAGACTTCACTCTCACCATCAGCAGACTGGAGCCTGAAGATTTTGTAG
TGTATTACTGTCAGCAGTATGGTAGCTCACCTCTCACTTTTCGGCGGAGG
GACCAAGGTGGAGATCAAACGAactgtggctgcaccatctgtttcatcttccgccatctgatga
gcagttgaaatctggaactgcctctgtttgtgtgcctgctgaataacttctatccagagagccaaagtacagtggaa
35 gtggataacgccctccaatcgggtaactccaggagagtgacacagcaggacagcagcaccctacagcc
tcagcagcacctgacgtgagcaaacgacactacgagaaacacaaagtctacgcctgcgaagtcacccatcaggg
cctgagctgccctgtcacaaagagcttcacaggggagagtgt

SEQ ID NO: 44

8.10.3 and 8.10.3-Ser Light Chain [Kappa chain] protein sequence

- 40 metpaqlflfllllwlpdttgEFVLTQSPGTLSPGERATLSCRASQSVSSSYLAWYQQ
KPGQAPRLLIYGAASSRATGIPDRFSGSGSGTDFTLTISRLEPEDFVYYCQQ
YGSSPLTFGGGKVEIKRtvaapsvfifppdsdeqlksqtasvvcllnnfyreakvqwkvdnalqsg
nsqesvtqcdskdstylsstlltllskadyekhkvyacevthqglsspvtksfnrgec

SEQ ID NO: 58

8.10.3C-Ser Heavy Chain [Gamma chain] protein sequence

- 5 melglcwvflvailegvqcEVQLVESGGGLVQPGGSLRLSCAASGFTFSFSMTWV
 RQAPGKGLEWVSYISRSSTISYADSVKGRFTISRDNNAKNSLYLQMNSLRD
 EDTAVYYCARDPLLAGATFFDYWGQGTLVTVSSAstkgpsvfplapcsrcstsestaalg
 clvkdyfpepvtvswngaltsgvhtfpavlgssgylsslsvvtvpsslgtktytcnvdhkpnsntkvdkrveskyg
 ppcppcpapeflggpsvflfppkpkdtlmsirtpetvcvvdvsqedpevqfnwyvdgvevhnaktkpreeqf
 nstyrvsvtlvhqdwlngkeyckckvsnkgplssiectiskakgqprepqvtylppsqeemtknqvsltlvkgf
 10 ypsdiavewesngqpennykttppvldsdgsfflysrftvdksrwqegnfvscsvmhcalhnhytqkslslspgk

SEQ ID NO: 60

8.10.3-CG2, 8.10.3-CG4 and 8.10.3C-Ser Light Chain [kappa chain] protein sequence

- 15 metpaqllfllllwlpdttgEIVLTQSPGTLSTLSPGERATLSCRASQSVSSSYLAWYQQ
 KPGQAPRLLIYGASSRATGIPDRFSGSGSDTFTLTISRLEPEDFAVYYCQ
 YGSSPLTFGGGKVEIKRtvaapsvflfppsdeqlksqtasvcllnnfypreakvqwkvdnalqsg
 nsqesvteqdkdstylssltltskadyekkhvyaecvthqglsspvtksfirgce

SEQ ID NO: 62

8.10.3-CG2 Heavy Chain [Gamma chain] protein sequence

- 20 melglcwvflvailegvqcEVQLVESGGGLVQPGGSLRLSCAASGFTFSFSMTWV
 RQAPGKGLEWVSYISRSSTISYADSVKGRFTISRDNNAKNSLYLQMNSLRD
 EDTAVYYCARDPLLAGATFFDYWGQGTLVTVSSAstkgpsvfplapcsrcstsestaalg
 clvkdyfpepvtvswngaltsgvhtfpavlgssgylsslsvvtvpssnfgtqytcnvdhkpnsntkvdktrkcc
 25 vecppcpappvagsvflfppkpkdtlmsirtpetvcvvdvshedpevqfnwyvdgvevhnaktkpreeqfn
 stfrvsvtlvhqdwlngkeyckckvsnkgplapiectisktkgqprepqvtylppsrreemtknqvsltlvkgfy
 psdiavewesngqpennykttppmldsdgsfflyskltvdksrwqqgnfvscsvmhcalhnhytqkslslspgk

SEQ ID NO: 90

8.10.3-Ser Heavy Chain [Gamma chain] protein sequence

- 30 melglcwvflvailegvqcEVQLVESGGGLVQPGGSLRLSCAASGFTFSFSMTWV
 RQAPGKGLEWVSYISRSSTISYADSVKGRFTISRDNNAKNSLYLQMNSLRD
 EDTAVYYCARDPLLAGATFFDYWGQGTLVTVSSAstkgpsvfplapcsrcstsestaalg
 clvkdyfpepvtvswngaltsgvhtfpavlgssgylsslsvvtvpsslgtktytcnvdhkpnsntkvdkrveskyg
 35 ppcppcpapeflggpsvflfppkpkdtlmsirtpetvcvvdvsqedpevqfnwyvdgvevhnaktkpreeqf
 nstyrvsvtlvhqdwlngkeyckckvsnkgplssiectiskakgqprepqvtylppsqeemtknqvsltlvkgf
 ypsdiavewesngqpennykttppvldsdgsfflysrftvdksrwqegnfvscsvmhcalhnhytqkslslspgk

SEQ ID NO: 94

8.10.3-CG4 Heavy Chain [Gamma chain] protein sequence

- melglcwvflvailegvqcEVQLVESGGGLVQPGGSLRLSCAASGFTFSFSMTWV
 5 RQAPGKGLEWVSYISSRSSTISYADSVKGRFTISRDNAKNSLYLQMNSLRD
 EDTAVYYCARD**P****L****L****A****G****A****T****F****F****D****Y**WGQGLTVTVSSAstkgpsvflpacsrstsestaalg
 clvkdyfpepvtvswngaltsgvhtfpavlqssglyslssvvtvpssslgtktytencvdkpsntkvdkrveskyg
 ppcpscpapeflggpsvflfppkpkdlmistrpevtcvvdvsqedpevqfnwyvdgvevhnaktkpreeqf
 10 nstyrvsvltvlhqdwlngkeykckvsnkglpssiektiskakgqprepqvyltppsgeemtknqvsltlvkglf
 ypsdiafewesngqpennyktppvldsdsfflyslrvtldksrwqqgnvfscsvmhleahlnhytkslslspgk

SEQ ID NO: 97

8.10.3FG1 Heavy Chain nucleotide sequence

- atggagttggggctgagctgggttttccctgtgtctattataaaagggtgccagtgGAGGTGCAGCTGGTG
 15 GAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCC
 TGTGTCAGCTCTGGATTACCTTCAGTAGTTTATAGTATGACTGGGTCC
 GCCAGGCTCCAGGAAAGGGGCTGGAGTGGGTTCATACATTAGTAGTA
 GAAGTAGTACCATATCCTACGCAGACTCTGTGAAGGGCCGATTACCA
 20 TCTCCAGAGACAATGCCAAGAACTCACTGTATCTGCAAAATGAACAGCC
 TGAGAGACGAGGACACGGCTGTGTATTACTGTGCGAGAGATCCTCTCT
 AGCGGGAAGCTACCTCTTTGACTACTGGGGCCAGGGAACCCTGGTCA
 CGTCTCCTCAGCCtccaccaaggcccacgtcttccctggcaccctctcccaagagcaccctctggg
 ggcacacggccctgggctgctgtgtcaaggactacttcccgaaccggtagcggtgtctggaactcaggccccc
 tgaccagcggcgtgcacaccttccggctgtcctacagtctcaggactctactcctcagcagcggtgtgaccgtgc
 25 cctccagcagcttgggcaaccagacactacatctgcaacgtgaatcacaagcccagcaacaccaagggtggacaagaa
 agttgagcccaaatctgtgacaaaactcacacatgccaccgtgccagcactgaactcctggggggaccgtcagt
 ctctctctcccccaaaacccaaggacacctcatgaatcctccggaccctgaggtgcacatgctgtgtgggacgtg
 agccacgaagacctgaggtcaagttcaactgtgtacgtggacggcggtgaggtgcataatgccaagacaagccgc
 gggagggagcagtaacaacagcagctaccgtgtgtgcagcgtctcaccgtctcaccaggactggcgtgaatggcaa
 30 ggaagtacaagtgcaaggtctccaacaagccctccagcccccactgagaaaacctctccaagaccaaggggcag
 cccggagaacacagcaggtgtacacctgccccatccgggaatgagctgaccaagaaccaaggtcagcgtgaactgoc
 tgtgtcaagggctctatccacagacatcgccgtggagtgaggagacaaatgggcagccggagaacaactacaagac
 caagcctccctgtgtgacacccgacggctctctctctacagcaagctcaacgttgacaaagcagcaggtggcagca
 ggggaacgtctctcatgctcctgtgatgcatgaggtctgcacaaccactacacgcgaagaagcctctcctctcctcc
 35 ggtaaatag

SEQ ID NO: 98

8.10.3FG1 Heavy chain (gamma chain) protein sequence

- melglcwvflvailegvqcEVQLVESGGGLVQPGGSLRLSCAASGFTFSFSMTWV
 40 RQAPGKGLEWVSYISSRSSTISYADSVKGRFTISRDNAKNSLYLQMNSLRD
 EDTAVYYCARD**P****L****L****A****G****A****T****F****F****D****Y**WGQGLTVTVSSAstkgpsvflpsskstsggtaal
 gclvkdyfpepvtvswngaltsgvhtfpavlqssglyslssvvtvpssslgtqtyicnvhnkpsntkvdldkvepk
 sdkthtppcpapeflggpsvflfppkpkdlmistrpevtcvvdvshdepvkfnwyvdgvevhnaktkpr
 45 eqynstyrvsvltvlhqdwlngkeykckvsnkalpapiektiskakgqprepqvyltppsrdeltknqvsltlv
 kgfypsdiawesngqpennyktppvldsdsfflyslrvtldksrwqqgnvfscsvmhleahlnhytkslsls
 pgk

SEQ ID NO: 33

9.7.2IF Heavy Chain [Gamma chain] nucleotide sequence

- atggagttgggctgagctgggtttcttctgttctgattataaaaggctgcagtgtcAGGTGCAGCTGGTG
GAGTCTGGGGGAGGCTTGGTCAAGCCTGGAGGGTCCCTGAGACTCTCC
5 TGTGCAGCCTCTGGATTACCTTCAGTGACTACTACATGAGCTGGATCC
GCCAGGCTCCAGGGAAGGGGCTGGAGTGGGTTCATACATTAGTAGTA
TGTGGTAGTACCATATACTACGCAGACTCTGTGAAGGGCCGATTACCACAT
CTCCAGGGAACAACGCCAAGAATTCACTGTATCTGCAAAATGAACAGCCT
GAGAGCCGAGGACACGCCGTGTATTACTGTGCGAGGCGTATAGGAGG
10 TATGACAGTCTTGGGGCCAAGGGACCACGGTACACGTCTCCTCAGCTTcca
ccaagggccaccctgcttccccctggcgcctgctctagaagcacctccgagagcacagcggccctggcctg
gtcaaggactactccccgaacccgtgacgggtgctgctggaactcaggcgcctgaccacggcgtgcacacctccc
agctgtctacagtctcaggaactctactccctcagcagcgtggtgacccgtccctccagcaactcggcgaaccaaac
ctacacctgcaactgtagalcaacaagccagcaaaccaagggtggacaagacagtgtagcgcaaatgtgtgtcagat
15 gcccacctgtgccagcaccacctgtggcaggaccgtcagcttctcttcccccaaaaacgaagacacctcatga
tctcccgaccctcagggtcagctgctgggtgtgtgacgtgagccacgaagaccccgaggtccagtccaactggtac
gtggaaggcgtggagggtgcataatgccaaagacaaagccacgggagcaggttcaacagcagcttccgtgtgtgta
cgctctcaacgtgtgtgcaaccaggactggtgacggcnaaggagtacaagtgcaggtctccaacaaagccctccca
gccccatcgagaaaacatctccaaacaaaggcagccccgagaaccacagggtgtacacctgcccccatccc
20 gggaaggagatgacaaagaccaggtcagcctgacctgcctggtcaaaagcttctaccacgcagcatcggcgtgga
gtgggagagcaatgggcagccggagaaacatcaagaccacacctccatgctgactccagcggctctcttcc
tctacagaagctccacgtgggacaagagcaggtggcagcaggggaacgttctctcatgctccgtgatgatgaggt
ctgcacaacctacacgcagaagagcctccctgctcctcggtaaa

25 SEQ ID NO: 34

9.7.2IF Heavy Chain [Gamma Chain] protein sequence

- mefglswvflvaiikqvqcQVQLVESGGGLVKPGGSLRLSCAASGFTFSDYVMSWI
RQAPGKGLEWVSYISSGSTIYYADSVKGRFTISRDNAKNSLYLQMNSLRA
EDTAVVYCARRIGGMDVWGQGTITVSSAstkgpsvflapcsrcstestaaclgclvdyf
30 pepvtvswnsalstgsvhtfpavlqssgylslssvvtvpssnfgtqytctnvdkhpsntkvdktverkcovecppe
pappvagpsvflfpkpkltdlmisrtpetvctvvvdshedpevqfnwvyvdgvevhnaktcpreeqfnstfvrvs
vlvvhqdwlngekeyckvsnklpapielktisktkgpqrepqvyltppsreemtknqvslclvklfgyfypsdiav
ewesngqpennykttppmldsdgsfilyskltvdksrwqqgnvfscvmhhealnhnytkslslspgk

35 SEQ ID NO: 35

9.7.2IF Light Chain [Kappa chain] nucleotide sequence

- atggacatgaggtcccgctcagctctctggggctcctgctactctgctccgaggtgccagatgGACATCC
AGATGACCCAGTCTCCATCCTCCCTGTCTGTCATCTGTAGGAGACAGAGT
CACCATCACTTGC CGGGCAAGTCAGAGCATTAGCGGCTTTTTTAATTTGG
40 TATCAGCAGAGACCAGGGAAGCCCTAAGCTCCTGATCTATGCTACA
TCCAGTTTACAAGTGGGGTCCCATCAAGGTTTCAGTGGCAGTGGATCTG
GGACAGATTTCACCTCTCACCATCAGCAGCTGCAACCTGAAGATTTTGC
AACTTACTACTGTCAACAGAGTTACAGTACCCCATTCACITTCGGCCCT
GGGACCAAAGTGGATATCAAACGAactgtgctgcaacctgtcttcaictctccgcacatgta
45 tgaagcagttgaaalcgtgaactgctctgtgtgtgctgctgaataactctatccagagagcccaaggtacagtga
aggtggataacgccctccaaalccgggttaactccagagaggtgtcacagacagagacageaaagacagacactaca
gcctcagcagcaacctgacgtgagcaagcagactacgagaacacaaagtctacgctgcgaagteaccatca
ggcctgagctcggcctcacaagagcttcaacaggggagagtgt

SEQ ID NO: 36

9.7.2IF Light Chain [Kappa chain] protein sequence

- 5 mdmrnpaqlglgllllwrgarcDIQMTQSPSSLSASVGDRTVITTCRASQISGFLIYWYQ
QRPKGAPKLLIYATSSLSQSGVPSRFSGSGSDFTLTISSLQPEDFATYYCQ
QSYSTPTFTFGPGTKVDIKRtvaapsvfifppdsdeqlksgtasvvcllnnfyreakvqkwkdnaqls
gnsqesvteqdkstdstyslssltlslkadyekhkvyacevthqglsspvtksfirgce

SEQ ID NO: 45

9.7.2 Heavy Chain [Gamma chain] nucleotide sequence

- 10 atggagatttggcgtagactgggttttccctgttgcattataaaagctgacagtcAGGTGCAGCTGGTG
GAGTCTGGGGAGGCTTGGTCAAGCCTGGAGGGTCCCTGAGACTCTCC
TGTGCAGCCTCTGGATTACCTTCAGTGACTACTACATGAGCTGGATCC
GCCAGCTCCAGGGAAGGGGCTGGAGTGGGTTTCATACATTAGTAGTA
15 GTGGTAGTACCATATACTACGCAGACTCTGTGAAGGGCCGATTACCAT
CTCAGGGACAACGCCAAGAATTCACTGTATCTGCAAAATGAACAGCCT
GAGAGCCGAGGACACGGCCGTGTATTACTGTGCGAGGGCTATAGGAGG
TATGGACGCTCTGGGGCCAAGGGACCACGGTCACCGTCTCCTCAGCTTcca
20 ccaagggccatccgtcttccctggcgcctgctctagaagcacctccgagagcacagcgccctggcgctcg
gtcaaggactacttcccgaaccgggtgacgggtgtcgtgaactcaggcgctctgaccagcgcggtgcacacttccc
agctgtctacagctcctcaggactctactcctcagcagcgtggtgacctgacctccagcagctgggcacgaagac
ctacacctgcaacgtatgatacacaagccagcaacaccaaggtggacaaagagaggtgagtcctcaaatatggtccccc
25 gccccatcatgccagcacctgagttcctggggggaccatcatgcttctctgttcccccaaacccaaggacactctcat
gatctccggaccctgaggtcacgtgcgtgggtgggtgacgtgacccaggaagaccccagggtccagtccaactggt
acgtgtaggctggtgaggtgataatgccaaagacaagccgcgggaggaagcagttcaacagcacgtacctgtggtg
cagcgtctcaccgtctgcaccaggactggtgacggcagggagtaacaggtgcaaggtctccaacaagggcctc
30 ccgtctccatcgagaaaacatctccaaagccaaaggcagccccgagagccacaggtgtacacctgcccccat
cccaggagagatgaccaagaaccaggtcagcctgacctgctggtcaaaagcttctaccccagacatcgccgt
ggagtgaggagaacaatgggcagccggagaaacaactacaagaccacgcctccgtgctggaactccgagcgctcctt
ttctctacagcaggttaaccgtggacaagagcaggtggcaggagggaatgctctctcatgctccgtgatgcatgag
gctctgcacaaccactacacagaaagacgctctcctgtctccgggtaaa

SEQ ID NO: 46

9.7.2 Heavy Chain [Gamma Chain] protein sequence

- 35 mefglsvvflvailkvqeqVQLVESGGGLVPGGSLRLSCAASGFTFSDDYYMSWI
RQAPGKGLEWVSYISSSGSTIYYADSVKGRFTISRDNKNSLYLQMNSLRA
EDTAVYYCARRIGGMDVWQGTTVTVSS Astkpsvfplapcsrstsestaalgc/vkdyf
 pepvtvswnsaltsghvhtfavlqssglyslsvvtvpssslgtktyctcnvdhkpantkvdkrveskygppcpse
 papelfggpsvfllppkpkdltlmsirtpentcvvvdvsqedpevqfnwvydgvvhnaktkpreeqfnstyrvv
40 svltvlthdwlngkeykckvsnklgppsielktiskakgqprepvytlppsqeemtknqvslctlvkgfypsdia
vewesngqpenntktpvldsdgfflysrllvtvdkswrqevgnvfscsvmhcalhnhytqksllspgk

SEO ID NO: 47

9.7.2 and 9.7.2-Ser Light Chain [Kappa chain] nucleotide sequence

5 atggacatgagggtccccgctcagctctggggctcctgctactctgctccgaggtgccagatg(GACATCC
AGATGACCCAGCTTCCATCTCCCTGTCTGCATCTGTGAGGACAGAGT
CACCATCACTTGGCCGGCAAGTCAAGCATTAGCGGCTTTTAAATTGG
TATCAGCAGAGACCAAGGAAAGCCCTAAGCTCTGTATCTATGCTACA
TCCAGTTTACAAAGTGGGGTCCCCATTAAAGGTTCACTGGCAGTGAATCTG
GGACAGATTTCACTCTACCATCAGCAGTCTGCAACCTGAAGATTTTGC
10 AACTTACTACTGTCAACAGAGTTACAGTACCCCAATTCACTTTCGCCGCTT
GGGACCAAAAGTGGATATCAACAGCAactggtgctgacacctgtctctatctccgcacatga
tgacagttgaaactggaactgctctgtgtgctgctgtaataacttctatccagagagccaaagtgaacgga
agggtgataacgccctccaactgggtaactccagagagagtgtgcaagagcaggacagcaaggacagcacctaca
gcttcagcagcactgacgctgacgaagcagcagcgaagaacaaaggtacgctgctgaggtcacccatca
15 ggctgagcgtcgcctctcaaaaggctctcaacaggctgacgaggtgt

SEO ID NO: 48

9.7.2 and 9.7.2-Ser Light Chain [Kappa chain] protein sequence

20 mdmrpvagllglglwlgarcDIQMTQSPSSLSASVGDRTITCRASQISIGFLIWYQ
QRP GKAPKLLIYATSSLSQVPLRFSGESGDFTLTISSLQPEDFATYYCQ
QSYSTPTFTFGDKVIDIKRtvaapsvfiippsdeqlksgtadvcllnnfypreakvqwkvdnlqsg
ngsqevsteqdgkdstylsstllskadyekhlkvaccvthqglsspvtkcfnr

SEO ID NO: 50

25 9.7.2C-Ser Heavy Chain [Gamma chain] protein sequence

30 mefgslswvflvliijkvgacQVQLVSGGGGLVKKPGKSLRLSCAASGFTFSDYFMSWI
RQAPGKGLEWVSYSXSSGSGTITYADSVKGRFTISRDNAKNSLYLQMNSRL
EDTAVPYGCAIRIGGMDYWGQGTFTVTVSSAstkgpsvfplapcserstsestaalgevkdyfp
epvtvswnsгалтсгvhtplamrlsgyslsstvpssslgltkytenvdhkpntkvdcrvesygpppcp
apeflgpgsvflfpplkplqvstentpvtcvvdsgepeqvfuwiydvgnvhnaktpeeqnfrsttyrvs
vltvlhqdwngkeylckkcvnslgppssiektikagpprepqvtltppsgeemtkqvsitclvfgkfyssda
ewscagennvktptpvlslsdsglyfslrvltdksrwgeqncfsvcmhealhhnhtyqtlslslspgk

SEO ID NO: 52

35 9.7.2C-Ser, 9.7.2-CG2 and 9.7.2-CG4 Light Chain [Kappa chain] protein sequence

mdmrvpaqlgllllwlrgrcdIQMTQSPSSLSASVGDRVTITCRASQSSIGFLIWIYQ
QKPGKAPKLLIYATSSLSQGVPSRFSGSGSGTDFTLITSSLPEDFATYYCQ
QSYSTPFTFGPGTKVDIKRtvaapsvfipfsspeqlgkstgsvcllnnfpreakvqwkvdnals
gnsqspvteqdskslystslstlksadyekkhvycvethglsspvtksfnrgec

SEQ ID NO: 66

9.7.2-CG2 Heavy Chain [Gamma chain] protein sequence

- 5 mefglswvflvaiikgvqcQVQLVESGGGLVKPGGSLRLSCAASGFTFSDYYMSWI
 RQAPGKGLEWVSYISSSGSTIYYADSVKGRFTISRDNAKNSLYLQMNSLRA
 EDTAVYYCAIRIGGMDYWGQGTTTVTVSSAstkgpsvfplapcsrcstsestaalgclvkdyfp
 epvtvswngaltsgvhtfpavqlqssglyslssvvtvpssnfgtktytcnvdhkpsntkvdtkverkcceppcp
 appvagspvsflfppkpkdtlmsrtpevtcvvvdvshdpevqfinwyvdgvevhnaktkpreeqfnstfrvsv
 10 ltvvhqdwlngkeyckckvsnkglpapiektisktkgqprepqvyltppsreemtknqvslclvkgfypsdiave
 wesngqpennyktppmldsdgsfflyskltvdksrwqegnvfscsvmhealhnhytqklsislspgk

SEQ ID NO: 70

9.7.2-CG4 Heavy Chain [Gamma chain] protein sequence

- 15 mefglswvflvaiikgvqcQVQLVESGGGLVKPGGSLRLSCAASGFTFSDYYMSWI
 RQAPGKGLEWVSYISSSGSTIYYADSVKGRFTISRDNAKNSLYLQMNSLRA
 EDTAVYYCARIGGMDYWGQGTTTVTVSSAstkgpsvfplapcsrcstsestaalgclvkdyfp
 epvtvswngaltsgvhtfpavqlqssglyslssvvtvpssslgktytcnvdhkpsntkvdtkverksygppepcsp
 apelfggpsvflfppkpkdtlmsrtpevtcvvvdvsqedpevqfinwyvdgvevhnaktkpreeqfnstfrvsv
 20 vitvlhqdwlngkeyckckvsnkglpssiektiskakgqprepqvyltppsqeemtknqvslclvkgfypsdiav
 ewesngqpennyktppvldsdgsfflysrlltvdksrwqegnvfscsvmhealhnhytqklsislspgk

SEQ ID NO: 86

9.7.2-Ser Heavy Chain [Gamma chain] protein sequence

- 25 mefglswvflvaiikgvqcQVQLVESGGGLVKPGGSLRLSCAASGFTFSDYYMSWI
 RQAPGKGLEWVSYISSSGSTIYYADSVKGRFTISRDNAKNSLYLQMNSLRA
 EDTAVYYCARRIGGMDYWGQGTTTVTVSSAstkgpsvflapcsrcstsestaalgclvkdyf
 pepvtvswngaltsgvhtfpavqlqssglyslssvvtvpssslgktytcnvdhkpsntkvdtkverksygppeppc
 papelfggpsvflfppkpkdtlmsrtpevtcvvvdvsqedpevqfinwyvdgvevhnaktkpreeqfnstfrvsv
 30 svltvlhqdwlngkeyckckvsnkglpssiektiskakgqprepqvyltppsqeemtknqvslclvkgfypsdiav
 vewesngqpennyktppvldsdgsfflysrlltvdksrwqegnvfscsvmhealhnhytqklsislspgk

What is claimed is:

1. A human monoclonal antibody or an antigen-binding portion thereof that specifically binds to M-CSF.
2. The human monoclonal antibody or antigen-binding portion according to claim 1, wherein said antibody or portion possesses at least one of the following properties:
 - a) binds to human secreted isoforms of M-CSF and membrane bound isoforms of M-CSF;
 - b) has a selectivity for M-CSF that is at least 100 times greater than its selectivity for GM-CSF or G-CSF;
 - c) binds to M-CSF with a K_D of 1.0×10^{-7} M or less;
 - d) has an off rate (k_{off}) for M-CSF of $2.0 \times 10^{-4} s^{-1}$ or smaller; or
 - e) binds human M-CSF in the presence of human *c-fms*.
3. The human monoclonal antibody or portion according to claim 2, wherein said antibody or fragment blocks binding to *c-fms* and binds M-CSF with a K_D of 1.0×10^{-7} M or less.
4. A human monoclonal antibody 8.10.3F or an antigen-binding portion thereof that specifically binds to M-CSF.
5. A human monoclonal antibody 9.14.4I or an antigen-binding portion thereof that specifically binds to M-CSF.
6. A humanized, chimeric or human monoclonal antibody or antigen-binding portion thereof that binds specifically to and inhibits human M-CSF, wherein the antibody or portion thereof has at least one property selected from the group consisting of:
 - a) cross-competes for binding to M-CSF with an antibody selected from the group consisting of: antibody 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-

Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 and 9.14.4G1;

b) competes for binding to M-CSF with an antibody selected from the group consisting of: 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 and 9.14.4G1;

c) binds to the same epitope of M-CSF as an antibody selected from the group consisting of: antibody 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 and 9.14.4G1;

d) binds to M-CSF with substantially the same K_D as an antibody selected from the group consisting of: antibody 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 and 9.14.4G1; and

e) binds to M-CSF with substantially the same off rate as an antibody selected from the group consisting of: antibody 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 and 9.14.4G1.

7. A monoclonal antibody that specifically binds M-CSF, wherein the antibody is selected from the group consisting of:

a) an antibody comprising the heavy chain amino acid sequence set forth in SEQ ID NO: 2 and the light chain amino acid sequence set forth in SEQ ID NO: 4, without the signal sequences;

b) an antibody comprising the heavy chain amino acid sequence set forth in SEQ ID NO: 6 and the light chain amino acid sequence set forth in SEQ ID NO: 8, without the signal sequences;

- c) an antibody comprising the heavy chain amino acid sequence set forth in SEQ ID NO: 10 and the light chain amino acid sequence set forth in SEQ ID NO: 12, without the signal sequences;
- d) an antibody comprising the heavy chain amino acid sequence set forth in SEQ ID NO: 14 and the light chain amino acid sequence set forth in SEQ ID NO: 16, without the signal sequences;
- e) an antibody comprising the heavy chain amino acid sequence set forth in SEQ ID NO: 18 and the light chain amino acid sequence set forth in SEQ ID NO: 20, without the signal sequences;
- f) an antibody comprising the heavy chain amino acid sequence set forth in SEQ ID NO: 22 and the light chain amino acid sequence set forth in SEQ ID NO: 24, without the signal sequences;
- g) an antibody comprising the heavy chain amino acid sequence set forth in SEQ ID NO: 26 and the light chain amino acid sequence set forth in SEQ ID NO: 28, without the signal sequences;
- h) an antibody comprising the heavy chain amino acid sequence set forth in SEQ ID NO: 38 and the light chain amino acid sequence set forth in SEQ ID NO: 28, without the signal sequences;
- i) an antibody comprising the heavy chain amino acid sequence set forth in SEQ ID NO: 54 and the light chain amino acid sequence set forth in SEQ ID NO: 56, without the signal sequences;
- j) an antibody comprising the heavy chain amino acid sequence set forth in SEQ ID NO: 74 and the light chain amino acid sequence set forth in SEQ ID NO: 56, without the signal sequences;
- k) an antibody comprising the heavy chain amino acid sequence set forth in SEQ ID NO: 78 and the light chain amino acid sequence set forth in SEQ ID NO: 56, without the signal sequences;
- l) an antibody comprising the heavy chain amino acid sequence set forth in SEQ ID NO: 82 and the light chain amino acid sequence set forth in SEQ ID NO: 28, without the signal sequences;

m) an antibody comprising the heavy chain amino acid sequence set forth in SEQ ID NO: 102 and the light chain amino acid sequence set forth in SEQ ID NO: 28, without the signal sequences;

n) an antibody comprising the heavy chain amino acid sequence set forth in SEQ ID NO: 30 and the light chain amino acid sequence set forth in SEQ ID NO: 32, without the signal sequences;

o) an antibody comprising the heavy chain amino acid sequence set forth in SEQ ID NO: 30 and the light chain amino acid sequence set forth in SEQ ID NO: 44, without the signal sequences;

p) an antibody comprising the heavy chain amino acid sequence set forth in SEQ ID NO: 58 and the light chain amino acid sequence set forth in SEQ ID NO: 60, without the signal sequences;

q) an antibody comprising the heavy chain amino acid sequence set forth in SEQ ID NO: 62 and the light chain amino acid sequence set forth in SEQ ID NO: 60, without the signal sequences;

r) an antibody comprising the heavy chain amino acid sequence set forth in SEQ ID NO: 90 and the light chain amino acid sequence set forth in SEQ ID NO: 44, without the signal sequences;

s) an antibody comprising the heavy chain amino acid sequence set forth in SEQ ID NO: 94 and the light chain amino acid sequence set forth in SEQ ID NO: 60, without the signal sequences;

t) an antibody comprising the heavy chain amino acid sequence set forth in SEQ ID NO: 98 and the light chain amino acid sequence set forth in SEQ ID NO: 32, without the signal sequences;

u) an antibody comprising the heavy chain amino acid sequence set forth in SEQ ID NO: 34 and the light chain amino acid sequence set forth in SEQ ID NO: 36, without the signal sequences;

v) an antibody comprising the heavy chain amino acid sequence set forth in SEQ ID NO: 46 and the light chain amino acid sequence set forth in SEQ ID NO: 48, without the signal sequences;

w) an antibody comprising the heavy chain amino acid sequence set forth in SEQ ID NO: 50 and the light chain amino acid sequence set forth in SEQ ID NO: 52, without the signal sequences;

x) an antibody comprising the heavy chain amino acid sequence set forth in SEQ ID NO: 66 and the light chain amino acid sequence set forth in SEQ ID NO: 52, without the signal sequences;

y) an antibody comprising the heavy chain amino acid sequence set forth in SEQ ID NO: 70 and the light chain amino acid sequence set forth in SEQ ID NO: 52, without the signal sequences; and

z) an antibody comprising the heavy chain amino acid sequence set forth in SEQ ID NO: 86 and the light chain amino acid sequence set forth in SEQ ID NO: 48, without the signal sequences.

8. The human monoclonal antibody or an antigen-binding portion thereof according to claim 1, wherein said antibody or antigen-binding portion comprises:

a) a heavy chain CDR1, CDR2 and CDR3 independently selected from the heavy chain of an antibody selected from the group consisting of: monoclonal antibody 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 and 9.14.4G1; or

b) a light chain CDR1, CDR2 and CDR3 independently selected from the light chain of an antibody selected from the group consisting of: monoclonal antibody 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 and 9.14.4G1.

9. A monoclonal antibody or an antigen-binding portion thereof that specifically binds M-CSF, wherein:

a) the heavy chain comprises the heavy chain CDR1, CDR2 and CDR3 of an antibody selected from the group consisting of: antibody 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 and 9.14.4G1;

b) the light chain comprises the heavy chain CDR1, CDR2 and CDR3 of an antibody selected from the group consisting of: antibody 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 and 9.14.4G1;

c) the antibody comprises a heavy chain of (a) and a light chain of (b); or

d) the antibody of (c) wherein the heavy chain and light chain CDR amino acid sequences are selected from the same antibody.

10. A monoclonal antibody or an antigen-binding portion thereof that specifically binds M-CSF, wherein the antibody comprises:

a) a heavy chain comprising the amino acid sequence from the beginning of the CDR1 through the end of the CDR3 of the heavy chain of an antibody selected from the group consisting of: antibody 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 and 9.14.4G1;

b) a light chain comprising the amino acid sequence from the beginning of the CDR1 through the end of the CDR3 of an antibody selected from the group consisting of: antibody 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 and 9.14.4G1;

c) the heavy chain of (a) and the light chain of (b); or

d) the heavy chain of (a) and the light chain of (b) amino acid sequences are selected from the same antibody.

11. The monoclonal antibody or antigen-binding portion according to claim 10 wherein said monoclonal antibody or portion comprises:

a) the heavy chain variable domain (VH) amino acid sequence, without a signal sequence, of an antibody selected from the group consisting of: antibody 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 and 9.14.4G1;

b) the light chain variable domain (VL) amino acid sequence, without a signal sequence, of an antibody selected from the group consisting of: antibody 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 and 9.14.4G1;

c) the VH amino acid sequence of (a) and the VL amino acid sequence of (b); or

d) the VH amino acid sequence of (a) and the VL amino acid sequence of (b), wherein the VH and VL are from the same antibody.

12. A polypeptide selected from the group consisting of:

a) a polypeptide comprising the amino acid sequence of SEQ ID NO: 2, without the signal sequence;

b) a polypeptide comprising the amino acid sequence of SEQ ID NO: 6, without the signal sequence;

c) a polypeptide comprising the amino acid sequence of SEQ ID NO: 10, without the signal sequence;

d) a polypeptide comprising the amino acid sequence of SEQ ID NO: 14, without the signal sequence;

e) a polypeptide comprising the amino acid sequence of SEQ ID NO: 18, without the signal sequence;

- f) a polypeptide comprising the amino acid sequence of SEQ ID NO: 22, without the signal sequence;
- g) a polypeptide comprising the amino acid sequence of SEQ ID NO: 26, without the signal sequence;
- h) a polypeptide comprising the amino acid sequence of SEQ ID NO: 30, without the signal sequence;
- i) a polypeptide comprising the amino acid sequence of SEQ ID NO: 34, without the signal sequence;
- j) a polypeptide comprising the amino acid sequence of SEQ ID NO: 38, without the signal sequence;
- k) a polypeptide comprising the amino acid sequence of SEQ ID NO: 46, without the signal sequence;
- l) a polypeptide comprising the amino acid sequence of SEQ ID NO: 50, without the signal sequence;
- m) a polypeptide comprising the amino acid sequence of SEQ ID NO: 54, without the signal sequence;
- n) a polypeptide comprising the amino acid sequence of SEQ ID NO: 58, without the signal sequence;
- o) a polypeptide comprising the amino acid sequence of SEQ ID NO: 62, without the signal sequence;
- p) a polypeptide comprising the amino acid sequence of SEQ ID NO: 66, without the signal sequence;
- q) a polypeptide comprising the amino acid sequence of SEQ ID NO: 70, without the signal sequence;
- r) a polypeptide comprising the amino acid sequence of SEQ ID NO: 74, without the signal sequence;
- s) a polypeptide comprising the amino acid sequence of SEQ ID NO: 78, without the signal sequence;
- t) a polypeptide comprising the amino acid sequence of SEQ ID NO: 82, without the signal sequence;
- u) a polypeptide comprising the amino acid sequence of SEQ ID NO: 86, without the signal sequence;

- v) a polypeptide comprising the amino acid sequence of SEQ ID NO: 90, without the signal sequence;
- w) a polypeptide comprising the amino acid sequence of SEQ ID NO: 94, without the signal sequence;
- x) a polypeptide comprising the amino acid sequence of SEQ ID NO: 98, without the signal sequence; and
- y) a polypeptide comprising the amino acid sequence of SEQ ID NO: 102, without the signal sequence.

13. The polypeptide according to claim 12 comprising the amino acid sequence of SEQ ID NO: 30, without the signal sequence.

14. The polypeptide according to claim 12 comprising the amino acid sequence of SEQ ID NO: 26, without the signal sequence.

15. A polypeptide selected from the group consisting of:

- a) a polypeptide comprising the amino acid sequence of SEQ ID NO: 4, without the signal sequence;
- b) a polypeptide comprising the amino acid sequence of SEQ ID NO: 8, without the signal sequence;
- c) a polypeptide comprising the amino acid sequence of SEQ ID NO: 12, without the signal sequence;
- d) a polypeptide comprising the amino acid sequence of SEQ ID NO: 16, without the signal sequence;
- e) a polypeptide comprising the amino acid sequence of SEQ ID NO: 20, without the signal sequence;
- f) a polypeptide comprising the amino acid sequence of SEQ ID NO: 24, without the signal sequence;
- g) a polypeptide comprising the amino acid sequence of SEQ ID NO: 28, without the signal sequence;
- h) a polypeptide comprising the amino acid sequence of SEQ ID NO: 32, without the signal sequence;
- i) a polypeptide comprising the amino acid sequence of SEQ ID NO: 36, without the signal sequence;

- j) a polypeptide comprising the amino acid sequence of SEQ ID NO: 44, without the signal sequence;
- k) a polypeptide comprising the amino acid sequence of SEQ ID NO: 48, without the signal sequence;
- l) a polypeptide comprising the amino acid sequence of SEQ ID NO: 52, without the signal sequence;
- m) a polypeptide comprising the amino acid sequence of SEQ ID NO: 56, without the signal sequence; and
- n) a polypeptide comprising the amino acid sequence of SEQ ID NO: 60, without the signal sequence.

16. The polypeptide according to claim 15 comprising the amino acid sequence of SEQ ID NO: 32, without the signal sequence.

17. The polypeptide according to claim 15 comprising the amino acid sequence of SEQ ID NO: 28, without the signal sequence.

18. A monoclonal antibody or an antigen-binding portion thereof that specifically binds M-CSF, wherein said antibody or antigen-binding portion comprises one or more of an FR1, FR2, FR3 or F4' amino acid sequence of an antibody selected from the group consisting of: antibody 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 and 9.14.4G1.

19. The human monoclonal antibody according to claim 1, wherein the antibody comprises:

- a) a heavy chain amino acid sequence that is at least 90% identical to the heavy chain amino acid sequence of monoclonal antibody 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 or 9.14.4G1, without the signal sequence;

- b) a light chain amino acid sequence that is at least 90% identical to the light chain amino acid sequence of monoclonal antibody 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 or 9.14.4G1, without the signal sequence; or
- c) the heavy chain amino acid sequence of (a) and the light chain amino acid sequence of (b).

20. The monoclonal antibody or antigen-binding portion according to any one of claims 1-19, wherein the C-terminal lysine of the heavy chain of said antibody or antigen-binding portion is not present.

21. A pharmaceutical composition, comprising the antibody or antigen-binding portion according to any one of claims 1-20 and a pharmaceutically acceptable carrier.

22. The pharmaceutical composition of claim 21 wherein the antibody is monoclonal anti-M-CSF antibody 9.14.4I or 8.10.3F.

23. A method for treating a condition selected from the group consisting of arthritis, psoriatic arthritis, ankylosing spondylitis, Reiter's syndrome, rheumatoid arthritis, gout, traumatic arthritis, rubella arthritis and acute synovitis and other arthritic conditions, sepsis, septic shock, endotoxic shock, gram negative sepsis, toxic shock syndrome, Alzheimer's disease, stroke, neurotrauma, asthma, adult respiratory distress syndrome, cerebral malaria, chronic pulmonary inflammatory disease, silicosis, pulmonary sarcoidosis, bone resorption disease, osteoporosis, restenosis, cardiac and renal reperfusion injury, thrombosis, glomerulonephritis, diabetes, graft vs. host reaction, allograft rejection, inflammatory bowel disease, Crohn's disease, ulcerative colitis, multiple sclerosis, muscle degeneration, eczema, contact dermatitis, psoriasis, sunburn and conjunctivitis shock in a subject, including a human, comprising the step of administering to said subject in need thereof a therapeutically effective amount of the antibody or antigen-binding portion thereof according to any one of claims 1-

20 or the pharmaceutical composition according to any one of claims 21-22, wherein said antibody or portion inhibits M-CSF.

24. The method of claim 23 wherein the condition is rheumatoid arthritis.

25. The method of claim 23 or 24 wherein the antibody utilized is monoclonal anti-M-CSF antibody 8.10.3F or 9.14.4I.

26. A method for treating a solid tumor such as a sarcoma, a carcinoma or a lymphoma in a subject, including a human, comprising the step of administering to said subject the antibody or antigen-binding portion thereof according to any one of claims 1-20 or the pharmaceutical composition according to any one of claims 21-22, wherein said antibody or antigen-binding portion inhibits M-CSF binding to *c-fms*.

27. An isolated cell line that produces the antibody or antigen-binding portion thereof according to any one of claims 1-20 or the heavy chain or light chain of said antibody or said antigen-binding portions thereof.

28. The cell line according to claim 27 that produces an antibody selected from the group consisting of: antibody 252, 88, 100, 3.8.3, 2.7.3, 1.120.1, 9.14.4I, 8.10.3F, 9.7.2IF, 9.14.4, 8.10.3, 9.7.2, 9.7.2C-Ser, 9.14.4C-Ser, 8.10.3C-Ser, 8.10.3-CG2, 9.7.2-CG2, 9.7.2-CG4, 9.14.4-CG2, 9.14.4-CG4, 9.14.4-Ser, 9.7.2-Ser, 8.10.3-Ser, 8.10.3-CG4, 8.10.3FG1 and 9.14.4G1 and an antibody that has the same amino acid sequence as one of the foregoing antibodies.

29. An isolated nucleic acid molecule comprising a nucleotide sequence that encodes the heavy chain or an antigen-binding portion thereof or the light chain or an antigen-binding portion thereof of an antibody according to any one of claims 1-20.

30. A vector comprising the nucleic acid molecule according to claim 29, wherein the vector optionally comprises an expression control sequence operably linked to said nucleic acid molecule.

31. A host cell comprising the vector according to claim 30 or the nucleic acid molecule according to claim 29.

32. The host cell according to claim 31 comprising a nucleic acid molecule encoding the heavy chain and a nucleic acid molecule encoding the light chain of an antibody or antigen-binding portion according to any one of claims 1-20.

33. A method of making an anti-M-CSF antibody or antigen-binding portion thereof, comprising culturing the host cell according to claim 32 or the cell line according to claim 28 under suitable conditions and recovering said antibody or antigen-binding portions.

34. A method for making a human monoclonal antibody that specifically binds to M-CSF, comprising the steps of:

- a) immunizing a non-human transgenic animal that is capable of producing human antibodies with M-CSF, an immunogenic portion of M-CSF or a cell or tissue expressing M-CSF;
- b) allowing said transgenic animal to mount an immune response to M-CSF; and
- c) isolating B lymphocytes from said transgenic animal.

1 / 24




FIG. 1A

2 / 24






FIG. 1B

3/24

FIG. 2A




4/24




5/24

FIG. 3A



6 / 24

FIG. 3B



7/24

FIG. 4A

Germine V=O12, J=JK3

252

-----GF-----T-----F-----V-----
 DIQMTQSPSSLSASVGRVTITC RASQISSYLN WYQKFGKAPKILLY RASSLSQGVPSRFSGSGSGTDFTLTISSLQPEDFATYYC QQSYSTPET
 FR1 CDR1 FR2 CDR2 FR3 CDR3

----- (residues 21-127 of SEQ ID NO: 4)

FGPGTKVDIK (SEQ ID NO: 103)

J

FIG. 4B

Germine V=O12, J=JK3

88

-----P-D-----L-----
 DIQMTQSPSSLSASVGRVTITC RASQISSYLN WYQKFGKAPKILLY RASSLSQGVPSRFSGSGSGTDFTLTISSLQPEDFATYYC QQSYSTPET
 FR1 CDR1 FR2 CDR2 FR3 CDR3

----- (residues 21-127 of SEQ ID NO: 8)

FGPGTKVDIK (SEQ ID NO: 103)

J

FIG. 4C

Germine V=I12, J=JK3

100

-----S-----D-I-----
 EIVMTQSPATLSVSPGERATLSC RASQSVSSNLA WYQKFGKAPKILLY GASTRATGIPARFSGSGSGTEFTLTISSLQSEDFAVYYC QQYNINRPET
 FR1 CDR1 FR2 CDR2 FR3 CDR3

----- (residues 21-127 of SEQ ID NO: 12)

FGPGTKVDIK (SEQ ID NO: 107)

J

FIG. 4D

Germline V=L5, J=JK3
 3.8.3
 (residues 23-130 of SEQ ID NO: 16)
 -----S-----D--G-----S--T--H-----T-----
 Germ DIQMTQSPSSVSASVGRVTITC RASQGISWLA WYQKQKAPKLLY RASSLOS GVPSRFSSGSGTDFTLTISSLQPEDFATYYC QANSFELT FGGKVDIKR
 (SEQ ID NO: 109) FR1 CDR1 FR2 CDR2 FR3 CDR3 J

FIG. 4E

Germline V=L5, J=JK4
 2.7.3
 (residues 23-130 of SEQ ID NO: 20)
 -----D-----R-----Q-----E-----N-----S-----T-----
 Germ DIQMTQSPSSVSASVGRVTITC RASQGISWLA WYQKQKAPKLLY RASSLOS GVPSRFSSGSGTDFTLTISSLQPEDFATYYC QANSFELT FGGKVEIKR
 (SEQ ID NO: 117) FR1 CDR1 FR2 CDR2 FR3 CDR3 J

FIG. 4F

Germline V=B3, J=JK1
 1.120.1
 -----I--F-----R-----N-----S-----
 Germ DIQMTQSPDLSAVSLGERATINC EGSSSVLYSSNNKNYLA WYQKQKQPKLLY MASTRFS GVPSRFSSGSGTDFTLTISSLQEDVAVYYC QQYSTWT
 (residues 21-134 of SEQ ID NO: 24) FR1 CDR1 FR2 CDR2 FR3 CDR3 J
 1.120.1
 Germ FGGTKVEIKR (SEQ ID NO: 112)

FIG. 4G

Germline V=3-11, D=D7-27, J=JH6

252

-----I-----G-----H-----
 QVQLVESGGGLVQPGGSLRLSCAAS GFTESSDYMS WIRQAPCKGLEWS YISSGGSTIYADSVKG RTTISRDNAKNSLYLQMNSLRRAEDTAVYYCAR ALGGMDV
 FR1 CDR1 FR2 CDR2 FR3 CDR3

252 ----- (residues 20-136 of SEQ ID NO: 2)

Germ NGQSTIVTVSSA (SEQ ID NO: 106)

FR4

FIG. 4H

Germline V=3-7, D=6-13, J=JH4

88

-----P-----RAY#
 EVQLVESGGGLVQPGGSLRLSCAAS GFTESSYMS WIRQAPCKGLEWS NIKODGEKYYVDVSKG RTTISRDNAKNSLYLQMNSLRRAEDTAVYYCAR GIMAGYFDY
 FR1 CDR1 FR2 CDR2 FR3 CDR3

88 ----- (residues 20-138 of SEQ ID NO: 6)

Germ NGQSTIVTVSSA (SEQ ID NO: 105)

FR4

FIG. 4I

Germline V=3-23, D=D1-26, J=JH4

100

-----R-R-E-----P-V-EG-R-GF-----
 EVQLVESGGGLVQPGGSLRLSCAAS GFTESSYMS WIRQAPCKGLEWS AISGGSGSTIYADSVKG RTTISRDNSKNTLYLQMNSLRRAEDTAVYYCAR ##YSGSYIEDY
 FR1 CDR1 FR2 CDR2 FR3 CDR3

100 ----- (residues 20-141 of SEQ ID NO: 10)

Germ NGQSTIVTVSSA (SEQ ID NO: 104)

FR4

10 / 24

FIG. 4J

Germline V=3-11, D=D7-27, J=JH4

3.8.3

Germ

QVQLVESGGGLVKEGGLRLSCAAS GFESDYMS WIRQAPGKLEWVS YISSSGSTIYADSVKG RFTISRDNKNSLYLQMSLRADFTAYYCAR #LTGDV
FR1 CDR1 FR2 CDR2 FR3 CDR3

3.8.3 (residues 20-135 of SEQ ID NO: 14)

Germ

WGQGLTVTVSSA (SEQ ID NO: 108)
FR4

FIG. 4K

Germline V=3-33, D=D1-26, J=JH4

2.7.3

Germ

QVQLVESGGGVQPRGSLRLSCAAS GETFSYGNH WVROAPGKLEWVA VIWYDGSNKYVADSVKG RETISRDNKNTLYLQMSLRADFTAYYCAR GYS#VEDY
FR1 CDR1 FR2 CDR2 FR3 CDR3

2.7.3 (residues 20-137 of SEQ ID NO: 18)

Germ

WGQGLTVTVSSA (SEQ ID NO: 110)
FR4

FIG. 4L

Germline V=1-18, D=D4-23, J=JH4

1.120.1

Germ

QVQLVSGAEVKKPGASVKYSCKAS GVTFTSYGIS WVROAPGQGLEWNG WISAYNGNTNVAOKLQ RVYMTDTSTSTAYMELRLSRSDDTAYYCA# #DYGGNVFDY
FR1 CDR1 FR2 CDR2 FR3 CDR3

1.120.1 (residues 20-139 of SEQ ID NO: 22)

Germ

WGQGLTVTVSSA (SEQ ID NO: 111)
FR4

11/24

FIG. 4M

Germline V-A27, J=JK4
 8.10.3
 Germ

 -F-----
 EVLVTSPTLSLSPGERATLSC RASQVSSSYLA WYQKCGQAPRLIY GASSRAT GIPDRFSGSGGTDETLTISRLEPEDFAVYC
 FR1 FR2 CDR1 CDR2 FR3

8.10.3 (residues 21-129 of SEQ ID NO: 44)

Germline V-VH3-48, D=D1-26, J=JH4b
 8.10.3
 Germ

 -F-----
 QQVGSPLT FGGKVEIKR (SEQ ID NO: 114)
 CDR3

FIG. 4N

Germline V-VH3-48, D=D1-26, J=JH4b
 8.10.3
 Germ

 -F-----
 EVLVSGGLVQPGGSLRLSCAAS GTFSSYSMN WVRCAPGKLEWYS YISSSSSTIYADVSK RETISRDNARNSLYLOMSLRDEPTAVYCAR
 FR1 CDR1 FR2 CDR2 FR3

8.10.3 (residues 20-141 of SEQ ID NO: 30)

Germline V-VH3-48, D=D1-26, J=JH4b
 8.10.3
 Germ

 -F-----
 WGGTLVTVSSA (SEQ ID NO: 113)
 CDR3

FIG. 4O

Germline V=012, J=JK3
 9.14.4
 (residues 23-130 of SEQ ID NO: 28)

Germline V=012, J=JK3
 9.14.4
 Germ

 -P-----
 DIQMTQSPFSLASVDGVITTC RASQISSYLN WYQKCPKAPKILY AASSLAS GVPSRFGSGGTDETLTISRLQPEDFAVYC QQSYSTPPT
 FR1 CDR1 FR2 CDR2 FR3 CDR3

FIG. 4P

Germline V=VH3-11, D=D7-27, J=JH4b

9.14.4

Germ QVQLVESGGGLVFPGGSLRLSCAAS GFTFSDYMS WIRQAPGKLEWVS YISSSGSTIYYADSVKG RTISRNNAKNSLYLQMNSTRAEDTAVYYCA #LTGDY
FR1 CDR1 FR2 CDR2 FR3 CDR3

9.14.4 ----- (residues 20-135 of SEQ ID NO: 38)

Germ WGQGTLLTVSSA (SEQ ID NO: 116)

J

FIG. 4Q

Germline V=012, J=JK3

9.7.2

(residues 23-130 of SEQ ID NO: 48)
Germ DIQMTQSPFSLASIVGDRVITTC RASQISSTLYN WYQKPGKAPKLLIY AASSIAS GVPSRFSGSGSGTDFTLTISLQPEDENFYTC QDSYSTPEF FGPGTKVDIKR
FR1 CDR1 FR2 CDR2 FR3 CDR3 J

FIG. 4R

Germline V=VH3-11, D=D6-13, J=JH4b

9.7.2

Germ QVQLVESGGGLVFPGGSLRLSCAAS GFTFSDYMS WIRQAPGKLEWVS YISSSGSTIYYADSVKG RTISRNNAKNSLYLQMNSTRAEDTAVYYCA #LTGDY
FR1 CDR1 FR2 CDR2 FR3 CDR3

9.7.2 ----- (residues 20-136 of SEQ ID NO: 46)

Germ WGQGTLLTVSSA (SEQ ID NO: 115)

J

12/24

FIG. 4S

Germline V=012, J=JK3

9.14.4I

(residues 23-130 of SEQ ID NO: 28)

Germ
 (SPQ ID NO: 103) FR1
 DIQWTOSTFSSLSASVGRVTITC
RASQSISSYIN
WYQKPGKAPKLLIY
AASSLQS
GVPKFSGSGSGTDTLTWISSLPEDFATYYC
QQSYSTPTT
EQPGTKVDIKR

FR1

CDR1

FR2

CDR2

FR3

CDR3

J

FIG. 4T

Germline V=VH3-11, D=D7-27, J=JH4b

9.14.4I

Germ

OYVLVESGGGLVPGGSLRLSCAAS
 GFTFSDYINS
WIRQAPKGLEWVS
YISSSGTIYADSVK
RTISRDNAKNSLYLQWNSLR
AEEDFATYYCAR
ILTGDY

FR1

CDR1

FR2

CDR2

FR3

CDR3

13/24

----- (residues 20-135 of SEQ ID NO: 26)

Germ
 9.14.4I
 WGGTTLTVSSA (SEQ ID NO: 116)

J

FIG. 4U

Germline V=J27, J=JK4

8.10.3F

Germ

EIVLTQSTGLSISSPERATISLC
 RASQSVSSSYIA
WYQKPGQAPRLIY
GASSRAT
GIPORFSGSGGTDTLTISRLEPEDFATYYC

FR1

CDR1

FR2

CDR2

FR3

8.10.3F

Germ

QOYGSSTPLT
 EGGKTKVEIKR
 (residues 21-129 of SEQ ID NO: 32)

CDR3

J

FIG. 4V

Germline V=VH3-48, D=D1-26, J=JH4b
8.10.3F

```

-----F-T-----R-S-----DELLA-ATP-----
Germline V=VH3-48, D=D1-26, J=JH4b
8.10.3F
-----F-T-----R-S-----DELLA-ATP-----
EVQLVESGGIVQPGGSLRLSCAAS GFTFFSYNN WVRQAPGKLEWVS YISSSSTIYVADSVKQ RFTISRNKAKNSLYLQNNISLRDFTAVYYCAR ##IVG##EFDY
FR1 CDR1 FR2 CDR2 FR3 CDR3
----- (residues 20-141 of SEQ ID NO: 30)
HGGGTIVTSSA (SEQ ID NO: 113)
J

```

14 / 24

FIG. 4W

Germline V=012, J=JK3
9.7.21F

(residues 23-130 of SEQ ID NO: 36)

```

-----GF-I-----R-----T-----J-----
Germline V=012, J=JK3
9.7.21F
-----GF-I-----R-----T-----J-----
DIQWTQSPSSLSASVGRVTTC RASQSISSYLN WYQOKPGKAPKLLIV AASSLQSGVPSRFGSGSGTDFTLTISIIQPEDEFTYVC QQSYSYPTFT FGGFTKVDIKR
FR1 CDR1 FR2 CDR2 FR3 CDR3 J
----- (residues 23-130 of SEQ ID NO: 36)
HGGGTIVTSSA (SEQ ID NO: 113)
J

```

FIG. 4X

Germline V=VH3-11, D=D6-13, J=JH6b
9.7.21F

```

-----R-G-----
Germline V=VH3-11, D=D6-13, J=JH6b
9.7.21F
-----R-G-----
QVQLVESGGIVQPGGSLRLSCAAS GFTFSDYNN WVRQAPGKLEWVS YISSSGCTIYVADSVKQ RFTISRNKAKNSLYLQNNISLRDFTAVYYCAR ##IVGNDV
FR1 CDR1 FR2 CDR2 FR3 CDR3
----- (residues 20-136 of SEQ ID NO: 34)
HGGGTIVTSSA (SEQ ID NO: 115)
J

```


FIG. 4Y

Germline V=012, J=JK3

9.7.2C-Ser

(residues 23-130 of SEQ ID NO: 52)

Germ

(SEQ ID NO: 103)

-----GF-I-----F-----R-G-----
 FRL CDR1 FR2 CDR2 FR3 CDR3 J
 DIOMTQSPFSSLSASVGRVITTC RASQSISSYLN WYQKQKAPKLLIY AASSLQS GVPKRFSGSGSGTDFTLTISSLQPEDFATYYC QQSISTPFT FPGTKVDIKR
 FRL CDR1 FR2 CDR2 FR3 CDR3 J

FIG. 4Z

Germline V=VH3-11, D=D6-13, J=JH6b

9.7.2C-Ser

Germ

(residues 20-136 of SEQ ID NO: 50)

-----GF-I-----F-----R-G-----
 FRL CDR1 CDR2 FR3
 QVQLVESGGGLYKPGSLRLSQAAS GFTFSDYMS WIRQAPGKGLWVS YISSGSGTIYADSVKRG RFTISRNANKSLYLQNSLRAPETAVYYCA FIGHMDV
 FRL CDR1 CDR2 FR3
 WSGQTITVTYSSA (SEQ ID NO: 115) J

FIG. 4AA

Germline V=012, J=JK3

9.14.4C-Ser

(residues 23-130 of SEQ ID NO: 56)

Germ

(SEQ ID NO: 103)

-----P-I-----F-----R-G-----
 FRL CDR1 CDR2 FR3
 DIOMTQSPFSSLSASVGRVITTC RASQSISSYLN WYQKQKAPKLLIY AASSLQS GVPKRFSGSGSGTDFTLTISSLQPEDFATYYC QQSISTPFT FPGTKVDIKR
 FRL CDR1 CDR2 FR3 CDR3 J

FIG. 4BB

Germline V=VH3-11, D=D7-27, J=JH4b

9.14.4C-Ser

Genin	CDR1	FR2	CDR2	FR3	CDR3	LTGID	Gen
9.14.4C-Ser	QVQLVESGGGLVKPGGSLRLSCAAS	GFTESDYMS	WIRQAPKGLFWIS	YISSSGSTIYYADSVK	RTISRDNAKNSLTQNNISRAETIAVYYCAR		G

9.14.4C-Ser ----- (residues 20-135 of SEQ ID NO: 54)

Germ WGQGT LVT VSSA (SEQ ID NO: 116)

FIG. 4CC

Germline V-A27. J=JK4

8.10.3C-Ser

8.10.3C-Ser	Germ		Germ	
	FR1	CDR1	FR2	CDR2
	EVLTQSEPTLSLSPGERATLSC	RASQSVSSSYIA	WYQQKFGQAPRLIY	GASSRAT
				CDR2
				FR3
				GIPDFSGSGSDFTLTISRLEPDEFAVYVC

Germ QQYGSPLT FGGKTVEIKR
CDR3 J (residues 21-129 of SEQ ID NO: 60)

8.10.3 ----- (SEQ ID NO: 114)

FIG. 4DD

Germline V=VH3-48, D=D1-26, J=JH4b

8.10.3C-Ser

[illegible]

8.10.3C-Ser ----- (residues 20-141 of SEQ ID NO: 58)

WGQGLTVTVSSA (SEQ ID NO: 113)

17/24

FIG. 4EE

Germine V-A27, J-JK4
8.10.3-CG2

Germ EIVLTQSPGTLSLSPGERATLSC RASQSVSSSYLA WYQKFGQAPRLIY CASSRAT GIPDRFSGSGGTDFLTISRLEPEDFAVYVC
FR1 FR2 CDR1 CDR2 FR3

8.10.3-CG2 ----- (residues 21-129 of SEQ ID NO: 60)

Germ QOYSSSPLT FGGKTVEIKR J
CDR3

FIG. 4FF

Germine V-VH3-48, D-D1-26, J-JH4b
8.10.3-CG2

Germ EVOLVEGGGLVQGGSLRLISCAAS GETFESSYMN WVROAFKGLWVS YISSSSSTIYADSVKG RETISRDNAKNSLYLQWNSIRDEPTAVYCAR ###IVG###FDY
FR1 FR2 CDR1 CDR2 FR3 CDR3

8.10.3-CG2 ----- (residues 20-141 of SEQ ID NO: 62)

Germ WGQGTLYTVSSA J
CDR3

FIG. 4GG

Germine V-012, J-JK3
9.7.2-CG2

(residues 23-130 of SEQ ID NO: 52)

Germ DIQMTQSPSSLSASVDGVAVITC RASQISSYLN WYQKPKGAPRLIY RASSLAS GVPNSFSGSGGTDFLTISRLEPEDFAVYVC QASYSFTFT FGCTKVDIKR
FR1 CDR1 FR2 CDR2 CDR3 J

18/24

FIG. 4HH

Germine V=VH3-11, D=D6-13, J=JH6b
 9.7.2-CG2 ----- R-G-----
 (residues 20-136 of SEQ ID NO: 66)

Germ QVQLVESGGGLVFRGSGSLRLSCAAS GFTSDYMS WIRQAPGKGLEWVS YISSSGSTIYYADSVKG RTTISRDNARKNSLYIQMNSLRADTAVYCA #IHGMDV
 (SEQ ID NO: 115) FR1 FR2 FR3 CDR2 CDR3

9.7.2-CG2 -----
 Germ WQQGTTVTIVSSA J

FIG. 4II

Germine V=VH3-11, D=D6-13, J=JH6b
 9.7.2-CG4 -----
 (residues 23-130 of SEQ ID NO: 52)

Germ DIONIQSPFSLASVGDVRVITTC RASQGISYLN WYQKPGKAPKLLI AASSLOS GVPSPRSGSGGTUFTLTISLQPEDFAIVYC QQSYSPTPT FQPGTKVDIKR
 (SEQ ID NO: 103) FR1 FR2 FR3 CDR1 CDR2 CDR3 J

FIG. 4JJ

Germine V=VH3-11, D=D6-13, J=JH6b
 9.7.2-CG4 ----- R-G-----
 (residues 20-135 of SEQ ID NO: 70)

Germ QVQLVESGGGLVFRGSGSLRLSCAAS GFTSDYMS WIRQAPGKGLEWVS YISSSGSTIYYADSVKG RTTISRDNARKNSLYIQMNSLRADTAVYCA #IHGMDV
 (SEQ ID NO: 115) FR1 FR2 FR3 CDR1 CDR2 CDR3

Germ WQQGTTVTIVSSA J

FIG. 4KK

Germline V=012, J=JK3

9.14.4-CG2
(residues 23-130 of SEQ ID NO: 56)

Germ
 (SEQ ID NO: 103)

DIQMTQPSLSASVGDRTVTC FR1 CDR1 FR2 CDR2 FR3 CDR3 J
 WTQQRPGKAPKALII AASSILOS GVPSRFGSGSGTDTLTISLSQEPEDPATYIC QQSYSTPFT EPGGTQVDIKR

FIG. 4LL

Germline V=VH3-11, D=D7-27, J=JH4b

9.14.4-CG2

Germ
 (SEQ ID NO: 103)

QVQLVESGGGLVPRGGSRLSCAS FR1 CDR1 FR2 CDR2 FR3 CDR3 J
 WTQQRPGKAPKALII AASSILOS GVPSRFGSGSGTDTLTISLSQEPEDPATYIC QQSYSTPFT EPGGTQVDIKR

9.14.4-CG2

(residues 20-135 of SEQ ID NO: 74)

Germ

WTQQRPGKAPKALII
 (SEQ ID NO: 116)

FIG. 4MM

Germline V=012, J=JK3

9.14.4-CG4
(residues 23-130 of SEQ ID NO: 56)

Germ
 (SEQ ID NO: 103)

DIQMTQPSLSASVGDRTVTC FR1 CDR1 FR2 CDR2 FR3 CDR3 J
 WTQQRPGKAPKALII AASSILOS GVPSRFGSGSGTDTLTISLSQEPEDPATYIC QQSYSTPFT EPGGTQVDIKR

FIG. 4NN

Germline V=VH3-11. D=D7-27. J=JH4b

9.14.4-CG4

Germ

<u>QVQLVESGGGLVKGPGGSLRLSCAAS</u>	<u>GFTFSDYYMS</u>	<u>WIRQAPKGLEWVS</u>	<u>YISSSGSTYYADSVKG</u>	<u>RFTISRDNAKNSLYLQMNSLR</u>	<u>EDTAVYYCAR</u>	<u>#LTGDI</u>
FR1	CDR1	FR2	CDR2	FR3		CDR3

9.14.4-CG4 ----- (residues 20-135 of SEQ ID NO: 78)

Germ
NGOGTLVTVSSA (SEQ ID NO: 116)

1

FIG. 400

Germline V=012. J=JK3

9.14.4-Ser

(residues 23-130 of SEQ ID NO: 28)

Germ	CDR1	CDR2	CDR3	J
DIQMTQSPSSLSASVGDRVTTC	RASQSTSSYLN	WYQOKPGKAPKLLIY	GVPSRFSGSGSGTDFTLTISLSLPEDFATYYC	QQSYSTPET
(SEQ ID NO: 103)	FR1	FR2	FR3	FGPGTKVDIKIR

H-T-I-d-

FIG. 4PP

Germline V=VH3-11, D=D7-27, J=JH4b

9.14.4-Ser

Germ

<u>QVQLVESGGGLV</u>	<u>KPGGSL</u>	<u>RLSCAAS</u>	<u>GTFTSDYMS</u>	<u>WIRQAPGKLEWVS</u>	<u>YISSSGSTYYADSVKG</u>	<u>RTGISRDNAKNSLY</u>	<u>LOMNSLR</u>	<u>AEATAVYYCAR</u>	<u>#LTGDI</u>
	FR1		CDR1	FR2	CDR2	FR3			CDR3

9.14.4-Ser ----- (residues 20-135 of SEQ ID NO: 82)

Germ WGOGTLVTVSSA (SEQ ID NO: 116)

1

FIG. 4QQ

Germline V=012, J=JK3

9.7.2-Ser

FR1-----GF-I-----R-----T-----L-----E-----FR3-----CDR3-----J

(residues 23-130 of SEQ ID NO: 40)

Germline DIQMTQSPSSLSASVGDRVTITC RASQISISYLN WYQKPKGAKPLLY AASSLOS GVPSRFGSGSGTDTLTISLQPEDFAVYVC QQSYSITPT FPGGTVVDLKE
(SEQ ID NO: 103) FR1 CDR1 FR2 CDR2 FR3 CDR3

FIG. 4RR

Germline V=VRH3-11, D=D6-13, J=JH6b

9.7.2-Ser

FR1-----CDR1-----FR2-----CDR2-----FR3-----R-Q----

(residues 20-136 of SEQ ID NO: 50)

Germline QVQLVESGGGLVKPGGSLRLSCAAS GTFSDYNS WIRQAPGKLEWVS YISSSGSTIYVDSYKG RFTTISRDNAKNSLYLNMSIRAEFTAVYCA #1#GMDV
(SEQ ID NO: 115) FR1 CDR1 FR2 CDR2 FR3 CDR3

9.7.2-Ser

WGQGTIVTVSSA (SEQ ID NO: 115)

J

FIG. 4SS

Germline V=A27, J=JK4

8.10.3-Ser

FR1-----F-----FR2-----FR3-----V-----

(residues 21-129 of SEQ ID NO: 44)

Germline EVLTQSPGTLISLSPGERATLSC RASQSVSSYLA WYQKPGQAPRLLY GASSRAT GIPORTFGSGSGTDTLTITISRLIEPEDFAVYVC
(SEQ ID NO: 114) FR1 CDR1 FR2 CDR2 FR3

8.10.3-Ser

QVGSSELT FGGGTVKVEIKR (SEQ ID NO: 114)

J

CDR3

FIG. 4TT

Germline V=VH3-48, D=D1-26, J=JH4b
8.10.3-Ser

-----F-T-----R-S-----DPLA-ATP-----
 EVQLVESGGGLVQPGGSLRLSCAAS GFTFSSYSN WVRQAPGKLEWVS YISSSSITYYADSVKG RTTISRQAKNSLYIQMNSLRDETFVYYCAR ###IVG##FDY
 CDR1 FR1 CDR2 FR3 CDR3

8.10.3-Ser ----- (residues 20-141 of SEQ ID NO: 90)

Germline V=VH3-48, D=D1-26, J=JH4b
8.10.3-Ser

EVQLVESGGGLVQPGGSLRLSCAAS GFTFSSYSN WVRQAPGKLEWVS YISSSSITYYADSVKG RTTISRQAKNSLYIQMNSLRDETFVYYCAR ###IVG##FDY
 CDR1 FR1 CDR2 FR3 CDR3

FIG. 4UU

Germline V=VH3-48, D=D1-26, J=JH4b
8.10.3-Ser

-----F-T-----R-S-----DPLA-ATP-----
 EVQLVESGGGLVQPGGSLRLSCAAS GFTFSSYSN WVRQAPGKLEWVS YISSSSITYYADSVKG RTTISRQAKNSLYIQMNSLRDETFVYYCAR ###IVG##FDY
 CDR1 FR1 CDR2 FR3 CDR3

8.10.3-Ser ----- (residues 21-129 of SEQ ID NO: 60)

Germline V=VH3-48, D=D1-26, J=JH4b
8.10.3-Ser

EVQLVESGGGLVQPGGSLRLSCAAS GFTFSSYSN WVRQAPGKLEWVS YISSSSITYYADSVKG RTTISRQAKNSLYIQMNSLRDETFVYYCAR ###IVG##FDY
 CDR1 FR1 CDR2 FR3 CDR3

FIG. 4W

Germline V=VH3-48, D=D1-26, J=JH4b
8.10.3-Ser

-----F-T-----R-S-----DPLA-ATP-----
 EVQLVESGGGLVQPGGSLRLSCAAS GFTFSSYSN WVRQAPGKLEWVS YISSSSITYYADSVKG RTTISRQAKNSLYIQMNSLRDETFVYYCAR ###IVG##FDY
 CDR1 FR1 CDR2 FR3 CDR3

8.10.3-Ser ----- (residues 20-141 of SEQ ID NO: 94)

Germline V=VH3-48, D=D1-26, J=JH4b
8.10.3-Ser

EVQLVESGGGLVQPGGSLRLSCAAS GFTFSSYSN WVRQAPGKLEWVS YISSSSITYYADSVKG RTTISRQAKNSLYIQMNSLRDETFVYYCAR ###IVG##FDY
 CDR1 FR1 CDR2 FR3 CDR3

FIG. 4WW

Germine V=012, J=JK3
 9.14.4G1
 (residues 23-130 of SEQ ID NO: 28)

Germ (SEQ ID NO: 103) DIQWTFQPSLSASVGRVTTC PASQTSYLN WYQKCKAPKLLIY RASSIOS CVPFRSCSGCTDFTLISLQPEDFATVYC QQSYSTPFT FGPCTKVDIKR

FR1 CDR1 FR2 FR3 CDR3 J

FIG. 4XX

Germine V=VH3-11, D=D7-27, J=JH4b
 9.14.4G1

Germ (SEQ ID NO: 103) QVQLVSGGGLYKFGGSLRLSCAS GFETSDYMS WIRQAFKGLWVS YISSGSIYVADSVK RTISRDNANKSYIQWSIRRETFATVYCAR #LTGDY

FRI FR2 CDR2 FR3 CDR3 J

9.14.4G1 (residues 20-135 of SEQ ID NO: 102)

Germ (SEQ ID NO: 116) WGQGLTVTSSA (SEQ ID NO: 116)

23 / 24

FIG. 4YY

Germine V=427, J=JK4
 8.10.3FG1

Germ (SEQ ID NO: 103) EIVLTQSPFTLSLSPGERATLSC RASQVSSYLA WYQKCKAPKLLIY QASSRAT GLPDRFSGSGGTDTLTISRLQPEDFATVYC

FRI CDR1 FR2 CDR2 FR3 J

8.10.3FG1 (residues 21-129 of SEQ ID NO: 32)

Germ (SEQ ID NO: 114) QQYGSSELT FGGKTVEIKR (SEQ ID NO: 114)

CDR3 J

FIG. 4ZZ

Germline V=VH3-48, D=D1-26, J=JH4b
8.10.3Fc1

Germline	EVQLVESGGGLVQPGGSLRLSQAAS	GFTESSYSMN	WVROAPGKGLQWVS	YISSSSTIYADSVKG	RTISRDNAKNSLYIQMNSLRDEDTAVYYCAR	###IVG##FDY
	FR1	CDR1	FR2	CDR2	FR3	CDR3
8.10.3Fc1	----- (residues 20-141 of SEQ ID NO: 96)					
Germline	WGQGTIVTSSA (SEQ ID NO: 113)					

SEQUENCE LISTING

<110> ABGENIX, INC.
 WARNER-LAMBERT COMPANY LLC
 BEDIAN, VAHE
 DEVALARAJA, MADHAV NARASIMHA
 FOLTZ, IAN
 HAAK-FRENDSCHO, MARY
 KELLERMANN, SIRID-AIMEE
 LOW, JOSEPH EDWIN
 MOBLEY, JAMES LESLIE

<120> ANTIBODIES TO M-CSF

<130> ABX-PF4 PCT

<140>

<141>

<150> 60/502,163

<151> 2003-09-10

<160> 117

<170> PatentIn Ver. 3.2

<210> 1

<211> 1383

<212> DNA

<213> Homo sapiens

<400> 1

```

atggagttgg  ggctgtgctg  gattttccct  gttgctatta  taaaagggtg  ccagtgctag  60
gtgcagctgg  tggagctctg  gggaggcttg  gtcaagcctg  gaggggtcc  gagactctcc  120
tgtgcagcct  ctggattcac  ctccagtgac  tactacatga  gctggatccg  ccaggctcca  180
gggaaggggc  tggagtggt  ttcatacat  agtggtagtg  gtagtaccat  atactacgca  240
gactctgtga  agggccgatt  caccatctcc  agggacaacg  ccaagaactc  actgtatctg  300
caaatgaaca  gccctgagagc  cgaggacacg  gccgtgtatc  actgtgcgag  agccctgggt  360
gggatggagc  tctggggcca  agggaccacg  gtccacgtct  cctcagcttc  caccaagggc  420
ccatccgtct  tccccctggc  gccctgtct  agaagcacct  ccgagagcac  agcggccctg  480
ggctgccctg  tcaaggacta  ctccccgaa  ccggtgacgg  tgtcgtggaa  ctcaggcgct  540
ctgaccagcg  gcgtgcacac  ctccccagct  gtccctacag  cctcaggact  ctactccctc  600
agcagcgtgg  tgaccgtgcc  ctccagcaac  ttccggcccc  agacctacac  ctgcaacgta  660
gatcacaagc  ccagcaaac  caaggtggac  aagacagtgg  agcgcaaatg  ttgtgtcgag  720
tgcccacgt  gccacgcacc  accctgtggc  ggaccgtcag  tcttccctct  ccccccaaaa  780
cccaaggaca  cctcatgat  ctcccggaac  cctgaggtca  cgtgcgtggt  ggtggacgtg  840
agccacgaag  accccgaggt  ccagttcaac  tggtaoigtg  acggcgtgga  ggtgcataat  900
gccaaagaca  agccacggga  ggagcagttc  aacagcacgt  tccgtgtggt  cagcgtctcc  960
accgttgtgc  accaggactg  gctgaacggc  aaggagtaca  agtgcaaggt  ctccaacaaa  1020
ggcctccacg  ccccatcga  gaaaaccatc  tccaaaacca  aagggcagcc  ccgagaacca  1080
caggtgtaca  cctctccccc  atcccgggag  gagatgacca  agaaccaggt  cagcctgacc  1140
tgctcggtca  aaggcttcta  cccacgcac  atgcgcgtgg  agtggggag  caatggggag  1200
ccggagaaca  actacaagac  cacacctccc  atgtggact  ccgacggctc  cttcttctc  1260
tacagcaagc  tcaccgtgga  caagagcagg  tggcagcagg  ggaacgtctt  ctcatgctcc  1320
gtgatgcag  aggcctctga  caaccactac  acgcagaaga  gcctctccct  gtctccgggt  1380
aaa

```

<210> 2
 <211> 461
 <212> PRT
 <213> Homo sapiens

<400> 2
 Met Glu Leu Gly Leu Cys Trp Ile Phe Leu Val Ala Ile Ile Lys Gly
 1 5 10 15
 Val Gln Cys Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys
 20 25 30
 Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
 35 40 45
 Ser Asp Tyr Tyr Met Ser Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu
 50 55 60
 Glu Trp Ile Ser Tyr Ile Ser Gly Ser Gly Ser Thr Ile Tyr Tyr Ala
 65 70 75 80
 Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
 85 90 95
 Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val
 100 105 110
 Tyr His Cys Ala Arg Ala Leu Gly Gly Met Asp Val Trp Gly Gln Gly
 115 120 125
 Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe
 130 135 140
 Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu
 145 150 155 160
 Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp
 165 170 175
 Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu
 180 185 190
 Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser
 195 200 205
 Ser Asn Phe Gly Thr Gln Thr Tyr Thr Cys Asn Val Asp His Lys Pro
 210 215 220
 Ser Asn Thr Lys Val Asp Lys Thr Val Glu Arg Lys Cys Cys Val Glu
 225 230 235 240
 Cys Pro Pro Cys Pro Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu
 245 250 255
 Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu
 260 265 270

Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Gln
 275 280 285

Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys
 290 295 300

Pro Arg Glu Glu Gln Phe Asn Ser Thr Phe Arg Val Val Ser Val Leu
 305 310 315 320

Thr Val Val His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys
 325 330 335

Val Ser Asn Lys Gly Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys
 340 345 350

Thr Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser
 355 360 365

Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys
 370 375 380

Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln
 385 390 395 400

Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Met Leu Asp Ser Asp Gly
 405 410 415

Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln
 420 425 430

Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn
 435 440 445

His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 450 455 460

<210> 3

<211> 702

<212> DNA

<213> Homo sapiens

<400> 3

atgagggtcc ctgctcagct cctgggggctc ctgctactct ggctccgagg tgccagatgt 60
 gacatccaga tgacccagtc tccatctctc ctgtctgcat ctgtaggaga cagagtcacc 120
 atcacttgcc gggcaagtca gagcattagc ggcttttttaa attggtatca gcagaaacca 180
 gggaaaagccc ctaagctcct gatctatgct acatccagtt tgcaaatggtg ggtcccatcc 240
 aggtttcagtg gcagtgagatc tgggacagat ttcaactctca ccatcagcag tctgcgaacct 300
 gaagattttg caacttatta ctgtcaacag agttacagtg tccattccac tttcggccct 360
 gggaccaaag tggatatcaa acgaactgtg gctgcacatc ctgtcttcat cttcccgcca 420
 tctgatgagc agttgaaatc tgggaactgct agcgttgtgt gctgtgtgaa taactcttat 480
 ccagagagagg ccaaagtaca gtggaaggtg gataacgccc tccaatcggtg taactccagc 540
 gagagtgatca cagagcagga cagcaaggac agcacctaca gctcagcag caccctgacg 600
 ctgagctaaag cagactacga gaaacacaaa gtctacgcct gcgaagtcac ccatcagggc 660
 ctgagctcgc ccgtcacaaa gagcttcaac aggggagagt gt 702

<210> 4
 <211> 234
 <212> PRT
 <213> Homo sapiens

<400> 4
 Met Arg Val Pro Ala Gln Leu Leu Gly Leu Leu Leu Trp Leu Arg
 1 5 10 15
 Gly Ala Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser
 20 25 30
 Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Ser
 35 40 45
 Ile Ser Gly Phe Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro
 50 55 60
 Lys Leu Leu Ile Tyr Ala Thr Ser Ser Leu Gln Ser Gly Val Pro Phe
 65 70 75 80
 Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
 85 90 95
 Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ser Tyr
 100 105 110
 Ser Val Pro Phe Thr Phe Gly Pro Gly Thr Lys Val Asp Ile Lys Arg
 115 120 125
 Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln
 130 135 140
 Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr
 145 150 155 160
 Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser
 165 170 175
 Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr
 180 185 190
 Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys
 195 200 205
 His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro
 210 215 220
 Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
 225 230

<210> 5
 <211> 1389
 <212> DNA

<213> Homo sapiens

<400> 5

```

atggaatttg  ggctgtgctg  ggttttcctt  gttgctattt  tagaagggtg  ccagtgtgag  60
gtgcagctgg  tggagtctgg  gggaggcttg  gtccagcctg  gggggtcctt  gagactctcc  120
tgtgcagcct  ctggattcac  ctttagtagc  tattggatga  gctgggtccg  ccaggctcca  180
gggaaggggc  tggagtgggt  ggccaacata  aagcaagatg  gaagtggagaa  atactatgtg  240
gactctgtga  agggccgatt  caccatctcc  agagacaacg  ccaagaactc  actgtatctg  300
caaatgaaca  gcctgagagc  cgaggacacg  gctgtgtatt  actgtgtctc  gggatataga  360
gcagctggta  gggcctactg  gggccaggga  accctggta  ccgtctctcc  agcttccacc  420
aaggggccat  ccgtcttccc  cctggcgccc  tgtcttagaa  gcacctccga  gagcacagcg  480
gccccgggct  gcctgggtcaa  ggactacttc  ccgaaacogg  tgaocgtgtc  gtggaaactca  540
ggcgctctga  ccagcggcgt  gcacaccttc  ccagctgtcc  tacagtctcc  aggactctac  600
tccctcagca  gcgtgggtgac  cgtgccctcc  agcaacttcg  gcaccagac  ctacacctgc  660
aacgtagatc  acaagcccag  caacaccaag  gtggacaaga  cagttgagcg  caaatgttgt  720
gtcgagtgcc  caccgtgccc  agcaccacct  gtggcaggac  cgctcagttt  cctcttcccc  780
ccaaaaccca  aggacacctc  catgatctcc  cggacccctg  aggtcacgtg  cgtggtgggt  840
gacgtgagcc  acgaagaccc  cgaggctccg  ttcaactgg  acgtggacgg  cgtggaggtg  900
cataatgcc  agacaagacc  acgggaggag  cagttcaaca  gcacgttccg  tgtggtcagc  960
gtctctacgg  ttgtgcacca  ggactgggtg  aacggcaagg  agtacaagt  caaggtctcc  1020
aacaaggccc  tcccagcccc  catcgagaaa  accatctcca  aaaccaaagg  gcagcccga  1080
gaaccacagg  tbtacacctc  gccccatccc  cggggaggaga  tgaccaagaa  ccaggtcagc  1140
ctgacctgcc  tggctcaagg  cttctacccc  agcagacatc  ccgtggagtg  ggagagcaat  1200
gggcagccgg  agaaacaact  caagaccaca  cctcccatgc  tggactccga  cggctccttc  1260
ttctcttaca  gcaagctcac  cgtggacaag  agcaggtggc  agcaggggaa  cgtcttctca  1320
tgctccgtga  tgcatgaggg  tctgcacaac  cactacacgc  agaagagcct  ctccctctct  1380
ccgggtaaa  1389

```

<210> 6

<211> 463

<212> PRT

<213> Homo sapiens

<400> 6

```

Met Glu Phe Gly Leu Cys Trp Val Phe Leu Val Ala Ile Leu Glu Gly
  1             5             10             15

Val Gln Cys Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln
  20             25             30

Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
  35             40             45

Ser Ser Tyr Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
  50             55             60

Glu Trp Val Ala Asn Ile Lys Gln Asp Gly Ser Glu Lys Tyr Tyr Val
  65             70             75             80

Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
  85             90             95

Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val
 100             105             110

```

Tyr Tyr Cys Ala Pro Gly Ile Ala Ala Ala Gly Arg Ala Tyr Trp Gly
 115 120 125
 Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser
 130 135 140
 Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala
 145 150 155 160
 Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val
 165 170 175
 Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala
 180 185 190
 Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val
 195 200 205
 Pro Ser Ser Asn Phe Gly Thr Gln Thr Tyr Thr Cys Asn Val Asp His
 210 215 220
 Lys Pro Ser Asn Thr Lys Val Asp Lys Thr Val Glu Arg Lys Cys Cys
 225 230 235 240
 Val Glu Cys Pro Pro Cys Pro Ala Pro Pro Val Ala Gly Pro Ser Val
 245 250 255
 Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr
 260 265 270
 Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu
 275 280 285
 Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys
 290 295 300
 Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Phe Arg Val Val Ser
 305 310 315 320
 Val Leu Thr Val Val His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys
 325 330 335
 Cys Lys Val Ser Asn Lys Gly Leu Pro Ala Pro Ile Glu Lys Thr Ile
 340 345 350
 Ser Lys Thr Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro
 355 360 365
 Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu
 370 375 380
 Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn
 385 390 395 400
 Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Met Leu Asp Ser
 405 410 415

Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg
 420 425 430

Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu
 435 440 445

His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 450 455 460

<210> 7
 <211> 702
 <212> DNA
 <213> Homo sapiens

<400> 7
 atgagggtcc ctgctcagct cctggggctc ctgctactct ggctccgagg tgcagatgt 60
 gacatccaga tgacccagtc tccatcctcc ctgtctgcat ctgttggaga cagagtcacc 120
 atcaactgtcc ggccaagtca ggacatttagc agttatttaa attgggtatca gcagaaacca 180
 gggaaagccc ctaagctcct gatctatgct gcattccagtt tgcaaaagtgg ggtccattta 240
 aggttcagtg gcagtggtatc tgggacagat ttcactctca ccatcagcag tctgcaacct 300
 gaagattttg caacttacta ctgtcaacag agttacagta ccccatcac tttcgccct 360
 gggaccaaa tggaatatac acgaactgtg gctgcacat ctgtcttcat cttccgccca 420
 tctgatgagc agttgaaatc tggaaactgct agcgttgtgt gcctgctgaa taactcttat 480
 cccagagagg ccaaaagtaca gtggaagggt gataacgccc tccaatcggg taactcccag 540
 gagagtgtca cagagcagga cagcaaggac agcacctaca gcctcagcag caacctgaag 600
 ctgagcaaa gacactcaga gaaacacaaa gtctacgctt gcgaagtcac ccatcagggc 660
 ctgagctcgc cgtgcacaaa gagcttcaac aggggagagt gt 702

<210> 8
 <211> 234
 <212> PRT
 <213> Homo sapiens

<400> 8
 Met Arg Val Pro Ala Gln Leu Leu Gly Leu Leu Leu Trp Leu Arg
 1 5 10 15

Gly Ala Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser
 20 25 30

Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Pro Ser Gln Asp
 35 40 45

Ile Ser Ser Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro
 50 55 60

Lys Leu Leu Ile Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Leu
 65 70 75 80

Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
 85 90 95

Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ser Tyr
 100 105 110

Ser Thr Pro Phe Thr Phe Gly Pro Gly Thr Lys Val Asp Ile Lys Arg
 115 120 125

Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln
 130 135 140

Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr
 145 150 155 160

Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser
 165 170 175

Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr
 180 185 190

Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys
 195 200 205

His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro
 210 215 220

Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
 225 230

<210> 9

<211> 1398

<212> DNA

<213> Homo sapiens

<400> 9

atggagtttg ggcctcgcgt gatattttctt gtggctatatt taaaagggtgt ccagtgtgag 60
 gtgcagctgt tggagtcctgg gggaggcctg gtacagcctg gggggctccct gagactctcc 120
 tgtgcagcct ctggattcac ctttagcagc tatgccatga gctgggtccg ccaggctcca 180
 gggaaggggc tggaatgggt ctacgtattt agtggctcgt gtggtaggac atactctcga 240

gactccgtga agggccgggt caccatctcc agagacaatt ccaagaacac gctgtatctg 300
 caaatgaaca gcttgagagc cgaggacacg gccgtatat tctgtgcggt agaaggctat 360
 agtgggcgct acggattttt tgactactgg gccaggggaa ccctagtcaac cgtctctcca 420
 gccctcacca agggcccatc ggtcttcccc ctgggcgcct gctctagaag caoctccag 480
 agcacagcgg cctctggcgt cctgggtcaag gactacttcc ccgaaccggt gcagggtgtcg 540
 tggaactcag gcgtctctgac cagcggcgtg cacaccttcc cagctgtctc acagtctcca 600
 ggactctact cctcagcag cgtgggtgacc gtgccctcca gcaacttcgg caccagacc 660
 tacacctgca acgtagatca caagcccagc aacaccaagg tggacaagac agttagagcg 720
 aaatggttgg tgcagtgccc accgtgccca gcaccacctg tggcaggacc gtcagtcttc 780
 ctcttcccc caaaaccocaa ggacacctc atgatctccc ggacccctga ggtcagctgc 840
 gtgggtggtg acgtgagcca cgaagacccc gaggtccagt tcaactggta cgtggacggc 900
 gtggaggtgc ataatgccaa gacaaagcca cgggaggagc agttcaacag caogttccgt 960
 gtggtcagcg tctctcacgt tgtgcaccag gactgggtga acgycaggga gtaccaagtc 1020
 aaggtctcca acaaggccct cccagccccc atcgagaaaa ccatctccaa atacaaaggg 1080
 cagccccgag aaccacaggt gtacacctg ccccatctcc gggaggagat gcccaagaa 1140
 caggtcagcc tgacctgcct ggtcaaaagg ttctacccca gcgacatcgc gctggagttg 1200
 gagagcaatg ggcagccgga gaacaaactc aagaccacac ctcccatgct ggactccgac 1260
 ggctcctctc tctctacag caagctcacc gtggacaaga gcaggtggca gcaggggaac 1320
 gtctctctcat gtcctgtgat gcatgaggct ctgcacaacc actacacgca gaagagcctc 1380
 tccctgtctc cgggtaaa 1398

<210> 10
 <211> 466
 <212> PRT
 <213> Homo sapiens

<400> 10
 Met Glu Phe Gly Leu Arg Trp Ile Phe Leu Val Ala Ile Leu Lys Gly
 1 5 10 15
 Val Gln Cys Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln
 20 25 30
 Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
 35 40 45
 Ser Ser Tyr Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
 50 55 60
 Glu Trp Val Ser Ala Ile Ser Gly Arg Gly Gly Arg Thr Tyr Phe Ala
 65 70 75 80
 Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn
 85 90 95
 Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val
 100 105 110
 Tyr Phe Cys Ala Val Glu Gly Tyr Ser Gly Arg Tyr Gly Phe Phe Asp
 115 120 125
 Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys
 130 135 140
 Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu
 145 150 155 160
 Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro
 165 170 175
 Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr
 180 185 190
 Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val
 195 200 205
 Val Thr Val Pro Ser Ser Asn Phe Gly Thr Gln Thr Tyr Thr Cys Asn
 210 215 220
 Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys Thr Val Glu Arg
 225 230 235 240
 Lys Cys Cys Val Glu Cys Pro Pro Cys Pro Ala Pro Pro Val Ala Gly
 245 250 255
 Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile

260 265 270
 Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
 275 280 285
 Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
 290 295 300
 Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Phe Arg
 305 310 315 320
 Val Val Ser Val Leu Thr Val Val His Gln Asp Trp Leu Asn Gly Lys
 325 330 335
 Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ala Pro Ile Glu
 340 345 350
 Lys Thr Ile Ser Lys Thr Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
 355 360 365
 Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
 370 375 380
 Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
 385 390 395 400
 Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Met
 405 410 415
 Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
 420 425 430
 Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
 435 440 445
 Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
 450 455 460
 Gly Lys
 465

<210> 11
 <211> 702
 <212> DNA
 <213> Homo sapiens

<400> 11
 atggaagccc cagctcagct tctcttctct ctgctaactct ggctcccaga taccactgga 60
 gaaatagtga tgacgcagtc tccagccacc ctgtctgtgt ctccaggga aagagccacc 120
 ctctctctgca gggccagtc gagtgtagc agcaacttag cctggtacca gcagaaacct 180
 ggccaggctc ccaggctcct catctatggt gcaccacca gggccagtggt tatccagac 240
 aggatcagtg gcagtggttc tggaaacagag ttcaacttca tcatcagcag cctgcagtct 300
 gaagattttg cagtttatta ctgtcagcag tctaataact ggccattcac ttctggccct 360
 gggaccaaaag tggatatcaa acgaactgtg gctgcacccat ctgtcttcat ctcccgcca 420
 tctgatgagc agttgaaatc tggaaactgt agcgttgtgt gctgtgtgaa taactctcat 480
 cccagagagg ccaaagtaca gtggaagggt gataacgccc tccaatcggg taactccag 540

gagagtgtca cagagcagga cagcaaggac agcacctaca gcttcagcag caccctgacg 600
 ctgagcaaa cagactacga gaaacacaaa gtctacgcct gcgaagtcac ccatcagggc 660
 ctgagctcgc ccgtcacaaa gagcttcaac aggggagagt gt 702

<210> 12

<211> 234

<212> PRT

<213> Homo sapiens

<400> 12

Met Glu Ala Pro Ala Gln Leu Leu Phe Leu Leu Leu Trp Leu Pro
 1 5 10 15

Asp Thr Thr Gly Glu Ile Val Met Thr Gln Ser Pro Ala Thr Leu Ser
 20 25 30

Val Ser Pro Gly Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser
 35 40 45

Val Ser Ser Asn Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro
 50 55 60

Arg Leu Leu Ile Tyr Gly Ala Ser Thr Arg Ala Ser Gly Ile Pro Asp
 65 70 75 80

Arg Ile Ser Gly Ser Gly Ser Gly Thr Glu Phe Thr Leu Ile Ile Ser
 85 90 95

Ser Leu Gln Ser Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Ser Asn
 100 105 110

Asn Trp Pro Phe Thr Phe Gly Pro Gly Thr Lys Val Asp Ile Lys Arg
 115 120 125

Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln
 130 135 140

Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr
 145 150 155 160

Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser
 165 170 175

Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr
 180 185 190

Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys
 195 200 205

His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro
 210 215 220

Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
 225 230

<210> 13

<400> 13

000

<210> 14

<211> 460

<212> PRT

<213> Homo sapiens

<400> 14

Met Glu Phe Gly Leu Ser Trp Val Phe Leu Val Ala Ile Ile Lys Gly
 1 5 10 15

Val Gln Cys Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys
 20 25 30

Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
 35 40 45

Ser Asp Tyr Tyr Met Ser Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu
 50 55 60

Glu Trp Phe Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala
 65 70 75 80

Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
 85 90 95

Ser Leu Ser Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val
 100 105 110

Tyr Tyr Cys Ala Arg Gly Leu Thr Gly Asp Tyr Trp Gly Gln Gly Thr
 115 120 125

Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro
 130 135 140

Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly
 145 150 155 160

Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn
 165 170 175

Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln
 180 185 190

Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser
 195 200 205

Asn Phe Gly Thr Gln Thr Tyr Thr Cys Asn Val Asp His Lys Pro Ser
 210 215 220

Asn Thr Lys Val Asp Lys Thr Val Glu Arg Lys Cys Cys Val Glu Cys
 225 230 235 240

Pro	Pro	Cys	Pro	Ala 245	Pro	Pro	Val	Ala 250	Gly	Pro	Ser	Val	Phe 255	Leu	Phe
Pro	Pro	Lys	Pro 260	Lys	Asp	Thr	Leu	Met 265	Ile	Ser	Arg	Thr	Pro 270	Glu	Val
Thr	Cys	Val 275	Val	Val	Asp	Val	Ser 280	His	Glu	Asp	Pro	Glu 285	Val	Gln	Phe
Asn 290	Trp	Tyr	Val	Asp	Gly	Val 295	Glu	Val	His	Asn	Ala 300	Lys	Thr	Lys	Pro
Arg 305	Glu	Glu	Gln	Phe	Asn 310	Ser	Thr	Phe	Arg	Val 315	Val	Ser	Val	Leu	Thr 320
Val	Val	His	Gln	Asp 325	Trp	Leu	Asn	Gly	Lys 330	Glu	Tyr	Lys	Cys	Lys	Val
Ser	Asn	Lys	Gly 340	Leu	Pro	Ala	Pro 345	Ile	Glu	Lys	Thr	Ile	Ser 350	Lys	Thr
Lys	Gly	Gln 355	Pro	Arg	Glu	Pro	Gln 360	Val	Tyr	Thr	Leu	Pro 365	Pro	Ser	Arg
Glu 370	Glu	Met	Thr	Lys	Asn	Gln 375	Val	Ser	Leu	Thr	Cys 380	Leu	Val	Lys	Gly
Phe 385	Tyr	Pro	Ser	Asp 390	Ile	Ala	Val	Glu	Trp	Glu 395	Ser	Asn	Gly	Gln	Pro 400
Glu	Asn	Asn	Tyr 405	Lys	Thr	Thr	Pro	Pro 410	Met	Leu	Asp	Ser	Asp 415	Gly	Ser
Phe	Phe	Leu	Tyr 420	Ser	Lys	Leu	Thr	Val 425	Asp	Lys	Ser	Arg	Trp 430	Gln	Gln
Gly 435	Asn	Val	Phe	Ser	Cys	Ser	Val 440	Met	His	Glu	Ala	Leu 445	His	Asn	His
Tyr 450	Thr	Gln	Lys	Ser	Leu	Ser	Leu 455	Ser	Pro	Gly	Lys 460				

<210> 15

<400> 15
000

```
<210> 16
<211> 236
<212> PRT
<213> Homo sapiens
```

<400> 16
Met Asp Met Arg Val Pro Ala Gln Leu Leu Gly Leu Leu Leu Leu Trp

1		5		10		15									
Phe	Pro	Gly	Ser	Arg	Cys	Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser
		20						25					30		
Val	Ser	Ala	Ser	Val	Gly	Asp	Arg	Val	Thr	Ile	Ser	Cys	Arg	Ala	Ser
		35				40						45			
Gln	Asp	Ile	Ser	Gly	Trp	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys
	50					55					60				
Ala	Pro	Lys	Leu	Leu	Ile	Ser	Ala	Thr	Ser	Ser	Leu	His	Ser	Gly	Val
	65				70				75					80	
Pro	Ser	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr
			85						90					95	
Ile	Ser	Ser	Leu	Gln	Pro	Glu	Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln
			100					105					110		
Thr	Asn	Ser	Phe	Pro	Phe	Thr	Phe	Gly	Pro	Gly	Thr	Lys	Val	Asp	Ile
			115					120				125			
Lys	Arg	Thr	Val	Ala	Ala	Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	Ser	Asp
	130					135					140				
Glu	Gln	Leu	Lys	Ser	Gly	Thr	Ala	Ser	Val	Val	Cys	Leu	Leu	Asn	Asn
	145				150					155				160	
Phe	Tyr	Pro	Arg	Glu	Ala	Lys	Val	Gln	Trp	Lys	Val	Asp	Asn	Ala	Leu
				165					170					175	
Gln	Ser	Gly	Asn	Ser	Gln	Glu	Ser	Val	Thr	Glu	Gln	Asp	Ser	Lys	Asp
			180					185					190		
Ser	Thr	Tyr	Ser	Leu	Ser	Ser	Thr	Leu	Thr	Leu	Ser	Lys	Ala	Asp	Tyr
		195					200						205		
Glu	Lys	His	Lys	Val	Tyr	Ala	Cys	Glu	Val	Thr	His	Gln	Gly	Leu	Ser
	210					215					220				
Ser	Pro	Val	Thr	Lys	Ser	Phe	Asn	Arg	Gly	Glu	Cys				
	225				230					235					

<210> 17

<400> 17
000
<210> 18
<211> 463
<212> PRT
<213> Homo sapiens

<400> 18

Met Glu Phe Gly Leu Ser Trp Val Phe Leu Val Ala Leu Leu Arg Gly
 1 5 10 15

 Cys Gln Cys Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln
 20 25 30

 Pro Gly Arg Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
 35 40 45

 Ser Ser Tyr Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
 50 55 60

 Glu Trp Val Ala Phe Ile Trp Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala
 65 70 75 80

 Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn
 85 90 95

 Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val
 100 105 110

 Tyr Tyr Cys Ala Arg Gly Tyr Arg Val Tyr Phe Asp Tyr Trp Gly Gln
 115 120 125

 Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
 130 135 140

 Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala
 145 150 155 160

 Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
 165 170 175

 Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
 180 185 190

 Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
 195 200 205

 Ser Ser Ser Leu Gly Thr Lys Thr Tyr Thr Cys Asn Val Asp His Lys
 210 215 220

 Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Ser Lys Tyr Gly Pro
 225 230 235 240

 Pro Cys Pro Ser Cys Pro Ala Pro Glu Phe Leu Gly Gly Pro Ser Val
 245 250 255

 Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr
 260 265 270

 Pro Glu Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu
 275 280 285

 Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys

290 295 300
 Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser
 305 310 315 320
 Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys
 325 330 335
 Cys Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile
 340 345 350
 Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro
 355 360 365
 Pro Ser Gln Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu
 370 375 380
 Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn
 385 390 395 400
 Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser
 405 410 415
 Asp Gly Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser Arg
 420 425 430
 Trp Gln Glu Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu
 435 440 445
 His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 450 455 460

<210> 19

 <400> 19
 000

 <210> 20
 <211> 236
 <212> PRT
 <213> Homo sapiens

 <400> 20
 Met Asp Met Arg Val Pro Ala Gln Leu Leu Gly Leu Leu Leu Trp
 1 5 10 15
 Phe Pro Gly Ser Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser
 20 25 30
 Val Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser
 35 40 45
 Gln Asp Ile Ser Ser Trp Leu Ala Trp Tyr Gln Arg Lys Pro Gly Lys
 50 55 60

Ala Pro Lys Leu Gln Ile Tyr Ala Ala Ser Ser Leu Glu Ser Gly Val
65 70 75 80

Pro Ser Arg Phe Asn Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Ser
85 90 95

Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln
100 105 110

Thr Asn Ser Ser Phe Pro Leu Thr Phe Gly Gly Gly Thr Lys Val Glu Ile
115 120 125

Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp
130 135 140

Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn
145 150 155 160

Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu
165 170 175

Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp
180 185 190

Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr
195 200 205

Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser
210 215 220

Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
225 230 235

<210> 21

<400> 21

000

<210> 22

<211> 464

<212> PRT

<213> Homo sapiens

<400> 22

Met Glu Trp Thr Trp Ser Phe Leu Phe Leu Val Ala Ala Ala Thr Gly
1 5 10 15

Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys
20 25 30

Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe
35 40 45

Thr Ser Tyr Gly Ile Ser Trp Val Arg Gln Ala Pro Gly Gln Gly Leu

50	55	60
Glu Trp Met Gly Trp Ile Ser Ala Tyr Asn Gly Asn Thr Asn Tyr Ala		
65	70	75
80		
Gln Lys Leu Gln Asp Arg Val Thr Met Thr Thr Asp Thr Ser Thr Thr		
85	90	95
Thr Ala Tyr Met Glu Leu Arg Ser Leu Arg Ser Asp Asp Thr Ala Val		
100	105	110
Tyr Tyr Cys Ala Arg Arg Ala Tyr Gly Ala Asn Phe Phe Asp Tyr Trp		
115	120	125
Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro		
130	135	140
Ser Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr		
145	150	155
160		
Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr		
165	170	175
Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro		
180	185	190
Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr		
195	200	205
Val Pro Ser Ser Asn Phe Gly Thr Gln Thr Tyr Thr Cys Asn Val Asp		
210	215	220
His Lys Pro Ser Asn Thr Lys Val Asp Lys Thr Val Glu Arg Lys Cys		
225	230	235
240		
Cys Val Glu Cys Pro Pro Cys Pro Ala Pro Pro Val Ala Gly Pro Ser		
245	250	255
Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg		
260	265	270
Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro		
275	280	285
Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala		
290	295	300
Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Phe Arg Val Val		
305	310	315
320		
Ser Val Leu Thr Val Val His Gln Asp Trp Leu Asn Gly Lys Glu Tyr		
325	330	335
Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ala Pro Ile Glu Lys Thr		
340	345	350

Ile Ser Lys Thr Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu
 355 360 365

Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys
 370 375 380

Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser
 385 390 395 400

Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Met Leu Asp
 405 410 415

Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser
 420 425 430

Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala
 435 440 445

Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 450 455 460

<210> 23

<400> 23

000

<210> 24

<211> 240

<212> PRT

<213> Homo sapiens

<400> 24

Met Val Leu Gln Thr Gln Val Phe Ile Ser Leu Leu Leu Trp Ile Ser
 1 5 10 15

Gly Ala Tyr Gly Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala
 20 25 30

Val Ser Leu Gly Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Gln Ser
 35 40 45

Ile Leu Phe Phe Ser Asn Asn Lys Asn Tyr Leu Ala Trp Tyr Arg Gln
 50 55 60

Lys Pro Gly Gln Pro Pro Asn Leu Leu Ile Tyr Trp Ala Ser Thr Arg
 65 70 75 80

Glu Ser Gly Val Pro Asp Arg Phe Ser Gly Ser Gly Thr Asp
 85 90 95

Phe Thr Leu Thr Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr
 100 105 110

Tyr Cys Gln Gln Tyr Tyr Ser Ser Pro Trp Thr Phe Gly Gln Gly Thr
 115 120 125
 Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe
 130 135 140
 Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys
 145 150 155 160
 Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val
 165 170 175
 Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln
 180 185 190
 Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser
 195 200 205
 Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His
 210 215 220
 Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
 225 230 235 240

<210> 25

<211> 1380

<212> DNA

<213> Homo sapiens

<400> 25

atggagtttg ggctgagctg ggttttccct gttgctatta taaaagggtg ccagtgctcag 60
 gtgcagctgg tggagtctgg gggaggcttg gtcaagcctg gagggtccct gagactctcc 120
 tgtgcagcct ctggattcac ctccagtgac tactatatga gctggatccg ccaggctcca 180
 gggaaaggac tggagtgggt ttcatacatt agtagtagtg gtagtaccat atactagca 240
 gactctgtga agggccgatt caccatctcc agggacaacg ccaagaactc actgtatctg 300
 caaatgaaca gcttgagagc cgaggacacg gcctgtatt actgtgcgag aggcctaact 360
 ggggactact ggggccaggg aacctcggtc acgctctcct cagcttccac caagggccca 420
 tcctctctcc ccttgccgcc ctgctctaga agcacctccg agagcacagc ggcctctggc 480
 tgctctggta aggactactt cccogaacgg gtgacggtgt cgtggaactc aggcctctcg 540
 accagcggcg tgcacacctt cccagctgtc ctacagtcct caggactcta ctccctcagc 600
 agcgtgggta cgtgcccctc cagcaacttc ggcacccaga cctacacctc caacgtagat 660
 cacaagccca gcaacaccaa ggtggacaag acagttgagc gcaaatgttg tgtcgagtg 720
 ccaccgtgct cagcaccacc tgtggcagga ccgtcagttc tcctctctcc cccaaaacc 780
 aaggacaccc tcatgatctc cggaccctct gaggtcacgt gcgtgggtgg ggcagtgagc 840
 ccaagaagcc ccagaggtcca gttcaactgg tacgtggagc gcgtggaggt gcatatgccc 900
 aagacaaagg cacgggagga gcagttcaac agcacgttcc gtgtggtcag cgtctccacc 960
 tgtgtgcaac aggaactggct gaacggcaag gactacaagt gcaaggctcc caacaaaggc 1020
 ctcccagccc ccactgagaa aaccatctcc aaaaccaaag ggcagccccc agaaccacag 1080
 gtgtacaccc tgcccccatc cggggaggag atgaccaaga accaggtcag cctgacctgc 1140
 ctggtcaaaag gctcttacc cagcgacatc gcctggaggt gggagagcaa tggcgacccg 1200
 gagaacaact acaagaccac acctcccatg ctggactccg accgctcctt ctctctctac 1260

agcaagctca ccgtggacaa gagcaggtgg cagcagggga acgtcttctc atgctccgtg 1320
 atgcatgagg ctctgcacaa ccactacacg cagaagagcc tctccctgtc tccgggtaaa 1380

<210> 26

<211> 460

<212> PRT

<213> Homo sapiens

<400> 26

Met Glu Phe Gly Leu Ser Trp Val Phe Leu Val Ala Ile Ile Lys Gly
 1 5 10 15
 Val Gln Cys Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys
 20 25 30
 Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
 35 40 45
 Ser Asp Tyr Tyr Met Ser Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu
 50 55 60
 Glu Trp Val Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala
 65 70 75 80
 Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
 85 90 95
 Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val
 100 105 110
 Tyr Tyr Cys Ala Arg Gly Leu Thr Gly Asp Tyr Trp Gly Gln Gly Thr
 115 120 125
 Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro
 130 135 140
 Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly
 145 150 155 160
 Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn
 165 170 175
 Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln
 180 185 190
 Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser
 195 200 205
 Asn Phe Gly Thr Gln Thr Tyr Thr Cys Asn Val Asp His Lys Pro Ser
 210 215 220
 Asn Thr Lys Val Asp Lys Thr Val Glu Arg Lys Cys Cys Val Glu Cys
 225 230 235 240
 Pro Pro Cys Pro Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu Phe

[illegible]

<210> 27

<211> 708

<212> DNA

<213> Homo sapiens

<400> 27

ttgacgatcat	gggttccegc	tcagctctctc	gggtctctctc	tactcttgct	ccgaggtgac	60
agatgtgaca	tcagatgac	tcagctctcca	tctctctctc	ctgcatctct	cgagacagac	120
ctccaccatc	ctgtccggcc	aagtgcagtc	atctagcagt	tattaaatgt	gtatcagcac	180
aaacacaggga	aagccctctaa	ctctctgtatc	catgtctgac	ccagtttgta	aagttggggtc	240
ccatcaaggt	tcagtggcag	tggatctgtg	acagatttca	ctctcaccat	cattagttctc	300
caactcgta	attttccaac	tactactgt	acacagagtt	acagatcccc	actattcttc	360
cgccctggga	cccaaaagt	tatcaaacga	actgctgtgt	cacacctctc	ctctcatctc	420
ggcccatctg	atgacagtt	gaattctgga	actcgtcttc	tgtgttgtct	ctctaaatac	480


```

ttctatccca gagaggccaa agtacagtgg aaggtggata acgccctcca atcgggtaac 540
tcccaggaga gtgtcacaga gcaggacagc aaggacagca cctacagcct cagcagcacc 600
ctgacgctga gcaaagcaga ctacgagaaa cacaaagtct acgctgcga agtcacccat 660
cagggcctga gctcgccgt cacaagagc ttcaacaggg gagagtgt 708

```

<210> 28

<211> 236

<212> PRT

<213> Homo sapiens

<400> 28

```

Met Asp Met Arg Val Pro Ala Gln Leu Leu Gly Leu Leu Leu Trp
  1              5              10              15

Leu Arg Gly Ala Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser
      20              25              30

Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Pro Ser
      35              40              45

Gln Ile Ile Ser Ser Leu Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys
      50              55              60

Ala Pro Lys Leu Leu Ile His Ala Ala Ser Ser Leu Gln Ser Gly Val
      65              70              75              80

Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
      85              90              95

Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln
      100             105             110

Ser Tyr Ser Thr Pro Phe Thr Phe Gly Pro Gly Thr Lys Val Asp Ile
      115             120             125

Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp
      130             135             140

Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn
      145             150             155             160

Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu
      165             170             175

Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp
      180             185             190

Ser Thr Tyr Ser Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr
      195             200             205

Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser
      210             215             220

Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
      225             230             235

```

<210> 29
 <211> 1398
 <212> DNA
 <213> Homo sapiens

<400> 29
 atggagttgg ggctgtgtg ggttttcctt gttgctattt tagaaggtgt ccagtgtgag 60
 gtgcagctgg tggagttcgg gggaggcttg gtacagcctg ggggggtccct gagactctcc 120
 tgtgcagcct ctggattcac ctccagtagt tttagataga cctgggtccg ccaggctcca 180
 ggaagggggc tggagttggg ttcatcatt agtagtagaa gtatgacct atctcatgca 240
 gactctgtga agggccgatt caccatctcc agagacaatg ccaagaactc actgtatctg 300
 caaatgaaca gctcgagaga cagggacacg gctgtgtatt actgtgcgag agatcctctt 360
 ctagcgggag ctaccttctt tgactactgg ggcaggga ccttggtcac cgtctctcca 420
 gcctccacca agggcccatc ggtcttcccc ctggcgccct gctccaggag cactcccgag 480
 agcacagcgg cctcgggctg cctgggtcaag gactacttcc ccgaaccggt gacggtgtcg 540
 ttggaactcag gcctctgtac cagcggcgtg cacaccttcc cagctgtctt acagctctca 600
 ggactctact cctcagcag cgtggtgacc gtgcccctcca gcaacttcgg caccagacc 660
 tacactctga acgtagatca caagcccagc aacaccaagg tggacaagac agttgagcgc 720
 aaatggttgg tcgagtgccc accgtgcccc gcaccacctg tggcaggacc gtcagtcttc 780
 ctcttcccc caaaaccaca ggacacctc atgatctccc ggacccctga ggtcacgtgc 840
 gtgggtgggg acgtgagcca cgaagacccc gagggtccagt tcaactggta cgtggacgag 900
 ctggagggtg ataattgccaa gacaaagcca cgggaggagc agttcaacag cagttccgt 960
 gtggtcagcg tctcacaggt tgtgcaccag gactggctga acggcaagga gtacaagtgc 1020
 aaggtctcca acaaggcctt cccagcccc atcagagaaa ccatctccaa aaccacaggg 1080
 cagccccag aaccacaggt gtacacctg ccccatccc gggaggagat cgaacgaac 1140
 caggtcagcc tgacctgcct ggtcaaaggc ttctaccaca cgcacatcgc cgtggaggtg 1200
 gagagcaatg gccagccgga gaacaaacta aagaccacac ctcccattgc ggactccgac 1260
 ggctctctat tctctacag caagctcacc gtggacaaga gcaggtggga gcaggggna 1320
 gtcttctcat gctccgtgat gcatgaggct ctgcacaacc actacacga gaagagcctc 1380
 tccctgtctc cgggtaaa 1398

<210> 30
 <211> 466
 <212> PRT
 <213> Homo sapiens

<400> 30
 Met Glu Leu Gly Leu Cys Trp Val Phe Leu Val Ala Ile Leu Glu Gly
 1 5 10 15
 Val Gln Cys Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln
 20 25 30
 Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
 35 40 45
 Ser Ser Phe Ser Met Thr Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
 50 55 60
 Glu Trp Val Ser Tyr Ile Ser Ser Arg Ser Ser Thr Ile Ser Tyr Ala
 65 70 75 80
 Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
 85 90 95

Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Asp Glu Asp Thr Ala Val
 100 105 110
 Tyr Tyr Cys Ala Arg Asp Pro Leu Leu Ala Gly Ala Thr Phe Phe Asp
 115 120 125
 Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys
 130 135 140
 Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu
 145 150 155 160
 Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro
 165 170 175
 Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr
 180 185 190
 Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val
 195 200 205
 Val Thr Val Pro Ser Ser Asn Phe Gly Thr Gln Thr Tyr Thr Cys Asn
 210 215 220
 Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys Thr Val Glu Arg
 225 230 235 240
 Lys Cys Cys Val Glu Cys Pro Pro Cys Pro Ala Pro Pro Val Ala Gly
 245 250 255
 Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
 260 265 270
 Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
 275 280 285
 Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
 290 295 300
 Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Phe Arg
 305 310 315 320
 Val Val Ser Val Leu Thr Val Val His Gln Asp Trp Leu Asn Gly Lys
 325 330 335
 Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ala Pro Ile Glu
 340 345 350
 Lys Thr Ile Ser Lys Thr Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
 355 360 365
 Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
 370 375 380
 Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
 385 390 395 400

Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Met
 405 410 415

Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
 420 425 430

Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
 435 440 445

Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
 450 455 460

Gly Lys
 465

<210> 31
 <211> 705
 <212> DNA
 <213> Homo sapiens

<400> 31
 atggaaaccc cagcgagct tctcttctc ctgctactct ggctccaga taccaccgga 60
 gaatttggtg tgacgcagtc tccaggcacc ctgtctttgt ctccagggga aagagccacc 120
 ctctctctga gggccagtc gagtgtagc agcagttact tagcctggta ccagcagaaa 180
 cctggccagg ctccaggct cctcatctat ggtgcatcca gcagggccac tggcatccca 240
 gacaggttca gtggcagtg gtctgggaca gacttcaact taccatcag cagactggag 300
 cctgaagatt ttgcagtgt ttactgtcag cagtatggta gctcacctct caotttcggc 360
 ggagggaacca aggtggagat caaacgaact gtggctgcac catctgtctt catcttcccg 420
 ccatctgatg agcagttgaa atctggaact gcctctgttg tgtgcctgct gaataacttc 480
 tatccagag aggccaaagt acagtgggaag gtggataacg cctcccaatc gggttaactcc 540
 caggagagtg tcacagagca ggcagcaag gacagcaact acagcctcag cagcaccctg 600
 acgctgagca aagcagacta cgagaaacac aaagtctacg cctgcgaagt caccatcag 660
 ggcctgagct cgcccgctac aaagagcttc aacaggggag agtgt 705

<210> 32
 <211> 235
 <212> PRT
 <213> Homo sapiens

<400> 32
 Met Glu Thr Pro Ala Gln Leu Leu Phe Leu Leu Leu Trp Leu Pro
 1 5 10 15

Asp Thr Thr Gly Glu Phe Val Leu Thr Gln Ser Pro Gly Thr Leu Ser
 20 25 30

Leu Ser Pro Gly Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser
 35 40 45

Val Ser Ser Ser Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala
 50 55 60

Pro Arg Leu Leu Ile Tyr Gly Ala Ser Ser Arg Ala Thr Gly Ile Pro
 65 70 75 80

Asp Arg Phe Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile
85 95

Ser Arg Leu Glu Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Tyr
100 105 110

Gly Ser Ser Pro Leu Thr Phe Gly Gly Thr Lys Val Glu Ile Lys
115 120 125

Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu
130 135 140

Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe
145 150 155 160

Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln
165 170 175

Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser
180 185 190

Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
195 200 205

Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser
210 215 220

Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
225 230 235

<210> 33

<211> 1383

<212> DNA

<213> Homo sapiens

<400> 33

atggagtttg ggcgtgagctg ggttttcoctt gttgctatta taaaagggtgt ccagtgctcag 60
gtgcagctgg tggagctctgg gggaggcttg gtcaagcctg gagggtccct gagactctcc 120
tgtgcagcct ctggattccac ctccagtgac tactacatga gctggatccg ccaggctcca 180
gggaaggggg tggagtggtt ttcatacatt agtagtagtg gtagtaccat atactacgca 240
gactctgtga agggccgatt caccatctcc agggacaacg ccaagaattc actgtatctg 300
caaatgaaca gcctgagagc cgaggacacg gccgtgtatt actgtgcgag gcgtatagga 360
ggatgggacg tctggggcca agggaccacg gtccacgtct cctcagcttc caccaaggcg 420
ccatccgtct tccccctggc gccctgctct agaagcacct ccgagagcac agcggccctg 480
ggctgctctg tcaaggacta ctccccgaa ccggtgacgg tgtcgtggaa ctcaggcgct 540
ctgaccagcg gcgtgcacac ctccccagct gtccacagct cctcaggact ctactccctc 600
agcagcgtgg tgaccgtgcc ctccagcaac ttccggcacc agacctacac ctgcaacgta 660
gatcacaagc ccagcaaac caaggtggag aagcagcttg agcgcaaatg ttgtgtcgag 720
tgcccaccgt gccacgaccc acctgtggca ggacogtccg tcttctctct ccccccaaaa 780
cccgaaggaca cctcatgat ctccccgacc cctgaggtca cgtgcgtggg ggtggacgtg 840
agccacgaag accccagagt ccagttcaac tggtagctgg acggcgtgga ggtgcataat 900
gccaagacaa agccacggga ggagcagttc aacagcacgt tccgtgtggg cagcgtcctc 960
accgtgtgac accaggaact gctgaacggc aaggagtaca agtgcgaagt ctccaaacaa 1020
ggctcccgac ccccatcga gaaaaccatc tccaaaacca aagggcagcc ccgagaacaa 1080
caggtgtaca cctgcctccc atccccggag gagatgacca agaaccaggt cagcctgacc 1140

```

tgcttggtca aagcttcta cccagcgac atcgccgtgg agtgggagag caatgggcag 1200
cgggagaaca actacaagac cacacctccc atgctggact cgaaggctc cttcttcttc 1260
tacagcaagc tcaccgtgga caagagcagg tggcagcagg ggaacgtctt ctcattgtcc 1320
gtgatgcatt aggccttgca caaccatac acgcagaaga gcctctccct gtctccgggt 1380
aaa
1383

```

<210> 34

<211> 461

<212> PRT

<213> Homo sapiens

<400> 34

```

Met Glu Phe Gly Leu Ser Trp Val Phe Leu Val Ala Ile Ile Lys Gly
  1             5             10             15

Val Gln Cys Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys
          20             25             30

Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
      35             40             45

Ser Asp Tyr Tyr Met Ser Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu
      50             55             60

Glu Trp Val Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala
      65             70             75             80

Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
          85             90             95

Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val
      100             105             110

Tyr Tyr Cys Ala Arg Arg Ile Gly Gly Met Asp Val Trp Gly Gln Gly
      115             120             125

Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe
      130             135             140

Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu
      145             150             155             160

Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp
      165             170             175

Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu
      180             185             190

Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser
      195             200             205

Ser Asn Phe Gly Thr Gln Thr Tyr Thr Cys Asn Val Asp His Lys Pro
      210             215             220

Ser Asn Thr Lys Val Asp Lys Thr Val Glu Arg Lys Cys Cys Val Glu
      225             230             235             240

```

Cys Pro Pro Cys Pro Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu
 245 250 255
 Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu
 260 265 270
 Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Gln
 275 280 285
 Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys
 290 295 300
 Pro Arg Glu Glu Gln Phe Asn Ser Thr Phe Arg Val Val Ser Val Leu
 305 310 315 320
 Thr Val Val His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys
 325 330 335
 Val Ser Asn Lys Gly Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys
 340 345 350
 Thr Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser
 355 360 365
 Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys
 370 375 380
 Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln
 385 390 395 400
 Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Met Leu Asp Ser Asp Gly
 405 410 415
 Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln
 420 425 430
 Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn
 435 440 445
 His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 450 455 460
 <210> 35
 <211> 708
 <212> DNA
 <213> Homo sapiens
 <400> 35
 atggacatga ggggtcccccgc tcagctctctg gggctctctgc tactctggct ccgaggtgcc 60
 agatgtgaca tccagatgac ccagctctcca tctcctctgt ctgcattctgt aggagacaga 120
 gtccacatca cttgcccgggc aagtcagagc attagcggct ttttaatttg gtatcagcag 180
 agaccaggga aagcccccctaa gctctctgac tatgctacat ccagttttaca aagtgggggc 240
 ccatcaaggt tcagtgccagc tggatctctgg acagatttca ctctcaccat cagcagctctg 300
 caacctgaag attttgcaac ttactactgt caacagagtt acagtacccc attcactttc 360
 ggccctggga ccaaagtga tatcaaacga actgtggctg caccatctgt cttcatcttc 420

```

ccgccatctg atgagcagtt gaaatctgga actgectctg ttgtgtgcct gctgaataac 480
ttctatccca gagaggccaa agtacagtgg aagggtggata acgccctcca atcggtgaac 540
tcccaggaga gtgtcacaga gcaggacagc aaggacagca cctacagcct cagcagcacc 600
ctgacgctga gcaaagcaga ctacgagaaa cacaaggtct acgectgcga agtcacccat 660
cagggcctga gctcgccgct cacaagagac ttcaacaggg gagagtgt 708

```

<210> 36

<211> 236

<212> PRT

<213> Homo sapiens

<400> 36

```

Met Asp Met Arg Val Pro Ala Gln Leu Leu Gly Leu Leu Leu Trp
  1                      5                      10                      15

```

```

Leu Arg Gly Ala Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser
      20                      25                      30

```

```

Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser
      35                      40                      45

```

```

Gln Ser Ile Ser Gly Phe Leu Ile Trp Tyr Gln Gln Arg Pro Gly Lys
      50                      55                      60

```

```

Ala Pro Lys Leu Leu Ile Tyr Ala Thr Ser Ser Leu Gln Ser Gly Val
      65                      70                      75                      80

```

```

Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
      85                      90                      95

```

```

Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln
      100                      105                      110

```

```

Ser Tyr Ser Thr Pro Phe Thr Phe Gly Pro Gly Thr Lys Val Asp Ile
      115                      120                      125

```

```

Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp
      130                      135                      140

```

```

Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn
      145                      150                      155                      160

```

```

Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu
      165                      170                      175

```

```

Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp
      180                      185                      190

```

```

Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr
      195                      200                      205

```

```

Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser
      210                      215                      220

```

```

Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
      225                      230                      235

```


<210> 37
 <211> 1383
 <212> DNA
 <213> Homo sapiens

<400> 37
 atggagtttg ggtgagctg ggttttcctt gttgctatta taaaagggtt ccagtgtcag 60
 gtgcagctgg tggagctctg gggaggcttg gtcaagctg gagggtccct gagactctcc 120
 tgtgcagcct ctggaattcac ctccagtgac tactatatga gctggatccg ccaggctcca 180
 gggaaagggac tggagtggtt ttcatacatt agtagtagtg gtatgacctat atactacgca 240
 gactctgtga agggcgagat caccatctcc agggacaacg ccaagaactc actgtatctg 300
 caaatgaaca gcctgagagc cgaggacacg gccgtgtatt actgtgcgag aggccttaact 360
 ggggactact gggggcaggg aacctgtgtc accgtctcct cagcttccac caagggccca 420
 tcctgtcttc ccctggcgcc ctgctctaga agcacctccg agagcacagc ggccctgggc 480
 tgctctgtca aggactactt cccgaacggc gtgacggtgt cgtggaactc aggcgctctg 540
 accagcggtg tgcacacctt cccagctgtc ctacagtctt caggactcta ctccctcagc 600
 agcgtggtga ccgtgcccct cagcagcttg ggcacgaaga cctacacctg caacgtagat 660
 cacaagccca gcaacaccaa ggtggacaag agagttgagt ccaaatatgg tccccatgc 720
 ccatcatgoc cagcacctga gtctctgggg ggaccatcag tcttccgtgt cccccaaaa 780
 cccaaggaca ctctcatgat ctcccgagcc cctgaggtca cgtgcgtgtt ggtggaactg 840
 agccaggaaag accccgaggt ccagttcaac tggtagcttg atggcgtgga ggtgcataat 900
 gccaaagaca agcccgcgga ggagcagttc aacagcacgt accgtgtggt cagcgtctct 960
 accgtctctc accaggactg gctgaacggc aaggagtaca agtgcaaggt ctccaacaaa 1020
 ggctctccgt cctccatcga gaaaaccatc tccaaagcca aagggcagcc ccgagagcca 1080
 caggtgtaca cctgtccccc atcccaggag gagatgacca agaaccaggt cagcgtgacc 1140
 tgctctgtca aaggcttcta cccacgcgac atgcgcgtgg agtgggagag caatgggacg 1200
 ccggagaaca actacaagac cagcctctcc gtgctggact ccgacggctc ctctctctcc 1260
 tacagcaggc taaccgtgga caagagcagg tggcaggagg ggaatgtctt ctcatgtccc 1320
 gtgatgcagc aggcctctga caaccactac acacagaaga gcctctccct gtctccgggt 1380
 aaa 1383

<210> 38
 <211> 461
 <212> FRT
 <213> Homo sapiens

<400> 38
 Met Glu Phe Gly Leu Ser Trp Val Phe Leu Val Ala Ile Ile Lys Gly
 1 5 10 15
 Val Gln Cys Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys
 20 25 30
 Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
 35 40 45
 Ser Asp Tyr Tyr Met Ser Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu
 50 55 60
 Glu Trp Val Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala
 65 70 75 80
 Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
 85 90 95

Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val
 100 105 110
 Tyr Tyr Cys Ala Arg Gly Leu Thr Gly Asp Tyr Trp Gly Gln Gly Thr
 115 120 125
 Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro
 130 135 140
 Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly
 145 150 155 160
 Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn
 165 170 175
 Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln
 180 185 190
 Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser
 195 200 205
 Ser Leu Gly Thr Lys Thr Tyr Thr Cys Asn Val Asp His Lys Pro Ser
 210 215 220
 Asn Thr Lys Val Asp Lys Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys
 225 230 235 240
 Pro Ser Cys Pro Ala Pro Glu Phe Leu Gly Gly Pro Ser Val Phe Leu
 245 250 255
 Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu
 260 265 270
 Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln
 275 280 285
 Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys
 290 295 300
 Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu
 305 310 315 320
 Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys
 325 330 335
 Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys
 340 345 350
 Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser
 355 360 365
 Gln Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys
 370 375 380
 Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln

385		390		395		400
Pro Glu Asn Asn Tyr Lys Thr Thr		Pro Pro Val Leu Asp Ser Asp Gly				
	405		410		415	
Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser Arg Trp Gln						
	420		425		430	
Glu Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn						
	435		440		445	
His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys						
	450		455		460	

<210> 39

<400> 39

000

<210> 40

<400> 40

000

<210> 41

<400> 41

000

<210> 42

<400> 42

000

<210> 43

<211> 705

<212> DNA

<213> Homo sapiens

<400> 43

atggaaaccc	cagcgcagct	tctcttctct	ctgctactct	ggctcccaga	taccaccgga	60
gaattttgtgt	tgacgcagtc	tccaggccac	ctgtctttgt	ctccaggggga	aagagccacc	120
ctctctctgca	gggccagtc	gagtggttagc	agcagttact	tagcctggta	ccagcagaaa	180
cctggccagg	ctccaggct	cctcatctat	gggtgatcca	gcagggccac	tggcatccca	240
gacaggttca	gtggcagtg	gtctgggaca	gacttcactc	tcaccatcag	cagactggag	300
cctgaagatt	ttgtagtgt	ttactgtcag	cagtatggta	getcacctct	cactttccgc	360
ggagggacca	agggtggag	caaacgaact	gtggctgcac	catctgtctt	catcttcccg	420
ccatctgatg	agcagttgaa	atctgggaact	gcctctgttg	tgtgcctgct	gaataacttc	480
tatcccagag	aggccaaagt	acagtgggaag	gtggataacg	ccctccaatc	gggtaactcc	540
caggagagtg	tcacagagca	ggacagcaag	gacagcacct	acagcctcag	cagcacccctg	600
acgttgagca	aagcagacta	cgagaaacac	aaagtctacg	cctgcgaagt	caccatcag	660

ggcctgagct cgcccgtcac aaagagcttc aacaggggag agtgt

705

<210> 44

<211> 235

<212> PRT

<213> Homo sapiens

<400> 44

Met Glu Thr Pro Ala Gln Leu Leu Phe Leu Leu Leu Trp Leu Pro
 1 5 10 15

Asp Thr Thr Gly Glu Phe Val Leu Thr Gln Ser Pro Gly Thr Leu Ser
 20 25 30

Leu Ser Pro Gly Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser
 35 40 45

Val Ser Ser Ser Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala
 50 55 60

Pro Arg Leu Leu Ile Tyr Gly Ala Ser Ser Arg Ala Thr Gly Ile Pro
 65 70 75 80

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile
 85 90 95

Ser Arg Leu Glu Pro Glu Asp Phe Val Val Tyr Tyr Cys Gln Gln Tyr
 100 105 110

Gly Ser Ser Pro Leu Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
 115 120 125

Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu
 130 135 140

Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe
 145 150 155 160

Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln
 165 170 175

Ser Gly Asn Ser Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser
 180 185 190

Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
 195 200 205

Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser
 210 215 220

Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
 225 230 235

<210> 45

<211> 1386

<212> DNA

<213> Homo sapiens

<400> 45

```

atggagtttg ggctgagctg ggttttccct gttgctatta taaaagggtt ccagtgtcag 60
gtgcagctgg tggagttctgg gggaggtctg gtcaagcctg gaggtccctc gagactctcc 120
tgtgcagcct ctggatttccac ctccagtgac tactacatga gctggatccg ccagggtccca 180
gggaaggggg tggagtgsgt ttccatacatt agtagtagtg gtagtaccat atactacgca 240
gactctgtga agggccgalt caccatctcc agggacaacg ccaagaatcc actgtatctg 300
caaatgaaca gcctgagagc cgaggacacg gccgtgtatt actgtggcag gcgtatagga 360
ggtatggagc tctggggcca agggaccacg gtccacgtct cctcagcttc caccgaaggc 420
ccatccgtct tccccctggc gccctgtctc agaagcacct ccgagagcac agcggccctg 480
ggctgcctgg tcaaggacta cttccccgaa ccggtgacgg tgtcgtggaa ctcaggcgct 540
ctgaccagcg gcgtgcacac cttcccagct gtccctacagt cctcaggact ctactccctc 600
agcagcgtgg tgaccgtgcc ctccagcagc ttgggcacga agacctacac ctgcaacgta 660
gatcacaaag ccagcaaacac caaggtggac aagagagttg agtccaaata tgggtcccca 720
tgccccatcat gccacgacac tgagttcctg gggggaccat cagtcttctt gttcccccca 780
aaacccaagg acactctcat gatctccogg acccctgagg tcacgtgcgt ggtggtggac 840
gtgagccagg aagaccocga ggtccagttc aactggtagc tggatggcgt ggaggtgcat 900
aatgccaaag caaagcccg cgaggagcag ttcaacacga cgtacogtgt ggtcagcgct 960
ctcaccgtcc tgaccacgga ctggctgaac ggcaaggagt acaagtgcga ggtctccacc 1020
aaagcgctcc cgtctccat cgagaaaaac atctccaaag ccaaagggca gccccgagag 1080
ccacaggtgt acacccctgcc cccatcccag gaggagatga ccaagaacca ggtcagcctg 1140
acctgcctgg tcaaaaggctt ctaccccagc gacatcgccg tggagtggga gagcaatggg 1200
cagccgggag acaactacaa gaccacgct cccgtgctgg actccgacg ctctctcttc 1260
ctctacagca ggctaaccgt ggacaagagc aggtggcagg aggggaatgt ttctctatgc 1320
tccgtgatgc atgaggtctc gcacaaccac tacacacaga agagcctctc cctgtctccg 1380
ggtaaa 1386

```

<210> 46

<211> 462

<212> PRT

<213> Homo sapiens

<400> 46

```

Met Gly Leu Phe Gly Leu Ser Trp Val Phe Leu Val Ala Ile Ile Lys Gly
  1             5             10             15

Val Gln Cys Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys
  20             25             30

Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
  35             40             45

Ser Asp Tyr Tyr Met Ser Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu
  50             55             60

Glu Trp Val Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala
  65             70             75             80

Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
  85             90             95

Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val
  100            105            110

```

Tyr Tyr Cys Ala Arg Arg Ile Gly Gly Met Asp Val Trp Gly Gln Gly
 115 120 125
 Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe
 130 135 140
 Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu
 145 150 155 160
 Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp
 165 170 175
 Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu
 180 185 190
 Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser
 195 200 205
 Ser Ser Leu Gly Thr Lys Thr Tyr Thr Cys Asn Val Asp His Lys Pro
 210 215 220
 Ser Asn Thr Lys Val Asp Lys Arg Val Glu Ser Lys Tyr Gly Pro Pro
 225 230 235 240
 Cys Pro Ser Cys Pro Ala Pro Glu Phe Leu Gly Gly Pro Ser Val Phe
 245 250 255
 Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro
 260 265 270
 Glu Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val
 275 280 285
 Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr
 290 295 300
 Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val
 305 310 315 320
 Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys
 325 330 335
 Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser
 340 345 350
 Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro
 355 360 365
 Ser Gln Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val
 370 375 380
 Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly
 385 390 395 400
 Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp
 405 410 415

Gly Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser Arg Trp
420 425 430

Gln Glu Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His
435 440 445

Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
450 455 460

<210> 47

<211> 708

<212> DNA

<213> Homo sapiens

<400> 47

atggacatga gggccccgc tcagctcctg gggctcctgc tactctggct ccgaggtgcc 60
agatgtgaca tccagatgac ccagctctoca tctccctgt ctgcattctgt aggagacaga 120
gtcaccatca cttgccgggc aagtcagagc attagcggct ttttaatttg gtatcagcag 180
agaccaggga aagccccata gctcctgac tatgctacat ccagtttaca aagtggggtc 240
ccattaaagt tcagtgccag tgaatctggg acagatttca ctctcaccat cagcagctgt 300
caacctgaag attttgcaac ttactactgt caacagagtt acagtacccc attcaatttc 360
ggccctggga ccaagtgga tatcaaacga actgtggctg caccattctgt ctcatcttc 420
cgcctatctg atgagcagtt gaaatctgga actgcctctg ttgtgtgct gtgtaataac 480
ttctatccca gagaggccaa agtcagtgga aagtggtgata acgcccctca atcggtgaac 540
tcccaggaga gtgtcacaga gcaggacagc aaggacagca cctacagcct cagcagcacc 600
ctgacagctg gcaaacgaga ctacgagaaa cacaagctct acgctgcca agtcacccat 660
cagggcctga gctcgccctg cacaagagag ttcaacaggg gagagtggt 708

<210> 48

<211> 236

<212> PRT

<213> Homo sapiens

<400> 48

Met Asp Met Arg Val Pro Ala Gln Leu Leu Gly Leu Leu Leu Trp
1 5 10 15

Leu Arg Gly Ala Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser
20 25 30

Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser
35 40 45

Gln Ser Ile Ser Gly Phe Leu Ile Trp Tyr Gln Gln Arg Pro Gly Lys
50 55 60

Ala Pro Lys Leu Leu Ile Tyr Ala Thr Ser Ser Leu Gln Ser Gly Val
65 70 75 80

Pro Leu Arg Phe Ser Gly Ser Glu Ser Gly Thr Asp Phe Thr Leu Thr
85 90 95

Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln

100 105 110
 Ser Tyr Ser Thr Pro Phe Thr Phe Gly Pro Gly Thr Lys Val Asp Ile
 115 120 125
 Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp
 130 135 140
 Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn
 145 150 155 160
 Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu
 165 170 175
 Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp
 180 185 190
 Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr
 195 200 205
 Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser
 210 215 220
 Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
 225 230 235

<210> 49

 <400> 49
 000

<210> 50

<211> 462

<212> PRT

<213> Homo sapiens

<400> 50

Met Glu Phe Gly Leu Ser Trp Val Phe Leu Val Ala Ile Ile Lys Gly
 1 5 10 15
 Val Gln Cys Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys
 20 25 30
 Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
 35 40 45
 Ser Asp Tyr Tyr Met Ser Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu
 50 55 60
 Glu Trp Val Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala
 65 70 75 80
 Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
 85 90 95

Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val
 100 105
 Tyr Tyr Cys Ala Ile Arg Ile Gly Gly Met Asp Val Trp Gly Gln Gly
 115 120 125
 Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe
 130 135 140
 Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu
 145 150 155 160
 Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp
 165 170 175
 Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu
 180 185 190
 Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser
 195 200 205
 Ser Ser Leu Gly Thr Lys Thr Tyr Thr Cys Asn Val Asp His Lys Pro
 210 215 220
 Ser Asn Thr Lys Val Asp Lys Arg Val Glu Ser Lys Tyr Gly Pro Pro
 225 230 235 240
 Cys Pro Pro Cys Pro Ala Pro Glu Phe Leu Gly Gly Pro Ser Val Phe
 245 250 255
 Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro
 260 265 270
 Glu Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val
 275 280 285
 Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr
 290 295 300
 Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val
 305 310 315 320
 Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys
 325 330 335
 Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser
 340 345 350
 Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro
 355 360 365
 Ser Gln Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val
 370 375 380
 Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly
 385 390 395 400

Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp
405 410 415

Gly Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser Arg Trp
420 425 430

Gln Glu Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His
435 440 445

Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
450 455 460

<210> 51

<400> 51
000

<210> 52

<211> 236

<212> PRT

<213> Homo sapiens

<400> 52

Met Asp Met Arg Val Pro Ala Gln Leu Leu Gly Leu Leu Leu Trp
1 5 10 15

Leu Arg Gly Ala Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser
20 25 30

Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser
35 40 45

Gln Ser Ile Ser Gly Phe Leu Ile Trp Tyr Gln Gln Lys Pro Gly Lys
50 55 60

Ala Pro Lys Leu Leu Ile Tyr Ala Thr Ser Ser Leu Gln Ser Gly Val
65 70 75 80

Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
85 90 95

Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln
100 105 110

Ser Tyr Ser Thr Pro Phe Thr Phe Gly Pro Gly Thr Lys Val Asp Ile
115 120 125

Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp
130 135 140

Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn
145 150 155 160

Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu

165										170					175					
Gln	Ser	Gly	Asn	Ser	Gln	Glu	Ser	Val	Thr	Glu	Gln	Asp	Ser	Lys	Asp					
180										185					190					
Ser	Thr	Tyr	Ser	Leu	Ser	Ser	Thr	Leu	Thr	Leu	Ser	Lys	Ala	Asp	Tyr					
195										200					205					
Glu	Lys	His	Lys	Val	Tyr	Ala	Cys	Glu	Val	Thr	His	Gln	Gly	Leu	Ser					
210										215					220					
Ser	Pro	Val	Thr	Lys	Ser	Phe	Asn	Arg	Gly	Glu	Cys									
225										230					235					

<210> 53

<400> 53
000

```
<210> 54
<211> 461
<212> PRT
<213> Homo sapiens
```

<400> 54
Met Glu Phe Gly Leu Ser Trp Val Phe Leu Val Ala Ile Ile Lys Gly
1 5 10 15

Val Gln Cys Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys
20 25 30

Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
35 40 45

Ser Asp Tyr Tyr Met Ser Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu
50 55 60

Glu Trp Val Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala
65 70 75 80

Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
85 90 95

Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val
100 105 110

Tyr Tyr Cys Ala Arg Gly Leu Thr Gly Asp Tyr Trp Gly Gln Gly Thr
115 120 125

Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro
130 135 140

Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly

145		150		155		160
Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn						
	165			170		175
Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln						
	180		185			190
Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser						
	195		200			205
Ser Leu Gly Thr Lys Thr Tyr Thr Cys Asn Val Asp His Lys Pro Ser						
	210		215			220
Asn Thr Lys Val Asp Lys Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys						
	225		230			235
Pro Pro Cys Pro Ala Pro Glu Phe Leu Gly Gly Pro Ser Val Phe Leu						
	245			250		255
Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu						
	260			265		270
Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln						
	275		280			285
Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys						
	290		295			300
Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu						
	305		310			315
Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys						
	325			330		335
Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys						
	340			345		350
Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser						
	355			360		365
Gln Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys						
	370			375		380
Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln						
	385			390		395
Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly						
	405			410		415
Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser Arg Trp Gln						
	420			425		430
Glu Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn						
	435			440		445

His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 450 455 460

<210> 55

<400> 55
 000

<210> 56
 <211> 236
 <212> PRT
 <213> Homo sapiens

<400> 56
 Met Asp Met Arg Val Pro Ala Gln Leu Leu Gly Leu Leu Leu Trp
 1 5 10 15
 Leu Arg Gly Ala Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser
 20 25 30
 Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Pro Ser
 35 40 45
 Gln Ile Ile Ser Ser Leu Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys
 50 55 60
 Ala Pro Lys Leu Leu Ile Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val
 65 70 75 80
 Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
 85 90 95
 Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln
 100 105 110
 Ser Tyr Ser Thr Pro Phe Thr Phe Gly Pro Gly Thr Lys Val Asp Ile
 115 120 125
 Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp
 130 135 140
 Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn
 145 150 155 160
 Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu
 165 170 175
 Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp
 180 185 190
 Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr
 195 200 205
 Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser
 210 215 220

Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
 225 230 235

<210> 57

<400> 57
 000

<210> 58
 <211> 467
 <212> PRT
 <213> Homo sapiens

<400> 58
 Met Glu Leu Gly Leu Cys Trp Val Phe Leu Val Ala Ile Leu Glu Gly
 1 5 10 15
 Val Gln Cys Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln
 20 25 30
 Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
 35 40 45
 Ser Ser Phe Ser Met Thr Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
 50 55 60
 Glu Trp Val Ser Tyr Ile Ser Ser Arg Ser Ser Thr Ile Ser Tyr Ala
 65 70 75 80
 Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
 85 90 95
 Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Asp Glu Asp Thr Ala Val
 100 105 110
 Tyr Tyr Cys Ala Arg Asp Pro Leu Leu Ala Gly Ala Thr Phe Phe Asp
 115 120 125
 Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys
 130 135 140
 Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu
 145 150 155 160
 Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro
 165 170 175
 Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr
 180 185 190
 Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val
 195 200 205

Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys Thr Tyr Thr Cys Asn
 210 215 220
 Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Ser
 225 230 235 240
 Lys Tyr Gly Pro Pro Cys Pro Pro Cys Pro Ala Pro Glu Phe Leu Gly
 245 250 255
 Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met
 260 265 270
 Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser Gln
 275 280 285
 Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val
 290 295 300
 His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr
 305 310 315 320
 Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly
 325 330 335
 Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile
 340 345 350
 Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val
 355 360 365
 Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr Lys Asn Gln Val Ser
 370 375 380
 Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu
 385 390 395 400
 Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro
 405 410 415
 Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val
 420 425 430
 Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser Cys Ser Val Met
 435 440 445
 His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser
 450 455 460
 Pro Gly Lys
 465

<210> 59

<400> 59

000

<210> 60
 <211> 235
 <212> PRT
 <213> Homo sapiens

<400> 60
 Met Glu Thr Pro Ala Gln Leu Leu Phe Leu Leu Leu Trp Leu Pro
 1 5 10 15
 Asp Thr Thr Gly Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser
 20 25 30
 Leu Ser Pro Gly Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser
 35 40 45
 Val Ser Ser Ser Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala
 50 55 60
 Pro Arg Leu Leu Ile Tyr Gly Ala Ser Ser Arg Ala Thr Gly Ile Pro
 65 70 75 80
 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile
 85 90 95
 Ser Arg Leu Glu Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Tyr
 100 105 110
 Gly Ser Ser Pro Leu Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
 115 120 125
 Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Ser Ser Asp Glu
 130 135 140
 Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe
 145 150 155 160
 Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln
 165 170 175
 Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser
 180 185 190
 Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
 195 200 205
 Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser
 210 215 220
 Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
 225 230 235

<210> 61
 <400> 61

000

<210> 62

<211> 466

<212> PRT

<213> Homo sapiens

<400> 62

Met Glu Leu Gly Leu Cys Trp Val Phe Leu Val Ala Ile Leu Glu Gly
 1 5 10 15

Val Gln Cys Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln
 20 25 30

Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
 35 40 45

Ser Ser Phe Ser Met Thr Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
 50 55 60

Glu Trp Val Ser Tyr Ile Ser Ser Arg Ser Ser Thr Ile Ser Tyr Ala
 65 70 75 80

Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
 85 90 95

Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Asp Glu Asp Thr Ala Val
 100 105 110

Tyr Tyr Cys Ala Arg Asp Pro Leu Leu Ala Gly Ala Thr Phe Asp
 115 120 125

Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys
 130 135 140

Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu
 145 150 155 160

Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro
 165 170 175

Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr
 180 185 190

Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val
 195 200 205

Val Thr Val Pro Ser Ser Asn Phe Gly Thr Gln Thr Tyr Thr Cys Asn
 210 215 220

Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys Thr Val Glu Arg
 225 230 235 240

Lys Cys Cys Val Glu Cys Pro Pro Cys Pro Ala Pro Pro Val Ala Gly
 245 250 255

Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
 260 265 270
 Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
 275 280 285
 Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
 290 295 300
 Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Phe Arg
 305 310 315 320
 Val Val Ser Val Leu Thr Val Val His Gln Asp Trp Leu Asn Gly Lys
 325 330 335
 Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ala Pro Ile Glu
 340 345 350
 Lys Thr Ile Ser Lys Thr Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
 355 360 365
 Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
 370 375 380
 Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
 385 390 395 400
 Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Met
 405 410 415
 Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
 420 425 430
 Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
 435 440 445
 Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
 450 455 460
 Gly Lys
 465

<210> 63

 <400> 63
 000

<210> 64

 <400> 64
 000

<210> 65

<400> 65
000

<210> 66
<211> 461
<212> PRT
<213> Homo sapiens

<400> 66
Met Glu Phe Gly Leu Ser Trp Val Phe Leu Val Ala Ile Ile Lys Gly
1 5 10 15

Val Gln Cys Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys
20 25 30

Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
35 40 45

Ser Asp Tyr Tyr Met Ser Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu
50 55 60

Glu Trp Val Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala
65 70 75 80

Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
85 90 95

Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val
100 105 110

Tyr Tyr Cys Ala Ile Arg Ile Gly Gly Met Asp Val Trp Gly Gln Gly
115 120 125

Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe
130 135 140

Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu
145 150 155 160

Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp
165 170 175

Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu
180 185 190

Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser
195 200 205

Ser Asn Phe Gly Thr Gln Thr Tyr Thr Cys Asn Val Asp His Lys Pro
210 215 220

Ser Asn Thr Lys Val Asp Lys Thr Val Glu Arg Lys Cys Cys Val Glu
225 230 235 240

Cys	Pro	Pro	Cys	Pro	Ala	Pro	Pro	Val	Ala	Gly	Pro	Ser	Val	Phe	Leu
			245						250					255	
Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu
			260					265					270		
Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Gln
			275				280					285			
Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys
			290			295					300				
Pro	Arg	Glu	Glu	Gln	Phe	Asn	Ser	Thr	Phe	Arg	Val	Val	Ser	Val	Leu
			305			310				315					320
Thr	Val	Val	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys
				325					330					335	
Val	Ser	Asn	Lys	Gly	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys
			340					345					350		
Thr	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser
			355				360					365			
Arg	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys
			370			375					380				
Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln
					390					395					400
Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Met	Leu	Asp	Ser	Asp	Gly
				405					410					415	
Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln
			420					425					430		
Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn
			435				440					445			
His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	Lys			
			450			455					460				

<210> 67

<400> 67
000

<210> 68

<400> 68
000

<210> 69

<400> 69
000

<210> 70
<211> 461
<212> PRT
<213> Homo sapiens

<400> 70
Met Glu Phe Gly Leu Ser Trp Val Phe Leu Val Ala Ile Ile Lys Gly
1 5 10 15
Val Gln Cys Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys
20 25 30
Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
35 40 45
Ser Asp Tyr Tyr Met Ser Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu
50 55 60
Glu Trp Val Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala
65 70 75 80
Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
85 90 95
Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val
100 105 110
Tyr Tyr Cys Ala Arg Ile Gly Gly Met Asp Val Trp Gly Gln Gly Thr
115 120 125
Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro
130 135 140
Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly
145 150 155 160
Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn
165 170 175
Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln
180 185 190
Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser
195 200 205
Ser Leu Gly Thr Lys Thr Tyr Thr Cys Asn Val Asp His Lys Pro Ser
210 215 220
Asn Thr Lys Val Asp Lys Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys
225 230 235 240
Pro Ser Cys Pro Ala Pro Glu Phe Leu Gly Gly Pro Ser Val Phe Leu
245 250 255

Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu
 260 265 270
 Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln
 275 280 285
 Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys
 290 295 300
 Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu
 305 310 315 320
 Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys
 325 330 335
 Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys
 340 345 350
 Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser
 355 360 365
 Gln Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys
 370 375 380
 Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln
 385 390 395 400
 Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly
 405 410 415
 Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser Arg Trp Gln
 420 425 430
 Glu Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn
 435 440 445
 His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 450 455 460

<210> 71

 <400> 71
 000

<210> 72

 <400> 72
 000

<210> 73

<400> 73
000

<210> 74
<211> 460
<212> PRT
<213> Homo sapiens

<400> 74
Met Glu Phe Gly Leu Ser Trp Val Phe Leu Val Ala Ile Ile Lys Gly
1 5 10 15
Val Gln Cys Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys
20 25 30
Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
35 40 45
Ser Asp Tyr Tyr Met Ser Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu
50 55 60
Glu Trp Val Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala
65 70 75 80
Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
85 90 95
Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val
100 105 110
Tyr Tyr Cys Ala Arg Gly Leu Thr Gly Asp Tyr Trp Gly Gln Gly Thr
115 120 125
Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro
130 135 140
Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly
145 150 155 160
Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn
165 170 175
Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln
180 185 190
Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser
195 200 205
Asn Phe Gly Thr Gln Thr Tyr Thr Cys Asn Val Asp His Lys Pro Ser
210 215 220
Asn Thr Lys Val Asp Lys Thr Val Glu Arg Lys Cys Cys Val Glu Cys
225 230 235 240
Pro Pro Cys Pro Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu Phe
245 250 255

Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val
 260 265 270
 Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Gln Phe
 275 280 285
 Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro
 290 295 300
 Arg Glu Glu Gln Phe Asn Ser Thr Phe Arg Val Val Ser Val Leu Thr
 305 310 315 320
 Val Val His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val
 325 330 335
 Ser Asn Lys Gly Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr
 340 345 350
 Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg
 355 360 365
 Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly
 370 375 380
 Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro
 385 390 395 400
 Glu Asn Asn Tyr Lys Thr Thr Pro Pro Met Leu Asp Ser Asp Gly Ser
 405 410 415
 Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln
 420 425 430
 Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His
 435 440 445
 Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 450 455 460

<210> 75

<400> 75

000

<210> 76

<400> 76

000

<210> 77

<400> 77

000

<210> 78
 <211> 461
 <212> FRT
 <213> Homo sapiens

<400> 78
 Met Glu Phe Gly Leu Ser Trp Val Phe Leu Val Ala Ile Ile Lys Gly
 1 5 10 15
 Val Gln Cys Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys
 20 25 30
 Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
 35 40 45
 Ser Asp Tyr Tyr Met Ser Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu
 50 55 60
 Glu Trp Val Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala
 65 70 75 80
 Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
 85 90 95
 Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val
 100 105 110
 Tyr Tyr Cys Ala Arg Gly Leu Thr Gly Asp Tyr Trp Gly Gln Gly Thr
 115 120 125
 Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro
 130 135 140
 Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly
 145 150 155 160
 Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn
 165 170 175
 Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln
 180 185 190
 Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser
 195 200 205
 Ser Leu Gly Thr Lys Thr Tyr Thr Cys Asn Val Asp His Lys Pro Ser
 210 215 220
 Asn Thr Lys Val Asp Lys Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys
 225 230 235 240
 Pro Ser Cys Pro Ala Pro Glu Phe Leu Gly Gly Pro Ser Val Phe Leu
 245 250 255
 Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu

260	265	270
Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln		
275	280	285
Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys		
290	295	300
Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu		
305	310	315
Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys		
325	330	335
Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys		
340	345	350
Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser		
355	360	365
Gln Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys		
370	375	380
Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln		
385	390	395
Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly		
405	410	415
Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser Arg Trp Gln		
420	425	430
Glu Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn		
435	440	445
His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys		
450	455	460

<210> 79

<400> 79

000

<210> 80

<400> 80

000

<210> 81

<400> 81

000

<210> 82

<211> 461

<212> PRT

<213> Homo sapiens

<400> 82

Met Glu Phe Gly Leu Ser Trp Val Phe Leu Val Ala Ile Ile Lys Gly
 1 5 10 15

Val Gln Cys Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys
 20 25 30

Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
 35 40 45

Ser Asp Tyr Tyr Met Ser Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu
 50 55 60

Glu Trp Val Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala
 65 70 75 80

Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
 85 90 95

Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val
 100 105 110

Tyr Tyr Cys Ala Arg Gly Leu Thr Gly Asp Tyr Trp Gly Gln Gly Thr
 115 120 125

Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro
 130 135 140

Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly
 145 150 155 160

Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn
 165 170 175

Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln
 180 185 190

Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser
 195 200 205

Ser Leu Gly Thr Lys Thr Tyr Thr Cys Asn Val Asp His Lys Pro Ser
 210 215 220

Asn Thr Lys Val Asp Lys Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys
 225 230 235 240

Pro Pro Cys Pro Ala Pro Glu Phe Leu Gly Gly Pro Ser Val Phe Leu
 245 250 255

Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu
 260 265 270

Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln
 275 280 285
 Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys
 290 295 300
 Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu
 305 310 315 320
 Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys
 325 330 335
 Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys
 340 345 350
 Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser
 355 360 365
 Gln Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys
 370 375 380
 Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln
 385 390 395 400
 Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly
 405 410 415
 Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser Arg Trp Gln
 420 425 430
 Glu Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn
 435 440 445
 His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 450 455 460

<210> 83

 <400> 83
 000

<210> 84

 <400> 84
 000

<210> 85

 <400> 85
 000

<210> 86

<211> 462

<212> PRT

<213> Homo sapiens

<400> 86

```

Met Glu Phe Gly Leu Ser Trp Val Phe Leu Val Ala Ile Ile Lys Gly
 1           5           10           15

Val Gln Cys Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys
           20           25           30

Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
           35           40           45

Ser Asp Tyr Tyr Met Ser Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu
 50           55           60

Glu Trp Val Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala
 65           70           75           80

Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
           85           90           95

Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val
           100          105          110

Tyr Tyr Cys Ala Arg Arg Ile Gly Gly Met Asp Val Trp Gly Gln Gly
           115          120          125

Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe
           130          135          140

Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu
           145          150          155          160

Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp
           165          170          175

Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu
           180          185          190

Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser
           195          200          205

Ser Ser Leu Gly Thr Lys Thr Tyr Thr Cys Asn Val Asp His Lys Pro
           210          215          220

Ser Asn Thr Lys Val Asp Lys Arg Val Glu Ser Lys Tyr Gly Pro Pro
           225          230          235          240

Cys Pro Pro Cys Pro Ala Pro Glu Phe Leu Gly Gly Pro Ser Val Phe
           245          250          255

Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro
           260          265          270

Glu Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val

```

275	280	285
Gln Phe Asn Trp Tyr Val Asp	Gly Val Glu Val His Asn Ala Lys Thr	
290	295	300
Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val		
305	310	315 320
Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys		
325	330	335
Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser		
340	345	350
Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro		
355	360	365
Ser Gln Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val		
370	375	380
Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly		
385	390	395 400
Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp		
405	410	415
Gly Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser Arg Trp		
420	425	430
Gln Glu Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His		
435	440	445
Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys		
450	455	460

<210> 87

<400> 87
000

<210> 88

<400> 88
000

<210> 89

<400> 89
000

<210> 90

<211> 467

<212> PRT

<213> Homo sapiens

<400> 90

Met Glu Leu Gly Leu Cys Trp Val Phe Leu Val Ala Ile Leu Glu Gly
 1 5 10 15

Val Gln Cys Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln
 20 25 30

Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
 35 40 45

Ser Ser Phe Ser Met Thr Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
 50 55 60

Glu Trp Val Ser Tyr Ile Ser Ser Arg Ser Ser Thr Ile Ser Tyr Ala
 65 70 75 80

Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
 85 90 95

Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Asp Glu Asp Thr Ala Val
 100 105 110

Tyr Tyr Cys Ala Arg Asp Pro Leu Leu Ala Gly Ala Thr Phe Phe Asp
 115 120 125

Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys
 130 135 140

Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu
 145 150 155 160

Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro
 165 170 175

Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr
 180 185 190

Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val
 195 200 205

Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys Thr Tyr Thr Cys Asn
 210 215 220

Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Ser
 225 230 235 240

Lys Tyr Gly Pro Pro Cys Pro Pro Cys Pro Ala Pro Glu Phe Leu Gly
 245 250 255

Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met
 260 265 270

Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser Gln
 275 280 285
 Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val
 290 295 300
 His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr
 305 310 315 320
 Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly
 325 330 335
 Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile
 340 345 350
 Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val
 355 360 365
 Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr Lys Asn Gln Val Ser
 370 375 380
 Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu
 385 390 395 400
 Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro
 405 410 415
 Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val
 420 425 430
 Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser Cys Ser Val Met
 435 440 445
 His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser
 450 455 460
 Pro Gly Lys
 465

<210> 91

 <400> 91
 000

<210> 92

 <400> 92
 000

<210> 93

 <400> 93
 000

<210> 94
 <211> 467
 <212> PRT
 <213> Homo sapiens

<400> 94
 Met Glu Leu Gly Leu Cys Trp Val Phe Leu Val Ala Ile Leu Glu Gly
 1 5 10 15
 Val Gln Cys Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln
 20 25 30
 Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
 35 40 45
 Ser Ser Phe Ser Met Thr Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
 50 55 60
 Glu Trp Val Ser Tyr Ile Ser Ser Arg Ser Ser Thr Ile Ser Tyr Ala
 65 70 75 80
 Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
 85 90 95
 Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Asp Glu Asp Thr Ala Val
 100 105 110
 Tyr Tyr Cys Ala Arg Asp Pro Leu Leu Ala Gly Ala Thr Phe Phe Asp
 115 120 125
 Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys
 130 135 140
 Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu
 145 150 155 160
 Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro
 165 170 175
 Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr
 180 185 190
 Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val
 195 200 205
 Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys Thr Tyr Thr Cys Asn
 210 215 220
 Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Ser
 225 230 235 240
 Lys Tyr Gly Pro Pro Cys Pro Ser Cys Pro Ala Pro Glu Phe Leu Gly
 245 250 255
 Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met

260 265 270
 Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser Gln
 275 280 285
 Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val
 290 295 300
 His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr
 305 310 315 320
 Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly
 325 330 335
 Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile
 340 345 350
 Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val
 355 360 365

 Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr Lys Asn Gln Val Ser
 370 375 380
 Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu
 385 390 395 400
 Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro
 405 410 415
 Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val
 420 425 430
 Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser Cys Ser Val Met
 435 440 445
 His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser
 450 455 460
 Pro Gly Lys
 465

<210> 95

 <400> 95
 000

<210> 96

 <400> 96
 000

<210> 97

<211> 1413

<212> DNA

<213> Homo sapiens

<400> 97

```

atggagttgg ggctgagctg ggttttccct gttgctatta taaaaggtgt ccagtgtag 60
gtgcagctgg tggagttctg gggaggcttg gtacagcctg ggggggtccct gagactctcc 120
tgtgcagcct ctggattcac cttcagtagt tttagtatga cctgggtccg ccaggctcca 180
ggaaaggggc tggagttgggt ttcatatcatt agtagtagaa gtagtacat atcctacgca 240
gaactctgtga agggcccgatt caccatctcc agagacaatg ccaagaactc actgtatctg 300
caaatgaaca gctctgagaga ctagcgaggag ctacctctct tgcactctgg ggcaggggaa ccttggtcac cgtctctcca 420
ctagcgaggag agggcccatc ggtcttcccc ctggcacctc cctccaaagag caactctggg 480
ggcacagcgg cctctggcgtg cctgggtcaag gactactctc cgaacccggt gacggtgtcg 540
tggaactcag ggcacctgac cagcggcgctg cacacctctc cggctgtctc acagtctcca 600
ggactctact cctcagcag cgtgggtgacc gtgccctcca gcagcttggg caccagacc 660
tacctctgca acgtgaatca caagcccagc aacaccaagg tggacaagaa agttgagccc 720
aaatcttgtg acaaaaactca cacatgccca cctgcccag cactgaact cctgggggga 780
ccgtcagctc tctctctccc cccaaaaccc aaggacaccc tcatgatctc ccggaccctc 840
gaggtcacat gcgtgggtgg ggcagtgagc cagcaagacc ctgaggtcaa gttcaactgg 900
tacgtggagc gcgtggaggt gcataatgcc aagacaaagc cgcgggagga gcagtacaac 960
agcacgtacc gttgtgctag cgtcctcacc gtcctgcacc aggaactggct gaattggcaa 1020
gagtacaagt gccaggtctc caacaaagcc ctcccagccc ccactcgagaa aacctctccc 1080
aaagccaaag ggcagccccg agaaccacag gtgtacaccc tgcccctacc ccgggatgag 1140
ctgaccaaga accaggtcag cctgacctgc cttggtcaaag gcttctatcc cagcgacatc 1200
ccctgtggagt gggagagcaa tgggcagccg gagaacaaat acaagaccac gctctccctg 1260
ctggactccg acggctcctt cttctctcac agcaagctca cctgggacaa gacgaggtgg 1320
cagcagggga acgtctcttc atgctccgtg atgcattgag ctctgcacaa ccactacacg 1380
cagaagagcc tctccctgtc tccgggtaaa tag 1413

```

<210> 98

<211> 470

<212> PRT

<213> Homo sapiens

<400> 98

```

Met Glu Leu Gly Leu Cys Trp Val Phe Leu Val Ala Ile Leu Glu Gly
  1             5             10            15

Val Gln Cys Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln
    20             25            30

Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
    35             40            45

Ser Ser Phe Ser Met Thr Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
    50             55            60

Glu Trp Val Ser Tyr Ile Ser Ser Arg Ser Ser Thr Ile Ser Tyr Ala
    65             70            75            80

Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
    85             90            95

Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Asp Glu Asp Thr Ala Val
   100           105           110

```

Tyr Tyr Cys Ala Arg Asp Pro Leu Leu Ala Gly Ala Thr Phe Phe Asp
 115 120 125
 Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys
 130 135 140
 Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly
 145 150 155 160
 Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro
 165 170 175
 Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr
 180 185 190
 Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val
 195 200 205
 Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn
 210 215 220
 Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro
 225 230 235 240
 Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu
 245 250 255
 Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp
 260 265 270
 Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp
 275 280 285
 Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly
 290 295 300
 Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn
 305 310 315 320
 Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp
 325 330 335
 Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro
 340 345 350
 Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu
 355 360 365
 Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn
 370 375 380
 Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile
 385 390 395 400
 Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr
 405 410 415

Thr Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys
 420 425 430

Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys
 435 440 445

Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu
 450 455 460

Ser Leu Ser Pro Gly Lys
 465 470

<210> 99

<400> 99

000

<210> 100

<400> 100

000

<210> 101

<211> 1395

<212> DNA

<213> Homo sapiens

<400> 101

atggagtttg ggcgtgagctg ggttttccct gtgtctatta taaaagggtt ccagtgctcag 60
 gtgcagctgg tggagtcctgg gggaggcttg gtcaagcctg gagggctcct gagactctcc 120
 tgtgcagcct ctggtattcac ctccagtgac tactatatga gctggatccg ccaggctcca 180
 ggggaaggac tggagtggtt ttcatacatt agtagtagtg gttagtaccat atactacga 240
 gactctgtga agggccgatt caccatctcc agggacaacg ccaagaactc actgtatctg 300
 caaatgaaca gcctgagagc cgaggacacg gccgtgtatt actgtgcgag aggcctaact 360
 ggggactact gggggcaggg aacctgtgtc accgtctcct cagcttccac caagggccca 420
 tcggtcttcc cctctggcacc ctctcccaag agcacctctg ggggcacacg ggcctctggg 480
 tgccctggta aggactactt ccccgaaacg gtgacgggtg cgtgggaactc aggcgcctg 540
 accagcggcg tgcacacctt cccggctgtc ctacagctct caggactcta ctccctcage 600
 agcgtgtgta ccgtgccttc cagcagcttg ggcacccaga cctacatctg caacgtgaat 660
 cacaagccca gcaacaccaa ggtggacaag aaagttaggc ccaaatcttg tgacaaaact 720
 cacacatgcc caccgtgcc agcacctgaa ctctggggg gaccgtcagt ctctccttc 780

cccccaaac ccaaggacac cctcatgac tcccgaccc ctgaggtcac atgcgtgggt 840
 gtggagctga gccacgaaga cctcaggtgc aagttcaact ggtacgtgga cggcgtggag 900
 gtgcataatg ccaagacaaa gcccgggag gagcagtaca acagcacgta ccgtgtgttc 960
 agcgtcctca ccgtcctgca ccaggactgg ctgaatggca aggagtacaa gtgcaaggtc 1020
 tccaacaaag cctctccagc ccccatcgag aaacacctct ccaagccaa agggcagccc 1080
 cgagaaccaa aggtgttacac cctgcoccca tcccgggatg agctgaccaa gaaccaagtc 1140
 agcctgacct gcctgtgtcaa aggtctctat cccagcgaca tcgcccgtga gtgggagagc 1200
 aatgggcagc cgagaaacaa ctacaagacc agcctcccgt tgcctggact cgacggctcc 1260
 ttcttctctc acagcaagct caccgtggac aagagcaggt ggcagcaggg gaacgtcttc 1320
 tcattgctccg tgatgcataa ggctctgcac aaccactaca cgcagaagag cctctccctg 1380
 ttctccggta aatag 1395

<210> 102
 <211> 464
 <212> PRT
 <213> Homo sapiens

<400> 102

Met Glu Phe Gly Leu Ser Trp Val Phe Leu Val Ala Ile Ile Lys Gly
 1 5 10 15

Val Gln Cys Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys
 20 25 30

Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
 35 40 45

Ser Asp Tyr Tyr Met Ser Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu
 50 55 60

Glu Trp Val Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala
 65 70 75 80

Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
 85 90 95

Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val
 100 105 110

Tyr Tyr Cys Ala Arg Gly Leu Thr Gly Asp Tyr Trp Gly Gln Gly Thr
 115 120 125

Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro
 130 135 140

Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly
 145 150 155 160

Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn
 165 170 175

Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln
 180 185 190

Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser
 195 200 205

Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser
 210 215 220

Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys Thr
 225 230 235 240

His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser
 245 250 255

Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg

260										265					270				
Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro				
		275					280				285								
Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala				
		290			295						300								
Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val				
		305			310				315						320				
Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr				
				325				330						335					
Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr				
		340						345				350							
Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu				
		355				360						365							
Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys				
		370				375						380							
Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser				
		385				390				395				400					
Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp				
				405				410						415					
Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser				
		420						425				430							
Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala				
		435				440						445							
Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	Lys				
		450				455						460							

```
<210> 103
<211> 107
<212> PRT
<213> Homo sapiens
```

```
<400> 103
Asp Ile Gln Met Thr Gln Ser Pro Ser Leu Ser Ala Ser Val Gly
   1                               10                      15
                        5
Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Ser Ile Ser Ser Tyr
                20                    25                  30
Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
    35              40              45
```

Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80
 Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ser Tyr Ser Thr Pro Phe
 85 90 95
 Thr Phe Gly Pro Gly Thr Lys Val Asp Ile Lys
 100 105

<210> 104
 <211> 120
 <212> PRT
 <213> Homo sapiens

<400> 104
 Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30
 Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ser Ala Ile Ser Ser Gly Ser Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Lys Tyr Ser Gly Ser Tyr Tyr Tyr Phe Asp Tyr Trp Gly Gln Gly
 100 105 110
 Thr Leu Val Thr Val Ser Ser Ala
 115 120

<210> 105
 <211> 119
 <212> PRT
 <213> Homo sapiens

<400> 105
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ala Asn Ile Lys Gln Asp Gly Ser Glu Lys Tyr Tyr Val Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80
 Leu Gln Met Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala
 85 90 95
 Arg Gly Ile Ala Ala Ala Gly Tyr Phe Asp Tyr Trp Gly Gln Gly Thr
 100 105 110
 Leu Val Thr Val Ser Ser Ala
 115

<210> 106
 <211> 117
 <212> PRT
 <213> Homo sapiens

<400> 106
 Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Tyr
 20 25 30
 Tyr Met Ser Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Ala Leu Gly Gly Met Asp Val Trp Gly Gln Gly Thr Thr Val
 100 105 110
 Thr Val Ser Ser Ala
 115

<210> 107
 <211> 107
 <212> PRT
 <213> Homo sapiens

<400> 107

Glu Ile Val Met Thr Gln Ser Pro Ala Thr Leu Ser Val Ser Pro Gly
 1 5 10 15
 Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Asn
 20 25 30
 Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
 35 40 45
 Tyr Gly Ala Ser Thr Arg Ala Thr Gly Ile Pro Ala Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Ser
 65 70 75 80
 Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Tyr Asn Asn Trp Pro Phe
 85 90 95
 Thr Phe Gly Pro Gly Thr Lys Val Asp Ile Lys
 100 105

<210> 108
 <211> 115
 <212> PRT
 <213> Homo sapiens

<400> 108
 Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Tyr
 20 25 30
 Tyr Met Ser Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Leu Thr Gly Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val
 100 105 110
 Ser Ser Ala
 115

<210> 109
 <211> 108
 <212> PRT

<213> Homo sapiens

<400> 109

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Val Ser Ala Ser Val Gly
 1 5 10 15
 Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ser Ser Trp
 20 25 30
 Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
 35 40 45
 Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80
 Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ala Asn Ser Phe Pro Phe
 85 90 95
 Thr Phe Gly Pro Gly Thr Lys Val Asp Ile Lys Arg
 100 105

<210> 110

<211> 117

<212> PRT

<213> Homo sapiens

<400> 110

Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30
 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ala Val Ile Trp Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Gly Tyr Ser Tyr Phe Asp Tyr Trp Gly Gln Gly Thr Leu Val
 100 105 110
 Thr Val Ser Ser Ala
 115

<210> 111
 <211> 118
 <212> PRT
 <213> Homo sapiens

<400> 111
 Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
 20 25 30
 Gly Ile Ser Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45
 Gly Trp Ile Ser Ala Tyr Asn Gly Asn Thr Asn Tyr Ala Gln Lys Leu
 50 55 60
 Gln Gly Arg Val Thr Met Thr Thr Asp Thr Ser Thr Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Arg Ser Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Asp Tyr Gly Gly Asn Tyr Phe Asp Tyr Trp Gly Gln Gly Thr Leu
 100 105 110
 Val Thr Val Ser Ser Ala
 115

<210> 112
 <211> 114
 <212> PRT
 <213> Homo sapiens

<400> 112
 Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
 1 5 10 15
 Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Gln Ser Val Leu Tyr Ser
 20 25 30
 Ser Asn Asn Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
 35 40 45
 Pro Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
 50 55 60
 Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
 65 70 75 80
 Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln
 85 90 95
 Tyr Tyr Ser Thr Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile

100 105 110

Lys Arg

<210> 113
 <211> 116
 <212> PRT
 <213> Homo sapiens

<400> 113
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30
 Ser Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ser Tyr Ile Ser Ser Ser Ser Ser Thr Ile Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Asp Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Ile Val Gly Phe Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr
 100 105 110
 Val Ser Ser Ala
 115

<210> 114
 <211> 109
 <212> PRT
 <213> Homo sapiens

<400> 114
 Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Leu Ser Pro Gly
 1 5 10 15
 Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser
 20 25 30
 Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu
 35 40 45
 Ile Tyr Gly Ala Ser Ser Arg Ala Thr Gly Ile Pro Asp Arg Phe Ser
 50 55 60
 Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg Leu Glu

65					70					75					80
Pro	Glu	Asp	Phe	Ala	Val	Tyr	Tyr	Cys	Gln	Gln	Tyr	Gly	Ser	Ser	Pro
				85					90					95	
Leu	Thr	Phe	Gly	Gly	Thr	Lys	Val	Glu	Ile	Lys	Arg				
			100				105								

```
<210> 115
<211> 114
<212> PRT
<213> Homo sapiens
```

```

400> 115
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
  1              5              10              15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Tyr
      20              25              30

Tyr Met Ser Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
      35              40              45

Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser Val
      50              55              60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
      65              70              75              80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
      85              90              95

Ala Ile Gly Met Asp Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser
      100             105             110

Ser Ala

```

```
<210> 116
<211> 115
<212> PRT
<213> Homo sapiens
```

<400> 116
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Tyr
20 25 30
Tyr Met Ser Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45
Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg Leu Thr Gly Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val
 100 105 110

Ser Ser Ala
 115

<210> 117

<211> 108

<212> PRT

<213> Homo sapiens

<400> 117

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Val Ser Ala Ser Val Gly
 1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ser Ser Trp
 20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
 35 40 45

Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ala Asn Ser Phe Pro Leu
 85 90 95

Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg
 100 105

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
7 April 2005 (07.04.2005)

PCT

(10) International Publication Number
WO 2005/030124 A3(51) International Patent Classification: ⁷ C07K 16/00,
C12N 15/00HAAK-FRENDSCHO, Mary [US/US]; 8472 Willow
Place, Newark, California 94560 (US).(21) International Application Number:
PCT/US2004/029390(74) Agents: HALEY, James F., Jr. et al.; c/o FISH &
NEAVE, 1251 Avenue of the Americas, New York, NY
10020 (US).(22) International Filing Date:
9 September 2004 (09.09.2004)(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/502,163 10 September 2003 (10.09.2003) US(71) Applicants (for all designated States except US):
WARNER-LAMBERT COMPANY LLC [US/US];
201 Tabor Road, Morris Plains, New Jersey 07950 (US).
ABGENIX, INC. [US/US]; 6701 Kaiser Drive, Fremont,
California 94555 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BEDIAN, Vahe
[US/US]; 3 Frog Hollow Road, East Lyme, Connecticut
06333 (US). DEVALARAJA, Madhav Narasimha
[IN/US]; 690 Trego Circle, Ann Arbor, Michigan 48103
(US). LOW, Joseph Edwin [US/US]; 4259 Fairway
Drive, Ann Arbor, Michigan 48103 (US). MOBLEY,
James Leslie [US/US]; 10195 Covington, Brighton,
Michigan 48114 (US). KELLERMANN, Srid-Aimee
[DE/US]; 1000 Alamanor Avenue, Menlo Park, California
94025 (US). FOLTZ, Ian [CA/CA]; 2108 Knightwood
Place, Burnaby, British Columbia V5A4B9 (CA).(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE,
SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN,
GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:
30 March 2006For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: ANTIBODIES TO M-CSF

(57) Abstract: The present invention relates to antibodies and antigen-binding portions thereof that specifically bind to a M-CSF, preferably human M-CSF, and that function to inhibit a M-CSF. The invention also relates to human anti-M-CSF antibodies and antigen-binding portions thereof. The invention also relates to antibodies that are chimeric, bispecific, derivatized, single chain antibodies or portions of fusion proteins. The invention also relates to isolated heavy and light chain immunoglobulins derived from human anti-M-CSF antibodies and nucleic acid molecules encoding such immunoglobulins. The present invention also relates to methods of making human anti-M-CSF antibodies, compositions comprising these antibodies and methods of using the antibodies and compositions for diagnosis and treatment. The invention also provides gene therapy methods using nucleic acid molecules encoding the heavy and/or light immunoglobulin molecules that comprise the human anti-M-CSF antibodies. The invention also relates to transgenic animals and transgenic plants comprising nucleic acid molecules of the present invention.

INTERNATIONAL SEARCH REPORT

PCT/US04/29390

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C07K 16/00; C12N 15/00

US CL : 530/387.3, 388.1, 388.15, 388.23; 800/6

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 530/387.3, 388.1, 388.15, 388.23; 800/6

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
Please See Continuation Sheet

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
<input checked="" type="checkbox"/> X	US 5,491,065 (HALENBECK et al) 13 February 1996 (13.02.1996), columns 6-7.	1
<input type="checkbox"/> Y		2, 34
<input checked="" type="checkbox"/> X	JP 05-095794 A (OMOTO et al) 20 April 1993 (20.04.1993), see entire document, especially page 2, paragraph [0009] of the attached computer translation.	1-2
<input type="checkbox"/> Y		34
<input checked="" type="checkbox"/> 34	WO 91/10741 (ROWLAND et al) 25 July 1991 (25.07.1991), see entire document, especially pages 3-15 and page 2, lines 25-38)	Y

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:	"T"
"A" document defining the general state of the art which is not considered to be of particular relevance	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or the theory underlying the invention
"B" earlier application or patent published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"C" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

14 October 2005 (14.10.2005)

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US
Commissioner for Patents
P.O. Box 1450
Alexandria, Virginia 22313-1450

Facsimile No. (703) 305-3230

Date of mailing of the international search report

06 JAN 2006

Authorized officer

David J. Blanchard

Telephone No. (571) 272-1600

INTERNATIONAL SEARCH REPORT

International appl.
No. 15
PCT/US04/29900

Continuation of B. FIELDS SEARCHED Item 3:

WEST, Medline, Embase, Cancerlit, Biosis, Biotechno, Geneseq, issued and published patents, PIR 79, Uniprot 03.

Search terms: SEQ ID numbers, M-CSF, c-fms, human monoclonal, xenogenic antibody, non-human transgenic, inventor search.